

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/12, C07K 14/47, A01K 67/027, C07K 16/18, C12Q 1/68, G06F 17/30		A1	(11) International Publication Number: WO 00/63375 (43) International Publication Date: 26 October 2000 (26.10.00)
(21) International Application Number: PCT/IB00/00562 (22) International Filing Date: 20 April 2000 (20.04.00) (30) Priority Data: 60/130,217 20 April 1999 (20.04.99) US (71) Applicant (for all designated States except US): GENSET [FR/FR]; Intellectual Property Dept., 24, rue Royale, F-75008 Paris (FR). (72) Inventors; and (75) Inventors/Applicants (for US only): BOUGUELERET, Lydie [FR/FR]; 108, avenue Victor Hugo, F-92170 Vanves (FR). DUFAURE-GARE, Isabelle [FR/FR]; 1 bis, rue d'Alesia, F-75014 Paris (FR). GREL, Pascale [FR/FR]; 45, boulevard Alsace-Lorraine, F-91170 Viry-Chatillon (FR). (74) Common Representative: GENSET; Intellectual Property Dept., 24, rue Royale, F-75008 Paris (FR).		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>	
(54) Title: DNA ENCODING A KINESIN-LIKE PROTEIN (HKLP) COMPRISING BIALLELIC MARKERS			
(57) Abstract The present invention is directed to polynucleotides encoding a human kinesin-like polypeptide as well as a regulatory region located at the 3'-end of said coding region. The invention also concerns polypeptides encoded by the kinesin-like gene. The invention also deals with antibodies directed specifically against such polypeptides that are useful as diagnostic reagents. The invention further encompasses biallelic markers of the <i>HKLP</i> gene useful in genetic analysis.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

DNA ENCODING A KINESIN-LIKE PROTEIN (HKLP) COMPRISING BIALLELIC MARKERS

FIELD OF THE INVENTION

The present invention is directed to polynucleotides encoding a human kinesin-like polypeptide as well as a regulatory region located at the 3'-end of said coding region. The invention
5 also concerns polypeptides encoded by the kinesin-like gene. The invention also deals with antibodies directed specifically against such polypeptides that are useful as diagnostic reagents. The invention further encompasses biallelic markers of the *HKLP* gene useful in genetic analysis.

BACKGROUND OF THE INVENTION

The kinesins are mechanochemical proteins utilizing chemical energy from ATP hydrolysis
10 to generate mechanical force. The kinesins can bind to and move on microtubules in the presence of ATP. The ability to move on microtubules has led to the classification of kinesins as microtubule motor proteins. The kinesins play important roles in intracellular transport and cell division.

Several kinesin proteins are involved in vesicle/organelle transport in neurons, and mutants of kinesin in *Drosophila* show impaired neuronal function. In humans, defects in kinesin-encoding
15 genes could cause neurological disorders or syndromes of clinical importance.

The kinesin proteins carry out or facilitate movements of the chromosomes and spindle in meiosis and mitosis. Defective meiotic kinesins in humans may be the causes of infertility, spontaneous abortion, neonatal chromosome disorders, and aneuploidy. In mitotically dividing cells, mutations in kinesin proteins could cause somatic abnormalities or cellular transformation, including
20 neoplasia.

Finally, the kinesins could be involved in developmental processes as the localization of some morphogens has been shown to be microtubule-dependent.

The KIF kinesin superfamily proteins have been identified as candidate motor proteins involved in organelle transport.

25 Among the KIFs, the murine KIF1A protein has been proposed as a transporter of synaptic vesicle precursors. KIF1A disruption assays in mice allowed to show that KIF1A is involved in the transport of a synaptic vesicle precursor and that KIF1A-mediated axonal transport plays a critical role in viability, maintenance, and function of neurons, particularly mature neurons (Yonekawa et al., 1998). The murine KIF1B protein is co-localized with mitochondria *in vivo* and could be
30 involved in the transport of mitochondria (Nangaku et al., 1994).

SUMMARY OF THE INVENTION

The present invention pertains to nucleic acid molecules comprising the genomic sequence of a novel human gene which encodes a kinesin-like protein and which has been named *HKLP* by the inventors. The *HKLP* presents homology with murine *KIF1A* and *KIF1B*. The *HKLP* genomic

sequence comprises regulatory sequence located downstream (3'-end) of the transcribed portion of said gene, these regulatory sequences being also part of the invention.

The invention also deals with the complete cDNA sequence encoding the HKLP protein, as well as with the corresponding translation product.

5 Oligonucleotide probes or primers hybridizing specifically with a *HKLP* genomic or cDNA sequence are also part of the present invention, as well as DNA amplification and detection methods using said primers and probes.

A further object of the invention consists of recombinant vectors comprising any of the nucleic acid sequences described above, and in particular of recombinant vectors comprising a
10 *HKLP* regulatory sequence or a sequence encoding a HKLP protein, as well as of cell hosts and transgenic non human animals comprising said nucleic acid sequences or recombinant vectors.

The invention is also directed to biallelic markers that are located within the *HKLP* genomic sequence or that are in linkage disequilibrium with the *HKLP* gene, these biallelic markers representing useful tools in order to identify a statistically significant association between specific
15 alleles of *HKLP* gene and diseases, for example cancer and neurological disorders. These association methods are within the scope of the invention.

Finally, the invention is directed to methods for the screening of substances or molecules that inhibit the expression of *HKLP*, as well as with methods for the screening of substances or molecules that interact with a HKLP polypeptide or that modulate the activity of a HKLP
20 polypeptide.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a block diagram of an exemplary computer system.

Figure 2 is a flow diagram illustrating one embodiment of a process 200 for comparing a new nucleotide or protein sequence with a database of sequences in order to determine the homology levels
25 between the new sequence and the sequences in the database.

Figure 3 is a flow diagram illustrating one embodiment of a process 250 in a computer for determining whether two sequences are homologous.

Figure 4 is a flow diagram illustrating one embodiment of an identifier process 300 for detecting the presence of a feature in a sequence.

30 BRIEF DESCRIPTION OF THE SEQUENCES PROVIDED IN THE SEQUENCE LISTING

SEQ ID Nos 1 and 2 contain the genomic sequence of the *HKLP* gene comprising the exons and introns, and the 3' regulatory region (downstream untranscribed region).

SEQ ID No 3 contains a cDNA sequence of the *HKLP* gene.

35 SEQ ID No 4 contains the amino acids sequence encoding by the cDNA of SEQ ID No 3.

SEQ ID Nos 5, 6, 7 and 8 respectively contain the nucleotide sequence of the amplicons 10-265, 10-266, 12-592 and 12-783.

SEQ ID No 9 contains a primer containing the additional PU 5' sequence described further in Example 2.

- 5 SEQ ID No 10 contains a primer containing the additional RP 5' sequence described further in Example 2.

In accordance with the regulations relating to Sequence Listings, the following codes have been used in the Sequence Listing to indicate the locations of biallelic markers within the sequences and to identify each of the alleles present at the polymorphic base. The code "r" in the sequences indicates that one allele of the polymorphic base is a guanine, while the other allele is an adenine. The code "y" in the sequences indicates that one allele of the polymorphic base is a thymine, while the other allele is a cytosine. The code "m" in the sequences indicates that one allele of the polymorphic base is an adenine, while the other allele is an cytosine. The code "k" in the sequences indicates that one allele of the polymorphic base is a guanine, while the other allele is a thymine. 10 The code "s" in the sequences indicates that one allele of the polymorphic base is a guanine, while the other allele is a cytosine. The code "w" in the sequences indicates that one allele of the polymorphic base is an adenine, while the other allele is an thymine. 15

The nucleotide code of the original allele for each biallelic marker is the following table:

Biallelic marker	Original allele
12-809-119	C
12-805-115	A
12-790-396	G
12-791-211	G
12-803-125	T
99-33040-321	T
12-810-77	A
12-787-103	A
12-793-383	T
12-792-233	A
99-41009-244	A
99-41009-111	C
12-593-174	T
12-589-152	T
12-785-200	T
12-785-393	A
12-588-103	G
12-603-191	T
12-586-414	G
12-602-196	T
12-602-350	C
12-587-379	A
12-596-124	G
12-808-52	A
12-808-75	G

DETAILED DESCRIPTION OF THE INVENTION

The present invention concerns polynucleotides and polypeptides related to the *HKLP* gene. Oligonucleotide probes and primers hybridizing specifically with a genomic or a cDNA sequence of *HKLP* are also part of the invention. A further object of the invention consists of recombinant
5 vectors comprising any of the nucleic acid sequences described in the present invention, and in particular recombinant vectors comprising a regulatory region of *HKLP* or a sequence encoding the *HKLP* protein, as well as cell hosts comprising said nucleic acid sequences or recombinant vectors. The invention also encompasses methods of screening of molecules which inhibit the expression of the *HKLP* gene or which modulate the activity of the *HKLP* protein. The invention also deals with
10 antibodies directed specifically against such polypeptides that are useful as diagnostic reagents.

The invention also concerns *HKLP*-related biallelic markers which can be used in any method of genetic analysis including linkage studies in families, linkage disequilibrium studies in populations and association studies of case-control populations. An important aspect of the present invention is that biallelic markers allow association studies to be performed to identify genes
15 involved in complex traits.

Definitions

Before describing the invention in greater detail, the following definitions are set forth to illustrate and define the meaning and scope of the terms used to describe the invention herein.

The terms "*HKLP* gene", when used herein, encompasses genomic, mRNA and cDNA
20 sequences encoding the *HKLP* protein, including the untranslated regulatory regions of the genomic DNA.

The term "heterologous protein", when used herein, is intended to designate any protein or polypeptide other than the *HKLP* protein. More particularly, the heterologous protein is a compound which can be used as a marker in further experiments with a *HKLP* regulatory region.

25 The term "isolated" requires that the material be removed from its original environment (e. g., the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or DNA or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotide could be part of a vector and/or such polynucleotide or polypeptide
30 could be part of a composition, and still be isolated in that the vector or composition is not part of its natural environment.

The term "purified" does not require absolute purity; rather, it is intended as a relative definition. Purification of starting material or natural material to at least one order of magnitude, preferably two or three orders, and more preferably four or five orders of magnitude is expressly
35 contemplated. As an example, purification from 0.1 % concentration to 10 % concentration is two orders of magnitude. The term "purified" is used herein to describe a polynucleotide or

polynucleotide vector of the invention which has been separated from other compounds including, but not limited to other nucleic acids, carbohydrates, lipids and proteins (such as the enzymes used in the synthesis of the polynucleotide), or the separation of covalently closed polynucleotides from linear polynucleotides. A polynucleotide is substantially pure when at least about 50%, preferably 60 to 75% of a sample exhibits a single polynucleotide sequence and conformation (linear versus covalently close). A substantially pure polynucleotide typically comprises about 50%, preferably 60 to 90% weight/weight of a nucleic acid sample, more usually about 95%, and preferably is over about 99% pure. Polynucleotide purity or homogeneity is indicated by a number of means well known in the art, such as agarose or polyacrylamide gel electrophoresis of a sample, followed by visualizing a single polynucleotide band upon staining the gel. For certain purposes higher resolution can be provided by using HPLC or other means well known in the art.

The term “polypeptide” refers to a polymer of amino acids without regard to the length of the polymer; thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide. This term also does not specify or exclude post-expression modifications of polypeptides, for example, polypeptides which include the covalent attachment of glycosyl groups, acetyl groups, phosphate groups, lipid groups and the like are expressly encompassed by the term polypeptide. Also included within the definition are polypeptides which contain one or more analogs of an amino acid (including, for example, non-naturally occurring amino acids, amino acids which only occur naturally in an unrelated biological system, modified amino acids from mammalian systems etc.), polypeptides with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally occurring.

The term “recombinant polypeptide” is used herein to refer to polypeptides that have been artificially designed and which comprise at least two polypeptide sequences that are not found as contiguous polypeptide sequences in their initial natural environment, or to refer to polypeptides which have been expressed from a recombinant polynucleotide.

The term “purified” is used herein to describe a polypeptide of the invention which has been separated from other compounds including, but not limited to nucleic acids, lipids, carbohydrates and other proteins. A polypeptide is substantially pure when at least about 50%, preferably 60 to 75% of a sample exhibits a single polypeptide sequence. A substantially pure polypeptide typically comprises about 50%, preferably 60 to 90% weight/weight of a protein sample, more usually about 95%, and preferably is over about 99% pure. Polypeptide purity or homogeneity is indicated by a number of means well known in the art, such as agarose or polyacrylamide gel electrophoresis of a sample, followed by visualizing a single polypeptide band upon staining the gel. For certain purposes higher resolution can be provided by using HPLC or other means well known in the art.

As used herein, the term “non-human animal” refers to any non-human vertebrate, birds and more usually mammals, preferably primates, farm animals such as swine, goats, sheep, donkeys, and horses, rabbits or rodents, more preferably rats or mice. As used herein, the term “animal” is used to

refer to any vertebrate, preferable a mammal. Both the terms "animal" and "mammal" expressly embrace human subjects unless preceded with the term "non-human".

As used herein, the term "antibody" refers to a polypeptide or group of polypeptides which are comprised of at least one binding domain, where an antibody binding domain is formed from the folding of variable domains of an antibody molecule to form three-dimensional binding spaces with an internal surface shape and charge distribution complementary to the features of an antigenic determinant of an antigen, which allows an immunological reaction with the antigen. Antibodies include recombinant proteins comprising the binding domains, as wells as fragments, including Fab, Fab', F(ab)₂, and F(ab')₂ fragments.

As used herein, an "antigenic determinant" is the portion of an antigen molecule, in this case a HKLP polypeptide, that determines the specificity of the antigen-antibody reaction. An "epitope" refers to an antigenic determinant of a polypeptide. An epitope can comprise as few as 3 amino acids in a spatial conformation which is unique to the epitope. Generally an epitope consists of at least 6 such amino acids, and more usually at least 8-10 such amino acids. Methods for determining the amino acids which make up an epitope include x-ray crystallography, 2-dimensional nuclear magnetic resonance, and epitope mapping e.g. the Pepscan method described by Geysen et al. 1984; PCT Publication No. WO 84/03564; and PCT Publication No. WO 84/03506.

Throughout the present specification, the expression "nucleotide sequence" may be employed to designate indifferently a polynucleotide or a nucleic acid. More precisely, the expression "nucleotide sequence" encompasses the nucleic material itself and is thus not restricted to the sequence information (i.e. the succession of letters chosen among the four base letters) that biochemically characterizes a specific DNA or RNA molecule.

As used interchangeably herein, the terms "nucleic acids", "oligonucleotides", and "polynucleotides" include RNA, DNA, or RNA/DNA hybrid sequences of more than one nucleotide in either single chain or duplex form. The term "nucleotide" as used herein as an adjective to describe molecules comprising RNA, DNA, or RNA/DNA hybrid sequences of any length in single-stranded or duplex form. The term "nucleotide" is also used herein as a noun to refer to individual nucleotides or varieties of nucleotides, meaning a molecule, or individual unit in a larger nucleic acid molecule, comprising a purine or pyrimidine, a ribose or deoxyribose sugar moiety, and a phosphate group, or phosphodiester linkage in the case of nucleotides within an oligonucleotide or polynucleotide. Although the term "nucleotide" is also used herein to encompass "modified nucleotides" which comprise at least one modifications (a) an alternative linking group, (b) an analogous form of purine, (c) an analogous form of pyrimidine, or (d) an analogous sugar, for examples of analogous linking groups, purine, pyrimidines, and sugars see for example PCT publication No. WO 95/04064. The polynucleotide sequences of the invention may be prepared by any known method, including synthetic, recombinant, *ex vivo* generation, or a combination thereof, as well as utilizing any purification methods known in the art.

A sequence which is "operably linked" to a regulatory sequence such as a promoter means that said regulatory element is in the correct location and orientation in relation to the nucleic acid to control RNA polymerase initiation and expression of the nucleic acid of interest. As used herein, the term "operably linked" refers to a linkage of polynucleotide elements in a functional relationship.

- 5 For instance, a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the coding sequence.

The terms "trait" and "phenotype" are used interchangeably herein and refer to any visible, detectable or otherwise measurable property of an organism such as symptoms of, or susceptibility to a disease for example. Typically the terms "trait" or "phenotype" are used herein to refer to
10 symptoms of, or susceptibility to a disease, a beneficial response to or side effects related to a treatment. Preferably, said trait can be, without to be limited to, cancers, developmental diseases, and neurological diseases.

The term "allele" is used herein to refer to variants of a nucleotide sequence. A biallelic polymorphism has two forms. Typically the first identified allele is designated as the original allele
15 whereas other alleles are designated as alternative alleles. Diploid organisms may be homozygous or heterozygous for an allelic form.

The term "heterozygosity rate" is used herein to refer to the incidence of individuals in a population which are heterozygous at a particular allele. In a biallelic system, the heterozygosity rate is on average equal to $2P_a(1-P_a)$, where P_a is the frequency of the least common allele. In order
20 to be useful in genetic studies, a genetic marker should have an adequate level of heterozygosity to allow a reasonable probability that a randomly selected person will be heterozygous.

The term "genotype" as used herein refers the identity of the alleles present in an individual or a sample. In the context of the present invention, a genotype preferably refers to the description of the biallelic marker alleles present in an individual or a sample. The term "genotyping" a sample
25 or an individual for a biallelic marker consists of determining the specific allele or the specific nucleotide carried by an individual at a biallelic marker.

The term "haplotype" refers to a combination of alleles present in an individual or a sample. In the context of the present invention, a haplotype preferably refers to a combination of biallelic marker alleles found in a given individual and which may be associated with a phenotype.

30 The term "polymorphism" as used herein refers to the occurrence of two or more alternative genomic sequences or alleles between or among different genomes or individuals. "Polymorphic" refers to the condition in which two or more variants of a specific genomic sequence can be found in a population. A "polymorphic site" is the locus at which the variation occurs. A single nucleotide polymorphism is the replacement of one nucleotide by another nucleotide at the polymorphic site.
35 Deletion of a single nucleotide or insertion of a single nucleotide also gives rise to single nucleotide polymorphisms. In the context of the present invention, "single nucleotide polymorphism"

preferably refers to a single nucleotide substitution. Typically, between different individuals, the polymorphic site may be occupied by two different nucleotides.

The term “biallelic polymorphism” and “biallelic marker” are used interchangeably herein to refer to a single nucleotide polymorphism having two alleles at a fairly high frequency in the
5 population. A “biallelic marker allele” refers to the nucleotide variants present at a biallelic marker site. Typically, the frequency of the less common allele of the biallelic markers of the present invention has been validated to be greater than 1%, preferably the frequency is greater than 10%, more preferably the frequency is at least 20% (i.e. heterozygosity rate of at least 0.32), even more preferably the frequency is at least 30% (i.e. heterozygosity rate of at least 0.42). A biallelic marker
10 wherein the frequency of the less common allele is 30% or more is termed a “high quality biallelic marker”.

The location of nucleotides in a polynucleotide with respect to the center of the polynucleotide are described herein in the following manner. When a polynucleotide has an odd number of nucleotides, the nucleotide at an equal distance from the 3' and 5' ends of the
15 polynucleotide is considered to be “at the center” of the polynucleotide, and any nucleotide immediately adjacent to the nucleotide at the center, or the nucleotide at the center itself is considered to be “within 1 nucleotide of the center.” With an odd number of nucleotides in a polynucleotide any of the five nucleotides positions in the middle of the polynucleotide would be considered to be within 2 nucleotides of the center, and so on. When a polynucleotide has an even
20 number of nucleotides, there would be a bond and not a nucleotide at the center of the polynucleotide. Thus, either of the two central nucleotides would be considered to be “within 1 nucleotide of the center” and any of the four nucleotides in the middle of the polynucleotide would be considered to be “within 2 nucleotides of the center”, and so on. For polymorphisms which involve the substitution, insertion or deletion of 1 or more nucleotides, the polymorphism, allele or
25 biallelic marker is “at the center” of a polynucleotide if the difference between the distance from the substituted, inserted, or deleted polynucleotides of the polymorphism and the 3' end of the polynucleotide, and the distance from the substituted, inserted, or deleted polynucleotides of the polymorphism and the 5' end of the polynucleotide is zero or one nucleotide. If this difference is 0 to 3, then the polymorphism is considered to be “within 1 nucleotide of the center.” If the difference
30 is 0 to 5, the polymorphism is considered to be “within 2 nucleotides of the center.” If the difference is 0 to 7, the polymorphism is considered to be “within 3 nucleotides of the center,” and so on.

The terms “complementary” or “complement thereof” are used herein to refer to the sequences of polynucleotides which is capable of forming Watson & Crick base pairing with another specified polynucleotide throughout the entirety of the complementary region. For the purpose of the
35 present invention, a first polynucleotide is deemed to be complementary to a second polynucleotide when each base in the first polynucleotide is paired with its complementary base. Complementary bases are, generally, A and T (or A and U), or C and G. “Complement” is used herein as a synonym

from “complementary polynucleotide”, “complementary nucleic acid” and “complementary nucleotide sequence”. These terms are applied to pairs of polynucleotides based solely upon their sequences and not any particular set of conditions under which the two polynucleotides would actually bind.

5 Variants and Fragments

1- Polynucleotides

The invention also relates to variants and fragments of the polynucleotides described herein, particularly of a *HKLP* gene containing one or more biallelic markers according to the invention.

10 Variants of polynucleotides, as the term is used herein, are polynucleotides that differ from a reference polynucleotide. A variant of a polynucleotide may be a naturally occurring variant such as a naturally occurring allelic variant, or it may be a variant that is not known to occur naturally. Such non-naturally occurring variants of the polynucleotide may be made by mutagenesis techniques, including those applied to polynucleotides, cells or organisms. Generally, differences are limited so that the nucleotide sequences of the reference and the variant are closely similar overall and, in many
15 regions, identical.

Variants of polynucleotides according to the invention include, without being limited to, nucleotide sequences which are at least 95% identical to a polynucleotide selected from the group consisting of the nucleotide sequences of SEQ ID Nos 1-3 or to any polynucleotide fragment of at least 8 consecutive nucleotides of a polynucleotide selected from the group consisting of the nucleotide
20 sequences of SEQ ID Nos 1-3, and preferably at least 99% identical, more particularly at least 99.5% identical, and most preferably at least 99.8% identical to a polynucleotide selected from the group consisting of the nucleotide sequences of SEQ ID Nos 1-3 or to any polynucleotide fragment of at least 8 consecutive nucleotides of a polynucleotide selected from the group consisting of the nucleotide sequences of SEQ ID No 1-3.

25 Nucleotide changes present in a variant polynucleotide may be silent, which means that they do not alter the amino acids encoded by the polynucleotide. However, nucleotide changes may also result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence. The substitutions, deletions or additions may involve one or more nucleotides. The variants may be altered in coding or non-coding regions or both. Alterations
30 in the coding regions may produce conservative or non-conservative amino acid substitutions, deletions or additions.

In the context of the present invention, particularly preferred embodiments are those in which the polynucleotides encode polypeptides which retain substantially the same biological function or activity as the mature *HKLP* protein, or those in which the polynucleotides encode
35 polypeptides which maintain or increase a particular biological activity, while reducing a second biological activity

A polynucleotide fragment is a polynucleotide having a sequence that is entirely the same as part but not all of a given nucleotide sequence, preferably the nucleotide sequence of a *HKLP* gene, and variants thereof. The fragment can be a portion of an intron or an exon of a *HKLP* gene. It can also be a portion of the regulatory regions of *HKLP*. Preferably, such fragments comprise at least one of the biallelic markers A1 to A32 or the complements thereto or a biallelic marker in linkage disequilibrium therewith.

Such fragments may be "free-standing", i.e. not part of or fused to other polynucleotides, or they may be comprised within a single larger polynucleotide of which they form a part or region. Indeed, several of these fragments may be present within a single larger polynucleotide.

Optionally, such fragments may consist of, or consist essentially of a contiguous span of at least 8, 10, 12, 15, 18, 20, 25, 35, 40, 50, 70, 80, 100, 250, 500 or 1000 nucleotides in length.

2- Polypeptides

The invention also relates to variants, fragments, analogs and derivatives of the polypeptides described herein, including mutated HKLP proteins.

The variant may be 1) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue and such substituted amino acid residue may or may not be one encoded by the genetic code, or 2) one in which one or more of the amino acid residues includes a substituent group, or 3) one in which the mutated HKLP is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or 4) one in which the additional amino acids are fused to the mutated HKLP, such as a leader or secretory sequence or a sequence which is employed for purification of the mutated HKLP or a preprotein sequence. Such variants are deemed to be within the scope of those skilled in the art.

A polypeptide fragment is a polypeptide having a sequence that entirely is the same as part but not all of a given polypeptide sequence, preferably a polypeptide encoded by a *HKLP* gene and variants thereof.

In the case of an amino acid substitution in the amino acid sequence of a polypeptide according to the invention, one or several amino acids can be replaced by "equivalent" amino acids. The expression "equivalent" amino acid is used herein to designate any amino acid that may be substituted for one of the amino acids having similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydrophobic nature of the polypeptide to be substantially unchanged. Generally, the following groups of amino acids represent equivalent changes: (1) Ala, Pro, Gly, Glu, Asp, Gln, Asn, Ser, Thr; (2) Cys, Ser, Tyr, Thr; (3) Val, Ile, Leu, Met, Ala, Phe; (4) Lys, Arg, His; (5) Phe, Tyr, Trp, His.

A specific embodiment of a modified HKLP peptide molecule of interest according to the present invention, includes, but is not limited to, a peptide molecule which is resistant to proteolysis, is a peptide in which the -CONH- peptide bond is modified and replaced by a (CH₂NH) reduced

bond, a (NHCO) retro inverso bond, a (CH₂-O) methylene-oxy bond, a (CH₂-S) thiomethylene bond, a (CH₂CH₂) carba bond, a (CO-CH₂) cetomethylene bond, a (CHOH-CH₂) hydroxyethylene bond), a (N-N) bound, a E-alcene bond or also a -CH=CH- bond. The invention also encompasses a human HKLP polypeptide or a fragment or a variant thereof in which at least one peptide bond has
5 been modified as described above.

Such fragments may be "free-standing", i.e. not part of or fused to other polypeptides, or they may be comprised within a single larger polypeptide of which they form a part or region. However, several fragments may be comprised within a single larger polypeptide.

As representative examples of polypeptide fragments of the invention, there may be
10 mentioned those which have a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, or 100 amino acids.

Identity Between Nucleic Acids Or Polypeptides

The terms "percentage of sequence identity" and "percentage homology" are used interchangeably herein to refer to comparisons among polynucleotides and polypeptides, and are
15 determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue
20 occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity. Homology is evaluated using any of the variety of sequence comparison algorithms and programs known in the art. Such algorithms and programs include, but are by no means limited to, TBLASTN, BLASTP, FASTA, TFASTA, and CLUSTALW
25 (Pearson and Lipman, 1988; Altschul et al., 1990; Thompson et al., 1994; Higgins et al., 1996; Altschul et al., 1990; Altschul et al., 1993). In a particularly preferred embodiment, protein and nucleic acid sequence homologies are evaluated using the Basic Local Alignment Search Tool ("BLAST") which is well known in the art (see, e.g., Karlin and Altschul, 1990; Altschul et al., 1990, 1993, 1997). In particular, five specific BLAST programs are used to perform the following
30 task:

(1) BLASTP and BLAST3 compare an amino acid query sequence against a protein sequence database;

(2) BLASTN compares a nucleotide query sequence against a nucleotide sequence database;

35 (3) BLASTX compares the six-frame conceptual translation products of a query nucleotide sequence (both strands) against a protein sequence database;

(4) TBLASTN compares a query protein sequence against a nucleotide sequence database translated in all six reading frames (both strands); and

(5) TBLASTX compares the six-frame translations of a nucleotide query sequence against the six-frame translations of a nucleotide sequence database.

- 5 The BLAST programs identify homologous sequences by identifying similar segments, which are referred to herein as "high-scoring segment pairs," between a query amino or nucleic acid sequence and a test sequence which is preferably obtained from a protein or nucleic acid sequence database. High-scoring segment pairs are preferably identified (i.e., aligned) by means of a scoring matrix, many of which are known in the art. Preferably, the scoring matrix used is the BLOSUM62 matrix
- 10 (Gonnet et al., 1992; Henikoff and Henikoff, 1993). Less preferably, the PAM or PAM250 matrices may also be used (see, e.g., Schwartz and Dayhoff, eds., 1978). The BLAST programs evaluate the statistical significance of all high-scoring segment pairs identified, and preferably selects those segments which satisfy a user-specified threshold of significance, such as a user-specified percent homology. Preferably, the statistical significance of a high-scoring segment pair is
- 15 evaluated using the statistical significance formula of Karlin (see, e.g., Karlin and Altschul, 1990).

Stringent Hybridization Conditions

For the purpose of defining such a hybridizing nucleic acid according to the invention, the stringent hybridization conditions are the followings :

- the hybridization step is realized at 65°C in the presence of 6 x SSC buffer, 5 x Denhardt's
- 20 solution, 0,5% SDS and 100µg/ml of salmon sperm DNA.

The hybridization step is followed by four washing steps :

- two washings during 5 min, preferably at 65°C in a 2 x SSC and 0.1%SDS buffer;
 - one washing during 30 min, preferably at 65°C in a 2 x SSC and 0.1% SDS buffer,
 - one washing during 10 min, preferably at 65°C in a 0.1 x SSC and 0.1%SDS buffer,
- 25 these hybridization conditions being suitable for a nucleic acid molecule of about 20 nucleotides in length. There is no need to say that the hybridization conditions described above are to be adapted according to the length of the desired nucleic acid, following techniques well known to the one skilled in the art. The suitable hybridization conditions may for example be adapted according to the teachings disclosed in the book of Hames and Higgins (1985).

Genomic Sequences Of The *HKLP* Gene

- 30 The present invention concerns the genomic sequence of *HKLP* comprising the 2 genomic contigs of SEQ ID Nos 1 and 2. The present invention encompasses *HKLP* gene, or *HKLP* genomic sequences consisting of, consisting essentially of, or comprising a sequence selected from the group consisting of SEQ ID Nos 1 and 2, a sequence complementary thereto, as well as fragments and
- 35 variants thereof. These polynucleotides may be purified, isolated, or recombinant.

The invention also encompasses a purified, isolated, or recombinant polynucleotides comprising a nucleotide sequence having at least 70, 75, 80, 85, 90, or 95% nucleotide identity with a nucleotide sequence selected from the group consisting of SEQ ID Nos 1 and 2 or a complementary sequence thereto or a fragment thereof. The nucleotide differences as regards to the
5 nucleotide sequences of SEQ ID Nos 1 and 2 may be generally randomly distributed throughout the entire nucleic acid. Nevertheless, preferred nucleic acids are those wherein the nucleotide differences as regards to the nucleotide sequences of SEQ ID Nos 1 and 2 are predominantly located outside the coding sequences contained in the exons. These nucleic acids, as well as their fragments and variants, may be used as oligonucleotide primers or probes in order to detect the presence of a
10 copy of the *HKLP* gene in a test sample, or alternatively in order to amplify a target nucleotide sequence within the *HKLP* sequences.

Another object of the invention consists of a purified, isolated, or recombinant nucleic acids that hybridizes with a nucleotide sequence selected from the group consisting of SEQ ID Nos 1 and 2 or a complementary sequence thereto or a variant thereof, under the stringent hybridization
15 conditions as defined above.

Particularly preferred nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, or 200 nucleotides of SEQ ID No 1 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, 10, 20, 30, 40 or 50 of the following
20 nucleotide positions of SEQ ID No 1: 1-39624, 39705-40589, 40666-43629, 43710-44203, 44311-45125, 45210-45440, 45622-45717, 45791-68580, 68675-70246, 70396-72421, 72601-73295, 73434-74648, 74898-83055, 83175-85192, 85279-85609, 85740-85906, 86070-88304, 88396-90585, 90705-91767, 91824-94380, 94490-96296, 96364-97184, 97270-101167, 101274-109465, 109581-110228, 110363-111819, 111882-113636, 113783-113945, 114186-117002, 117075-
25 119676, and 119677-121162.

Additional preferred nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 2 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide
30 positions of SEQ ID No 2: 1-1600, 1751-2138, 2332-2539, 2659-3829 and 8885-10884.

Additional preferred nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 1, or 2, or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 nucleotide positions of any one of
35 the following ranges of nucleotide positions of

(a) SEQ ID No 1: 1-1000, 1001-2000, 2001-3000, 3001-4000, 4001-5000, 5001-6000, 6001-7000, 7001-8000, 8001-9000, 9001-10000, 10001-11000, 11001-12000, 12001-13000, 13001-

14000, 14001-15000, 15001-16000, 16001-17000, 17001-18000, 18001-19000, 19001-20000, 20001-21000, 21001-22000, 22001-23000, 23001-24000, 24001-25000, 25001-26000, 26001-27000, 27001-28000, 28001-29000, 29001-30000, 30001-31000, 31001-32000, 32001-33000, 33001-34000, 34001-35000, 35001-36000, 36001-37000, 37001-38000, 38001-39000, 39001-39624, 39705-40589, 40666-43629, 43710-44203, 44311-45125, 45210-45440, 45622-45717, 45791-68580, 68675-70246, 70396-72421, 72601-73295, 73434-74648, 74898-83055, 83175-85192, 85279-85609, 85740-85906, 86070-88304, 88396-90585, 90705-91767, 91824-94380, 94490-96296, 96364-97184, 97270-101167, 101274-109465, 109581-110228, 110363-111819, 111882-113636, 113783-113945, 114186-117002, 117075-119676, and 119677-121162; and

10 (b) SEQ ID No 2: 1-1600, 1751-2138, 2332-2539, 2659-3829 and 8885-10884.

Additional preferred nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 1 or the complements thereof, wherein said contiguous span comprises a G at position 7159 of SEQ ID No 1. Further preferred

15 nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 2 or the complements thereof, wherein said contiguous span comprises a C either at position 2551 or 4500 of SEQ ID No 2. It should be noted that nucleic acid fragments of any size and sequence may also be comprised by the polynucleotides described in this

20 section.

The *HKLP* genomic nucleic acid comprises at least 48 exons. The exon positions in SEQ ID Nos 1 and 2 are detailed below in the Table A. The first exon which has been identified in the cDNA of the present invention is not comprised in the genomic sequence described in the present invention. The sequence of the first exon begins at the position 1 of SEQ ID No 3 and ends at the

25 position 292. The genomic sequence of SEQ ID Nos 1 and 2 comprises respectively 44 and 4 exons.

Thus, the invention embodies purified, isolated, or recombinant polynucleotides comprising a nucleotide sequence selected from the group consisting of the exons of the *HKLP* gene, or a sequence complementary thereto. The invention also deals with purified, isolated, or recombinant nucleic acids comprising a combination of at least two exons of the *HKLP* gene, wherein the

30 polynucleotides are arranged within the nucleic acid, from the 5'-end to the 3'-end of said nucleic acid, in the same order as in SEQ ID Nos 1 and 2.

The position of the introns is detailed in Table A. Thus, the invention embodies purified, isolated, or recombinant polynucleotides comprising a nucleotide sequence selected from the group consisting of the introns of the *HKLP* gene, or a sequence complementary thereto.

35 Thus, the present invention deals with a purified or isolated nucleic acid encoding a *HKLP* protein having the amino acid sequence of SEQ ID No 4 or a peptide fragment or variant thereof. In a specific embodiment, such a purified or isolated nucleic acid comprises a polynucleotide selected

from the group consisting of SEQ ID Nos 1 and 2, or a complementary sequence thereto or a fragment or a variant thereof.

Table A

Exon	Position in SEQ ID No 1		Intron	Position in SEQ ID No 1	
	Beginning	End		Beginning	End
2	4844	4920	2	4921	7057
3	7058	7237	3	7238	10468
4	10469	10534	4	10535	15939
5	15940	16118	5	16119	16711
6	16712	16823	6	16824	20053
7	20054	20131	7	20132	20792
8	20793	20858	8	20859	21557
9	21558	21575	9	21576	23970
10	23971	24046	10	24047	24863
11	24864	24942	11	24943	26528
12	26529	26671	12	26672	27639
13	27640	27681	13	27682	30862
14	30863	31074	14	31075	39624
15	39625	39704	15	39705	40589
16	40590	40665	16	40666	43629
17	43630	43709	17	43710	44203
18	44204	44310	18	44311	45125
19	45126	45209	19	45210	45440
20	45441	45621	20	45622	45717
21	45718	45790	21	45791	68580
22	68581	68674	22	68675	70246
23	70247	70395	23	70396	72421
24	72422	72600	24	72601	73295
25	73296	73433	25	73434	74648
26	74649	74897	26	74898	83055
27	83056	83174	27	83175	85192
28	85193	85278	28	85279	85609
29	85610	85739	29	85740	85906
30	85907	86069	30	86070	88304
31	88305	88395	31	88396	90585
32	90586	90704	32	90705	91767
33	91768	91823	33	91824	94380
34	94381	94489	34	94490	96296
35	96297	96363	35	96364	97184
36	97185	97269	36	97270	101167
37	101168	101273	37	101274	109465
38	109466	109580	38	109581	110228
39	110229	110362	39	110363	111819
40	111820	111881	40	111882	113636
41	113637	113782	41	113783	113945
42	113946	114185	42	114186	117002
43	117003	117074	43	117075	119676
44	119677	119798	44	119677	121162
Exon	Position in SEQ ID No 2		Intron	Position in SEQ ID No 2	
	Beginning	End		Beginning	End
45	1601	1750	44	1	1600
46	2139	2331	45	1751	2138
47	2540	2658	46	2332	2539
48	3830	8884	47	2659	3829

The *HKLP* genomic sequence is covered by two fragments. Indeed, one segment is unknown in the intron 44. The inventors think that this segment, which seems to comprise 20 to 30 nucleotides, forms a superstructure which prevents the sequencing. This superstructure comprises two polyG at each end of the segment.

5 While this section is entitled "Genomic Sequences of *HKLP*," it should be noted that nucleic acid fragments of any size and sequence may also be comprised by the polynucleotides described in this section, flanking the genomic sequences of *HKLP* on either side or between two or more such genomic sequences.

HKLP cDNA Sequences

10 The expression of the *HKLP* gene has been shown to lead to the production of at least one mRNA species, the nucleic acid sequence of which is set forth in SEQ ID No 3.

Another object of the invention is a purified, isolated, or recombinant nucleic acid comprising the nucleotide sequence of SEQ ID No 3, complementary sequences thereto, as well as allelic variants, and fragments thereof. Moreover, preferred polynucleotides of the invention include
15 purified, isolated, or recombinant *HKLP* cDNAs consisting of, consisting essentially of, or comprising the sequence of SEQ ID No 3. Particularly preferred embodiments of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 3 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of
20 the following nucleotide positions of SEQ ID No 3: 391-1619 and 6988-10682. Additional preferred embodiments of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 3 or the complements thereof, wherein said contiguous span comprises a nucleotide selected in the group consisting of a C at position 5487, and a C at
25 position 6265 of SEQ ID No 3.

The invention also pertains to a purified or isolated nucleic acid having at least 95% of nucleotide identity with the nucleotide sequence of SEQ ID No 3 or a fragment thereof or a complementary sequence thereto, advantageously 99 %, preferably 99.5% nucleotide identity and most preferably 99.8% nucleotide identity with the nucleotide sequence of SEQ ID No 3 or a
30 fragment thereof or a complementary sequence thereto.

Another object of the invention consists of a purified, isolated, or recombinant nucleic acids that hybridizes with the nucleotide sequence of SEQ ID No 3 or a complementary sequence thereto or a variant thereof, under the stringent hybridization conditions as defined above.

The cDNA of SEQ ID No 3 includes a 5'-UTR region starting from the nucleotide at
35 position 1 and ending at the nucleotide in position 186 of SEQ ID No 3. The cDNA of SEQ ID No 3 includes a 3'-UTR region starting from the nucleotide at position 5638 and ending at the nucleotide

at position 10682 of SEQ ID No 3. The polyadenylation site starts from the nucleotide at position 10631 and ends at the nucleotide in position 10636 of SEQ ID No 3.

Consequently, the invention concerns a purified, isolated, and recombinant nucleic acids comprising a nucleotide sequence of the 3'UTR of the *HKLP* cDNA, a sequence complementary thereto, or an allelic variant thereof.

While this section is entitled " *HKLP* cDNA Sequences," it should be noted that nucleic acid fragments of any size and sequence may also be comprised by the polynucleotides described in this section, flanking the genomic sequences of *HKLP* on either side or between two or more such genomic sequences.

10

Coding Regions

The *HKLP* open reading frame is contained in the corresponding mRNA of SEQ ID No 3. More precisely, the effective *HKLP* coding sequence (CDS) includes the region between nucleotide position 187 (first nucleotide of the ATG codon) and nucleotide position 5637 (end nucleotide of the TGA codon) of SEQ ID No 3.

15

The present invention also embodies isolated, purified, and recombinant polynucleotides which encode a polypeptides comprising a contiguous span of at least 6 amino acids, preferably at least 8 or 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, or 100 amino acids of SEQ ID No 4, wherein said contiguous span includes at least 1, 2, 3, 5 or 10 of the amino acid positions 1-478 of the SEQ ID No 4.

20

The above disclosed polynucleotide that contains the coding sequence of the *HKLP* gene may be expressed in a desired host cell or a desired host organism, when this polynucleotide is placed under the control of suitable expression signals. The expression signals may be either the expression signals contained in the regulatory regions in the *HKLP* gene of the invention or in contrast the signals may be exogenous regulatory nucleic sequences. Such a polynucleotide, when placed under the suitable expression signals, may also be inserted in a vector for its expression and/or amplification.

25

Regulatory Sequences Of *HKLP*

As mentioned, the genomic sequence of the *HKLP* gene contains regulatory sequences in the non-coding 3'-flanking region that border the *HKLP* coding region. The 3'-regulatory sequence of the *HKLP* gene is localized between nucleotide position 8885 and nucleotide position 10884 of SEQ ID No 2. Polynucleotides derived from the 3' regulatory region are useful in order to detect the presence of at least a copy of a nucleotide sequence of SEQ ID No 2 or a fragment thereof in a test sample.

30

In order to identify the relevant biologically active polynucleotide fragments or variants of the 3' regulatory region from SEQ ID No 2, the one skill in the art will refer to the book of Sambrook et al.(Sambrook, 1989) which describes the use of a recombinant vector carrying a marker gene (i.e.

35

beta galactosidase, chloramphenicol acetyl transferase, etc.) the expression of which will be detected when placed under the control of a biologically active polynucleotide fragments or variants of SEQ ID No 2. The level of reporter protein is assayed and compared to the level obtained from a vector which lacks an insert in the cloning site. The presence of an elevated expression level in the vector
5 containing the insert with respect to the control vector indicates the presence of a biologically active polynucleotide in the insert.

Polynucleotides carrying the regulatory elements located at the 3' end of the *HKLP* coding region may be advantageously used to control the transcriptional and translational activity of an heterologous polynucleotide of interest.

10 Thus, the present invention also concerns a purified or isolated nucleic acid comprising a polynucleotide of the 3' regulatory regions, or a sequence complementary thereto or a biologically active fragment or variant thereof.

Preferred fragments of the 3' regulatory region are at least 50, 100, 150, 200, 300 or 400 bases in length.

15 By "biologically active" polynucleotide derivatives of SEQ ID No 2 are polynucleotides comprising or alternatively consisting in a fragment of said polynucleotide which is functional as a regulatory region for expressing a recombinant polypeptide or a recombinant polynucleotide in a recombinant cell host. It could act either as an enhancer or as a repressor.

For the purpose of the invention, a nucleic acid or polynucleotide is "functional" as a
20 regulatory region for expressing a recombinant polypeptide or a recombinant polynucleotide if said regulatory polynucleotide contains nucleotide sequences which contain transcriptional and translational regulatory information, and such sequences are "operably linked" to nucleotide sequences which encode the desired polypeptide or the desired polynucleotide.

The regulatory polynucleotides of the invention may be prepared from the nucleotide
25 sequence of SEQ ID No 2 by cleavage using suitable restriction enzymes, as described for example in the book of Sambrook et al.(1989). The regulatory polynucleotides may also be prepared by digestion of SEQ ID No 2 by an exonuclease enzyme, such as Bal31 (Wabiko et al., 1986). These regulatory polynucleotides can also be prepared by nucleic acid chemical synthesis, as described elsewhere in the specification.

30 The regulatory polynucleotides according to the invention may be part of a recombinant expression vector that may be used to express a coding sequence in a desired host cell or host organism. The recombinant expression vectors according to the invention are described elsewhere in the specification.

A preferred 3'-regulatory polynucleotide of the invention includes the 3'-untranslated region
35 (3'-UTR) of the *HKLP* cDNA, or a biologically active fragment or variant thereof.

Polynucleotide Constructs

The terms "polynucleotide construct" and "recombinant polynucleotide" are used interchangeably herein to refer to linear or circular, purified or isolated polynucleotides that have been artificially designed and which comprise at least two nucleotide sequences that are not found as
5 contiguous nucleotide sequences in their initial natural environment.

DNA Construct That Enables Directing Temporal And Spatial *HKLP* Gene Expression In Recombinant Cell Hosts And In Transgenic Animals.

In order to study the physiological and phenotypic consequences of a lack of synthesis of the *HKLP* protein, both at the cell level and at the multi cellular organism level, the invention also
10 encompasses DNA constructs and recombinant vectors enabling a conditional expression of a specific allele of the *HKLP* genomic sequence or cDNA and also of a copy of this genomic sequence or cDNA harboring substitutions, deletions, or additions of one or more bases as regards to the *HKLP* nucleotide sequence of SEQ ID Nos 1-3, or a fragment thereof, these base substitutions, deletions or additions being located either in an exon, an intron or a regulatory sequence, but
15 preferably in an exon of the *HKLP* genomic sequence or within the *HKLP* cDNA of SEQ ID No 3. In a preferred embodiment, the *HKLP* sequence comprises a biallelic marker of the present invention. In a preferred embodiment, the *HKLP* sequence comprises a biallelic marker of the present invention, preferably one of the biallelic markers A1 to A32.

The present invention embodies recombinant vectors comprising any one of the
20 polynucleotides described in the present invention.

A first preferred DNA construct is based on the tetracycline resistance operon *tet* from *E. coli* transposon Tn10 for controlling the *HKLP* gene expression, such as described by Gossen et al.(1992, 1995) and Furth et al.(1994). Such a DNA construct contains seven *tet* operator sequences from Tn10 (*tet*op) that are fused to a minimal promoter, said minimal promoter being operably
25 linked to a polynucleotide of interest that codes either for a sense or an antisense oligonucleotide or for a polypeptide, including a *HKLP* polypeptide or a peptide fragment thereof. This DNA construct is functional as a conditional expression system for the nucleotide sequence of interest when the same cell also comprises a nucleotide sequence coding for either the wild type (tTA) or the mutant (rTA) repressor fused to the activating domain of viral protein VP16 of herpes simplex virus, placed
30 under the control of a promoter, such as the HCMVIE1 enhancer/promoter or the MMTV-LTR. Indeed, a preferred DNA construct of the invention comprise both the polynucleotide containing the *tet* operator sequences and the polynucleotide containing a sequence coding for the tTA or the rTA repressor.

In a specific embodiment, the conditional expression DNA construct contains the sequence
35 encoding the mutant tetracycline repressor rTA, the expression of the polynucleotide of interest is silent in the absence of tetracycline and induced in its presence.

DNA Constructs Allowing Homologous Recombination: Replacement Vectors

A second preferred DNA construct will comprise, from 5'-end to 3'-end: (a) a first nucleotide sequence that is comprised in the *HKLP* genomic sequence; (b) a nucleotide sequence comprising a positive selection marker, such as the marker for neomycine resistance (*neo*); and (c) a second nucleotide sequence that is comprised in the *HKLP* genomic sequence, and is located on the genome downstream the first *HKLP* nucleotide sequence (a).

In a preferred embodiment, this DNA construct also comprises a negative selection marker located upstream the nucleotide sequence (a) or downstream the nucleotide sequence (c).

Preferably, the negative selection marker consists of the thymidine kinase (*tk*) gene (Thomas et al., 1986), the hygromycine beta gene (Te Riele et al., 1990), the *hprt* gene (Van der Lugt et al., 1991; Reid et al., 1990) or the Diphtheria toxin A fragment (*Dt-A*) gene (Nada et al., 1993; Yagi et al., 1990). Preferably, the positive selection marker is located within a *HKLP* exon sequence so as to interrupt the sequence encoding a *HKLP* protein. These replacement vectors are described, for example, by Thomas et al. (1986; 1987), Mansour et al. (1988) and Koller et al. (1992).

The first and second nucleotide sequences (a) and (c) may be indifferently located within a *HKLP* regulatory sequence, an intronic sequence, an exon sequence or a sequence containing both regulatory and/or intronic and/or exon sequences. The size of the nucleotide sequences (a) and (c) ranges from 1 to 50 kb, preferably from 1 to 10 kb, more preferably from 2 to 6 kb and most preferably from 2 to 4 kb.

DNA Constructs Allowing Homologous Recombination: Cre-LoxP System.

These new DNA constructs make use of the site specific recombination system of the P1 phage. The P1 phage possesses a recombinase called Cre which interacts specifically with a 34 base pairs *loxP* site. The *loxP* site is composed of two palindromic sequences of 13 bp separated by a 8 bp conserved sequence (Hoess et al., 1986). The recombination by the Cre enzyme between two *loxP* sites having an identical orientation leads to the deletion of the DNA fragment.

The Cre-*loxP* system used in combination with a homologous recombination technique has been first described by Gu et al. (1993, 1994). Briefly, a nucleotide sequence of interest to be inserted in a targeted location of the genome harbors at least two *loxP* sites in the same orientation and located at the respective ends of a nucleotide sequence to be excised from the recombinant genome. The excision event requires the presence of the recombinase (Cre) enzyme within the nucleus of the recombinant cell host. The recombinase enzyme may be brought at the desired time either by (a) incubating the recombinant cell hosts in a culture medium containing this enzyme, by injecting the Cre enzyme directly into the desired cell, such as described by Araki et al. (1995), or by lipofection of the enzyme into the cells, such as described by Baubonis et al. (1993); (b) transfecting the cell host with a vector comprising the *Cre* coding sequence operably linked to a promoter functional in the recombinant cell host, which promoter being optionally inducible, said vector being

introduced in the recombinant cell host, such as described by Gu et al.(1993) and Sauer et al.(1988);
(c) introducing in the genome of the cell host a polynucleotide comprising the *Cre* coding sequence
operably linked to a promoter functional in the recombinant cell host, which promoter is optionally
inducible, and said polynucleotide being inserted in the genome of the cell host either by a random
5 insertion event or an homologous recombination event, such as described by Gu et al.(1994).

In a specific embodiment, the vector containing the sequence to be inserted in the *HKLP*
gene by homologous recombination is constructed in such a way that selectable markers are flanked
by *loxP* sites of the same orientation, it is possible, by treatment by the *Cre* enzyme, to eliminate the
selectable markers while leaving the *HKLP* sequences of interest that have been inserted by an
10 homologous recombination event. Again, two selectable markers are needed: a positive selection
marker to select for the recombination event and a negative selection marker to select for the
homologous recombination event. Vectors and methods using the *Cre-loxP* system are described by
Zou et al.(1994).

Thus, a third preferred DNA construct of the invention comprises, from 5'-end to 3'-end: (a)
15 a first nucleotide sequence that is comprised in the *HKLP* genomic sequence; (b) a nucleotide
sequence comprising a polynucleotide encoding a positive selection marker, said nucleotide
sequence comprising additionally two sequences defining a site recognized by a recombinase, such
as a *loxP* site, the two sites being placed in the same orientation; and (c) a second nucleotide
sequence that is comprised in the *HKLP* genomic sequence, and is located on the genome
20 downstream of the first *HKLP* nucleotide sequence (a).

The sequences defining a site recognized by a recombinase, such as a *loxP* site, are
preferably located within the nucleotide sequence (b) at suitable locations bordering the nucleotide
sequence for which the conditional excision is sought. In one specific embodiment, two *loxP* sites
are located at each side of the positive selection marker sequence, in order to allow its excision at a
25 desired time after the occurrence of the homologous recombination event.

In a preferred embodiment of a method using the third DNA construct described above, the
excision of the polynucleotide fragment bordered by the two sites recognized by a recombinase,
preferably two *loxP* sites, is performed at a desired time, due to the presence within the genome of
the recombinant host cell of a sequence encoding the *Cre* enzyme operably linked to a promoter
30 sequence, preferably an inducible promoter, more preferably a tissue-specific promoter sequence and
most preferably a promoter sequence which is both inducible and tissue-specific, such as described
by Gu et al.(1994).

The presence of the *Cre* enzyme within the genome of the recombinant cell host may result
of the breeding of two transgenic animals, the first transgenic animal bearing the *HKLP*-derived
35 sequence of interest containing the *loxP* sites as described above and the second transgenic animal
bearing the *Cre* coding sequence operably linked to a suitable promoter sequence, such as described
by Gu et al.(1994).

Spatio-temporal control of the Cre enzyme expression may also be achieved with an adenovirus based vector that contains the Cre gene thus allowing infection of cells, or *in vivo* infection of organs, for delivery of the Cre enzyme, such as described by Anton and Graham (1995) and Kanegae et al.(1995).

- 5 The DNA constructs described above may be used to introduce a desired nucleotide sequence of the invention, preferably a *HKLP* genomic sequence or a *HKLP* cDNA sequence, and most preferably an altered copy of a *HKLP* genomic or cDNA sequence, within a predetermined location of the targeted genome, leading either to the generation of an altered copy of a targeted gene (knock-out homologous recombination) or to the replacement of a copy of the targeted gene by
10 another copy sufficiently homologous to allow an homologous recombination event to occur (knock-in homologous recombination). In a specific embodiment, the DNA constructs described above may be used to introduce a *HKLP* genomic sequence or a *HKLP* cDNA sequence. Optionally, said sequence comprises at least one biallelic marker of the present invention, preferably at least one biallelic marker selected from the group consisting of A1 to A32.

15 **Nuclear Antisense DNA Constructs**

- Other compositions containing a vector of the invention comprising an oligonucleotide fragment of the nucleic sequence SEQ ID No 3, preferably a fragment including the start codon of the *HKLP* gene, as an antisense tool that inhibits the expression of the corresponding *HKLP* gene. Preferred methods using antisense polynucleotide according to the present invention are the
20 procedures described by Sczakiel et al.(1995) or those described in PCT Application No WO 95/24223.

- Preferably, the antisense tools are chosen among the polynucleotides (15-200 bp long) that are complementary to the 5' end of the *HKLP* mRNA. In one embodiment, a combination of different antisense polynucleotides complementary to different parts of the desired targeted gene are
25 used.

Preferred antisense polynucleotides according to the present invention are complementary to a sequence of the mRNAs of *HKLP* that contains either the translation initiation codon ATG or a splicing site. Further preferred antisense polynucleotides according to the invention are complementary of the splicing site of the *HKLP* mRNA.

- 30 Preferably, the antisense polynucleotides of the invention have a 3' polyadenylation signal that has been replaced with a self-cleaving ribozyme sequence, such that RNA polymerase II transcripts are produced without poly(A) at their 3' ends, these antisense polynucleotides being incapable of export from the nucleus, such as described by Liu et al.(1994). In a preferred embodiment, these *HKLP* antisense polynucleotides also comprise, within the ribozyme cassette, a
35 histone stem-loop structure to stabilize cleaved transcripts against 3'-5' exonucleolytic degradation, such as the structure described by Eckner et al.(1991).

Oligonucleotide Probes And Primers

Polynucleotides derived from the *HKLP* gene are useful in order to detect the presence of at least a copy of a nucleotide sequence of SEQ ID Nos 1-3, or a fragment, complement, or variant thereof in a test sample.

- 5 Particularly preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 a nucleotide of SEQ ID No 1 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 1: 1-39624, 39705-40589, 40666-43629, 43710-44203, 44311-45125,
 10 45210-45440, 45622-45717, 45791-68580, 68675-70246, 70396-72421, 72601-73295, 73434-74648, 74898-83055, 83175-85192, 85279-85609, 85740-85906, 86070-88304, 88396-90585, 90705-91767, 91824-94380, 94490-96296, 96364-97184, 97270-101167, 101274-109465, 109581-110228, 110363-111819, 111882-113636, 113783-113945, 114186-117002, 117075-119676, and 119677-121162. Additional preferred probes and primers of the invention include isolated, purified,
 15 or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 2 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 2: 1-1600, 1751-2138, 2332-2539, 2659-3829 and 8885-10884.

- Additional preferred probes and primers of the invention include isolated, purified, or
 20 recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 1 or 2, or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 nucleotide positions of any one of the following ranges of nucleotide positions of:

- (a) SEQ ID No 1: 1-1000, 1001-2000, 2001-3000, 3001-4000, 4001-5000, 5001-6000,
 25 6001-7000, 7001-8000, 8001-9000, 9001-10000, 10001-11000, 11001-12000, 12001-13000, 13001-14000, 14001-15000, 15001-16000, 16001-17000, 17001-18000, 18001-19000, 19001-20000, 20001-21000, 21001-22000, 22001-23000, 23001-24000, 24001-25000, 25001-26000, 26001-27000, 27001-28000, 28001-29000, 29001-30000, 30001-31000, 31001-32000, 32001-33000, 33001-34000, 34001-35000, 35001-36000, 36001-37000, 37001-38000, 38001-39000, 39001-39624, 39705-40589, 40666-43629, 43710-44203, 44311-45125, 45210-45440, 45622-45717,
 30 45791-68580, 68675-70246, 70396-72421, 72601-73295, 73434-74648, 74898-83055, 83175-85192, 85279-85609, 85740-85906, 86070-88304, 88396-90585, 90705-91767, 91824-94380, 94490-96296, 96364-97184, 97270-101167, 101274-109465, 109581-110228, 110363-111819, 111882-113636, 113783-113945, 114186-117002, 117075-119676, and 119677-121162; and
 35 (b) SEQ ID No 2: 1-1600, 1751-2138, 2332-2539, 2659-3829 and 8885-10884.

Additional preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40,

50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 1 or the complements thereof, wherein said contiguous span comprises a G at position 7159 of SEQ ID No 1. Further preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 2 or the complements thereof, wherein said contiguous span comprises a C either at position 2551 or 4500 of SEQ ID No 2.

Another preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 3 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 3: 391-1619 and 6988-10682. Additional preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 3 or the complements thereof, wherein said contiguous span comprises a nucleotide selected in the group consisting of a C at position 5487, and a C at position 6265 of SEQ ID No 3.

Thus, the invention also relates to nucleic acid probes characterized in that they hybridize specifically, under the stringent hybridization conditions defined above, with a nucleic acid selected from the group consisting of the nucleotide sequences:

a) 1-39624, 39705-40589, 40666-43629, 43710-44203, 44311-45125, 45210-45440, 45622-45717, 45791-68580, 68675-70246, 70396-72421, 72601-73295, 73434-74648, 74898-83055, 83175-85192, 85279-85609, 85740-85906, 86070-88304, 88396-90585, 90705-91767, 91824-94380, 94490-96296, 96364-97184, 97270-101167, 101274-109465, 109581-110228, 110363-111819, 111882-113636, 113783-113945, 114186-117002, 117075-119676, and 119677-121162 of SEQ ID No 1 or a variant thereof or a sequence complementary thereto;

b) 1-1600, 1751-2138, 2332-2539, 2659-3829 and 8885-10884 of SEQ ID No 2 or a variant thereof or a sequence complementary thereto; and

c) 391-1619 and 6988-10682 of SEQ ID No 3 or a variant thereof or a sequence complementary thereto.

Additionally, another preferred embodiment of a probe according to the invention consists of a nucleic acid comprising a biallelic marker selected from the group consisting of A1 to A32 or the complements thereto, for which the receptive locations in the sequence listing are provided in Table 2.

The invention also relates to a purified and/or isolated nucleotide sequence comprising a polymorphic base of a *HKLP*-related biallelic marker, preferably of a biallelic marker selected from the group consisting of A1 to A32, and the complements thereof. The sequence has between 8 and 1000 nucleotides in length, and preferably comprises at least 8, 10, 12, 15, 18, 20, 25, 35, 40, 50, 60,

70, 80, 100, 250, 500 or 1000 contiguous nucleotides, to the extent that such lengths are consistent with the specific sequence, of a nucleotide sequence selected from the group consisting of SEQ ID Nos 1-3 and 5-8 or a variant thereof or a complementary sequence thereto. In one embodiment the invention encompasses isolated, purified, and recombinant polynucleotides comprising, consisting of, or consisting essentially of a contiguous span of 8 to 50 nucleotides of any one of SEQ ID Nos 1-3 and 5-8 and the complement thereof, wherein said span includes a *HKLP*-related biallelic marker in said sequence; optionally, wherein said *HKLP*-related biallelic marker is selected from the group consisting of A1 to A32, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, wherein said *HKLP*-related biallelic marker is selected from the group consisting of A1 to A22 and A25 to A32, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, wherein said *HKLP*-related biallelic marker is selected from the group consisting of A23 and A24, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; These nucleotide sequences comprise the polymorphic base of either allele 1 or allele 2 of the considered biallelic marker. Optionally, said biallelic marker may be within 6, 5, 4, 3, 2, or 1 nucleotides of the center of said polynucleotide or at the center of said polynucleotide; optionally, wherein said contiguous span is 18 to 35 nucleotides in length and said biallelic marker is within 4 nucleotides of the center of said polynucleotide; optionally, wherein said polynucleotide consists of said contiguous span and said contiguous span is 25 nucleotides in length and said biallelic marker is at the center of said polynucleotide; optionally, wherein the 3' end of said contiguous span is present at the 3' end of said polynucleotide; and optionally, wherein the 3' end of said contiguous span is located at the 3' end of said polynucleotide and said biallelic marker is present at the 3' end of said polynucleotide. Optionally, said polynucleotide may further comprise a label. Optionally, said polynucleotide can be attached to solid support. In a further embodiment, the polynucleotides defined above can be used alone or in any combination. In a preferred embodiment, said probes consists of, or consists essentially of a sequence selected from the following sequences: P1 to P30 and the complementary sequences thereto.

In another embodiment the invention encompasses isolated, purified and recombinant polynucleotides comprising, consisting of, or consisting essentially of a contiguous span of 8 to 50 nucleotides of SEQ ID Nos 1-3 and 5-8 or the complements thereof, wherein the 3' end of said contiguous span is located at the 3' end of said polynucleotide, and wherein the 3' end of said polynucleotide is located or at least 2, 4, 6, 8, 10, 12, 15, 18, 20, 25, 50, 100, 250, 500, or 1000 nucleotides upstream of a *HKLP*-related biallelic marker in said sequence, preferably within within 20 nucleotides upstream of a *HKLP*-related biallelic marker in said sequence; optionally, wherein said *HKLP*-related biallelic marker is selected from the group consisting of A1 to A32, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, wherein said *HKLP*-related biallelic marker is selected from the group consisting of A1

to A22 and A25 to A32, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, wherein said *HKLP*-related biallelic marker is selected from the group consisting of A23 and A24, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, wherein the 3' end of said polynucleotide
5 is located 1 nucleotide upstream of said *HKLP*-related biallelic marker in said sequence; and optionally, wherein said polynucleotide consists essentially of a sequence selected from the following sequences: D1 to D30 and E1 to E30.

In a further embodiment, the invention encompasses isolated, purified, or recombinant polynucleotides comprising, consisting of, or consisting essentially of a sequence selected from the
10 following sequences: B1 to B25 and C1 to C25.

In an additional embodiment, the invention encompasses the use of any polynucleotide for, or polynucleotides for use in determining the identity of the nucleotide at a *HKLP*-related biallelic marker or the complements thereof, as well as polynucleotides for use or use of polynucleotides in amplifying segments of nucleotides comprising a *HKLP*-related biallelic marker or the complements
15 thereof; Optionally, said determining may be performed in hybridization assay, sequencing assays, and enzyme-based mismatch detection assays; Optionally, said amplifying may be performed by a PCR or LCR. optionally, wherein said *HKLP*-related biallelic marker is selected from the group consisting of A1 to A32, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, wherein said *HKLP*-related biallelic marker is selected from
20 the group consisting of A1 to A22 and A25 to A32, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, wherein said *HKLP*-related biallelic marker is selected from the group consisting of A23 and A24, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; Optionally, said polynucleotide may be attached to a solid support, array, or addressable array; Optionally, said
25 polynucleotide may be labeled.

The invention concerns the use of the polynucleotides according to the invention for determining the identity of the nucleotide at a *HKLP*-related biallelic marker, preferably in hybridization assay, sequencing assay, microsequencing assay, or an enzyme-based mismatch detection assay and in amplifying segments of nucleotides comprising a *HKLP*-related biallelic
30 marker. In addition, the polynucleotides of the invention for use or the use of polynucleotides in determining the identity of one or more nucleotides at a *HKLP*-related biallelic marker encompass polynucleotides with any further limitation described in this disclosure, or those following, specified alone or in any combination.

The primers and probes can be prepared by any suitable method, including, for example,
35 cloning and restriction of appropriate sequences and direct chemical synthesis by a method such as the phosphodiester method of Narang et al.(1979), the phosphodiester method of Brown et al.(1979),

the diethylphosphoramidite method of Beaucage et al.(1981) and the solid support method described in EP 0 707 592. The disclosures of all these documents are incorporated herein by reference.

The formation of stable hybrids depends on the melting temperature (T_m) of the DNA. The T_m depends on the length of the primer or probe, the ionic strength of the solution and the G+C content. The higher the G+C content of the primer or probe, the higher is the melting temperature because G:C pairs are held by three H bonds whereas A:T pairs have only two. The GC content in the probes of the invention usually ranges between 10 and 75 %, preferably between 35 and 60 %, and more preferably between 40 and 55 %.

Detection probes are generally nucleic acid sequences or uncharged nucleic acid analogs such as, for example peptide nucleic acids which are disclosed in International Patent Application WO 92/20702, morpholino analogs which are described in U.S. Patents Numbered 5,185,444; 5,034,506 and 5,142,047. The probe may have to be rendered "non-extendable" in that additional dNTPs cannot be added to the probe. In and of themselves analogs usually are non-extendable and nucleic acid probes can be rendered non-extendable by modifying the 3' end of the probe such that the hydroxyl group is no longer capable of participating in elongation. For example, the 3' end of the probe can be functionalized with the capture or detection label to thereby consume or otherwise block the hydroxyl group. Alternatively, the 3' hydroxyl group simply can be cleaved, replaced or modified, U.S. Patent Application Serial No. 07/049,061 filed April 19, 1993 describes modifications, which can be used to render a probe non-extendable.

A probe or a primer according to the invention has between 8 and 1000 nucleotides in length, or is specified to be at least 12, 15, 18, 20, 25, 35, 40, 50, 60, 70, 80, 100, 250, 500 or 1000 nucleotides in length. More particularly, the length of these probes and primers can range from 8, 10, 15, 20, or 30 to 100 nucleotides, preferably from 10 to 50, more preferably from 15 to 30 nucleotides. Shorter probes and primers tend to lack specificity for a target nucleic acid sequence and generally require cooler temperatures to form sufficiently stable hybrid complexes with the template. Longer probes and primers are expensive to produce and can sometimes self-hybridize to form hairpin structures. The appropriate length for primers and probes under a particular set of assay conditions may be empirically determined by one of skill in the art. A preferred probe or primer consists of a nucleic acid comprising a polynucleotide selected from the group of the nucleotide sequences of P1 to P30 and the complementary sequence thereto, B1 to B25, C1 to C25, D1 to D30, E1 to E30, for which the respective locations in the sequence listing are provided in Tables 1, 2, 3 and 4.

Any of the polynucleotides of the present invention can be labeled, if desired, by incorporating a label detectable by spectroscopic, photochemical, biochemical, immunochemical, or chemical means. For example, useful labels include radioactive substances (^{32}P , ^{35}S , ^3H , ^{125}I), fluorescent dyes (5-bromodesoxyuridin, fluorescein, acetylaminofluorene, digoxigenin) or biotin. Preferably, polynucleotides are labeled at their 3' and 5' ends. Examples of non-radioactive labeling

of nucleic acid fragments are described in the French patent No. FR-7810975 or by Urdea et al (1988) or Sanchez-Pescador et al (1988). In addition, the probes according to the present invention may have structural characteristics such that they allow the signal amplification, such structural characteristics being, for example, branched DNA probes as those described by Urdea et al. in 1991
5 or in the European patent No. EP 0 225 807 (Chiron).

A label can also be used to capture the primer, so as to facilitate the immobilization of either the primer or a primer extension product, such as amplified DNA, on a solid support. A capture label is attached to the primers or probes and can be a specific binding member which forms a binding pair with the solid's phase reagent's specific binding member (e.g. biotin and streptavidin).
10 Therefore depending upon the type of label carried by a polynucleotide or a probe, it may be employed to capture or to detect the target DNA. Further, it will be understood that the polynucleotides, primers or probes provided herein, may, themselves, serve as the capture label. For example, in the case where a solid phase reagent's binding member is a nucleic acid sequence, it may be selected such that it binds a complementary portion of a primer or probe to thereby
15 immobilize the primer or probe to the solid phase. In cases where a polynucleotide probe itself serves as the binding member, those skilled in the art will recognize that the probe will contain a sequence or "tail" that is not complementary to the target. In the case where a polynucleotide primer itself serves as the capture label, at least a portion of the primer will be free to hybridize with a nucleic acid on a solid phase. DNA Labeling techniques are well known to the skilled technician.

20 The probes of the present invention are useful for a number of purposes. They can be notably used in Southern hybridization to genomic DNA. The probes can also be used to detect PCR amplification products. They may also be used to detect mismatches in the *HKLP* gene or mRNA using other techniques.

Any of the polynucleotides, primers and probes of the present invention can be conveniently
25 immobilized on a solid support. Solid supports are known to those skilled in the art and include the walls of wells of a reaction tray, test tubes, polystyrene beads, magnetic beads, nitrocellulose strips, membranes, microparticles such as latex particles, sheep (or other animal) red blood cells, duracytes and others. The solid support is not critical and can be selected by one skilled in the art. Thus, latex particles, microparticles, magnetic or non-magnetic beads, membranes, plastic tubes, walls of
30 microtiter wells, glass or silicon chips, sheep (or other suitable animal's) red blood cells and duracytes are all suitable examples. Suitable methods for immobilizing nucleic acids on solid phases include ionic, hydrophobic, covalent interactions and the like. A solid support, as used herein, refers to any material which is insoluble, or can be made insoluble by a subsequent reaction. The solid support can be chosen for its intrinsic ability to attract and immobilize the capture reagent.
35 Alternatively, the solid phase can retain an additional receptor which has the ability to attract and immobilize the capture reagent. The additional receptor can include a charged substance that is oppositely charged with respect to the capture reagent itself or to a charged substance conjugated to

the capture reagent. As yet another alternative, the receptor molecule can be any specific binding member which is immobilized upon (attached to) the solid support and which has the ability to immobilize the capture reagent through a specific binding reaction. The receptor molecule enables the indirect binding of the capture reagent to a solid support material before the performance of the assay or during the performance of the assay. The solid phase thus can be a plastic, derivatized plastic, magnetic or non-magnetic metal, glass or silicon surface of a test tube, microtiter well, sheet, bead, microparticle, chip, sheep (or other suitable animal's) red blood cells, duracytes® and other configurations known to those of ordinary skill in the art. The polynucleotides of the invention can be attached to or immobilized on a solid support individually or in groups of at least 2, 5, 8, 10, 12, 15, 20, or 25 distinct polynucleotides of the invention to a single solid support. In addition, polynucleotides other than those of the invention may be attached to the same solid support as one or more polynucleotides of the invention.

Consequently, the invention also deals with a method for detecting the presence of a nucleic acid comprising a nucleotide sequence selected from a group consisting of SEQ ID Nos 1-3, a fragment or a variant thereof and a complementary sequence thereto in a sample, said method comprising the following steps of:

a) bringing into contact a nucleic acid probe or a plurality of nucleic acid probes which can hybridize with a nucleotide sequence included in a nucleic acid selected from the group consisting of the nucleotide sequences of SEQ ID Nos 1-3, a fragment or a variant thereof and a complementary sequence thereto and the sample to be assayed; and

b) detecting the hybrid complex formed between the probe and a nucleic acid in the sample.

The invention further concerns a kit for detecting the presence of a nucleic acid comprising a nucleotide sequence selected from a group consisting of SEQ ID Nos 1-3, a fragment or a variant thereof and a complementary sequence thereto in a sample, said kit comprising:

a) a nucleic acid probe or a plurality of nucleic acid probes which can hybridize with a nucleotide sequence included in a nucleic acid selected from the group consisting of the nucleotide sequences of SEQ ID Nos 1-3, a fragment or a variant thereof and a complementary sequence thereto; and

b) optionally, the reagents necessary for performing the hybridization reaction.

In a first preferred embodiment of this detection method and kit, said nucleic acid probe or the plurality of nucleic acid probes are labeled with a detectable molecule. In a second preferred embodiment of said method and kit, said nucleic acid probe or the plurality of nucleic acid probes has been immobilized on a substrate. In a third preferred embodiment, the nucleic acid probe or the plurality of nucleic acid probes comprise either a sequence which is selected from the group consisting of the nucleotide sequences of P1 to P30 and the complementary sequence thereto, B1 to B25, C1 to C25, D1 to D30, E1 to E30 or a biallelic marker selected from the group consisting of A1 to A32 and the complements thereto.

Olig nucleotide Arrays

A substrate comprising a plurality of oligonucleotide primers or probes of the invention may be used either for detecting or amplifying targeted sequences in the *HKLP* gene and may also be used for detecting mutations in the coding or in the non-coding sequences of the *HKLP* gene.

5 Any polynucleotide provided herein may be attached in overlapping areas or at random locations on the solid support. Alternatively the polynucleotides of the invention may be attached in an ordered array wherein each polynucleotide is attached to a distinct region of the solid support which does not overlap with the attachment site of any other polynucleotide. Preferably, such an ordered array of polynucleotides is designed to be "addressable" where the distinct locations are
10 recorded and can be accessed as part of an assay procedure. Addressable polynucleotide arrays typically comprise a plurality of different oligonucleotide probes that are coupled to a surface of a substrate in different known locations. The knowledge of the precise location of each polynucleotides location makes these "addressable" arrays particularly useful in hybridization assays. Any addressable array technology known in the art can be employed with the
15 polynucleotides of the invention. One particular embodiment of these polynucleotide arrays is known as the Genechips™, and has been generally described in US Patent 5,143,854; PCT publications WO 90/15070 and 92/10092. These arrays may generally be produced using mechanical synthesis methods or light directed synthesis methods which incorporate a combination of photolithographic methods and solid phase oligonucleotide synthesis (Fodor et al., 1991). The
20 immobilization of arrays of oligonucleotides on solid supports has been rendered possible by the development of a technology generally identified as "Very Large Scale Immobilized Polymer Synthesis" (VLSIPS™) in which, typically, probes are immobilized in a high density array on a solid surface of a chip. Examples of VLSIPS™ technologies are provided in US Patents 5,143,854; and 5,412,087 and in PCT Publications WO 90/15070, WO 92/10092 and WO 95/11995, which
25 describe methods for forming oligonucleotide arrays through techniques such as light-directed synthesis techniques. In designing strategies aimed at providing arrays of nucleotides immobilized on solid supports, further presentation strategies were developed to order and display the oligonucleotide arrays on the chips in an attempt to maximize hybridization patterns and sequence information. Examples of such presentation strategies are disclosed in PCT Publications WO
30 94/12305, WO 94/11530, WO 97/29212 and WO 97/31256.

In another embodiment of the oligonucleotide arrays of the invention, an oligonucleotide probe matrix may advantageously be used to detect mutations occurring in the *HKLP* gene and in its regulatory region. For this particular purpose, probes are specifically designed to have a nucleotide sequence allowing their hybridization to the genes that carry known mutations (either by deletion,
35 insertion or substitution of one or several nucleotides). By known mutations, it is meant, mutations on the *HKLP* gene that have been identified according, for example to the technique used by Huang et al.(1996) or Samson et al.(1996).

Another technique that is used to detect mutations in the *HKLP* gene is the use of a high-density DNA array. Each oligonucleotide probe constituting a unit element of the high density DNA array is designed to match a specific subsequence of the *HKLP* genomic DNA or cDNA. Thus, an array consisting of oligonucleotides complementary to subsequences of the target gene sequence is used to determine the identity of the target sequence with the wild gene sequence, measure its amount, and detect differences between the target sequence and the reference wild gene sequence of the *HKLP* gene. In one such design, termed 4L tiled array, is implemented a set of four probes (A, C, G, T), preferably 15-nucleotide oligomers. In each set of four probes, the perfect complement will hybridize more strongly than mismatched probes. Consequently, a nucleic acid target of length L is scanned for mutations with a tiled array containing 4L probes, the whole probe set containing all the possible mutations in the known wild reference sequence. The hybridization signals of the 15-mer probe set tiled array are perturbed by a single base change in the target sequence. As a consequence, there is a characteristic loss of signal or a "footprint" for the probes flanking a mutation position. This technique was described by Chee et al. in 1996, which is herein incorporated by reference.

Consequently, the invention concerns an array of nucleic acid molecules comprising at least one polynucleotide described above as probes and primers. Preferably, the invention concerns an array of nucleic acid comprising at least two polynucleotides described above as probes and primers.

A further object of the invention consists of an array of nucleic acid sequences comprising either at least one of the sequences selected from the group consisting of P1 to P30, B1 to B25, C1 to C25, D1 to D30, E1 to E30, the sequences complementary thereto, a fragment thereof of at least 8 consecutive nucleotides thereof, and at least one sequence comprising a biallelic marker selected from the group consisting of A1 to A32 and the complements thereto.

The invention also pertains to an array of nucleic acid sequences comprising either at least two of the sequences selected from the group consisting of P1 to P30, B1 to B25, C1 to C25, D1 to D30, E1 to E30, the sequences complementary thereto, a fragment thereof of at least 8 consecutive nucleotides thereof, and at least two sequences comprising a biallelic marker selected from the group consisting of A1 to A32 and the complements thereof.

Amplification of the *HKLP* gene.

1. DNA extraction

As for the source of the genomic DNA to be subjected to analysis, any test sample can be foreseen without any particular limitation. These test samples include biological samples which can be tested by the methods of the present invention described herein and include human and animal body fluids such as whole blood, serum, plasma, cerebrospinal fluid, urine, lymph fluids, and various external secretions of the respiratory, intestinal and genitourinary tracts, tears, saliva, milk, white blood cells, myelomas and the like; biological fluids such as cell culture supernatants; fixed

tissue specimens including tumor and non-tumor tissue and lymph node tissues; bone marrow aspirates and fixed cell specimens. The preferred source of genomic DNA used in the context of the present invention is from peripheral venous blood of each donor.

The techniques of DNA extraction are well-known to the skilled technician. Such
5 techniques are described notably by Mackey et al. (1998).

2. DNA amplification

DNA amplification techniques are well-known to those skilled in the art. Amplification techniques that can be used in the context of the present invention include, but are not limited to, the ligase chain reaction (LCR) described in EP-A- 320 308, WO 9320227 and EP-A-439 182, the
10 disclosures of which are incorporated herein by reference, the polymerase chain reaction (PCR, RT-PCR) and techniques such as the nucleic acid sequence based amplification (NASBA) described in Guatelli JC, et al. (1990) and in Compton J. (1991), Q-beta amplification as described in European Patent Application no 4544610, strand displacement amplification as described in Walker et al. (1996) and EP A 684 315 and, target mediated amplification as described in PCT Publication WO
15 9322461, the disclosure of which is incorporated herein by reference.

LCR and Gap LCR are exponential amplification techniques, both depend on DNA ligase to join adjacent primers annealed to a DNA molecule. In Ligase Chain Reaction (LCR), probe pairs are used which include two primary (first and second) and two secondary (third and fourth) probes, all of which are employed in molar excess to target. The first probe hybridizes to a first segment of the
20 target strand and the second probe hybridizes to a second segment of the target strand, the first and second segments being contiguous so that the primary probes abut one another in 5' phosphate-3'hydroxyl relationship, and so that a ligase can covalently fuse or ligate the two probes into a fused product. In addition, a third (secondary) probe can hybridize to a portion of the first probe and a fourth (secondary) probe can hybridize to a portion of the second probe in a similar abutting fashion.
25 Of course, if the target is initially double stranded, the secondary probes also will hybridize to the target complement in the first instance. Once the ligated strand of primary probes is separated from the target strand, it will hybridize with the third and fourth probes which can be ligated to form a complementary, secondary ligated product. It is important to realize that the ligated products are functionally equivalent to either the target or its complement. By repeated cycles of hybridization
30 and ligation, amplification of the target sequence is achieved. A method for multiplex LCR has also been described (WO 9320227). Gap LCR (GLCR) is a version of LCR where the probes are not adjacent but are separated by 2 to 3 bases.

For amplification of mRNAs, it is within the scope of the present invention to reverse transcribe mRNA into cDNA followed by polymerase chain reaction (RT-PCR); or, to use a single
35 enzyme for both steps as described in U.S. Patent No. 5,322,770 or, to use Asymmetric Gap LCR

(RT-AGLCR) as described by Marshall et al. (1994). AGLCR is a modification of GLCR that allows the amplification of RNA.

The PCR technology is the preferred amplification technique used in the present invention. A variety of PCR techniques are familiar to those skilled in the art. For a review of PCR
5 technology, see White (1997) and the publication entitled "PCR Methods and Applications" (1991, Cold Spring Harbor Laboratory Press). In each of these PCR procedures, PCR primers on either side of the nucleic acid sequences to be amplified are added to a suitably prepared nucleic acid sample along with dNTPs and a thermostable polymerase such as Taq polymerase, Pfu polymerase, or Vent polymerase. The nucleic acid in the sample is denatured and the PCR primers are
10 specifically hybridized to complementary nucleic acid sequences in the sample. The hybridized primers are extended. Thereafter, another cycle of denaturation, hybridization, and extension is initiated. The cycles are repeated multiple times to produce an amplified fragment containing the nucleic acid sequence between the primer sites. PCR has further been described in several patents including US Patents 4,683,195, 4,683,202 and 4,965,188. Each of these publications is
15 incorporated by reference.

One of the aspects of the present invention is a method for the amplification of the human *HKLP* gene, particularly of the genomic sequences of SEQ ID No 1 and 2 or of the cDNA sequence of SEQ ID No 3, or a fragment or a variant thereof in a test sample, preferably using the PCR technology. The method comprises the steps of contacting a test sample suspected of containing the
20 target *HKLP* encoding sequence or portion thereof with amplification reaction reagents comprising a pair of amplification primers, and eventually in some instances a detection probe that can hybridize with an internal region of amplicon sequences to confirm that the desired amplification reaction has taken place.

Thus, the present invention also relates to a method for the amplification of a human *HKLP*
25 gene sequence, particularly of a portion of the genomic sequences of SEQ ID Nos 1 and 2 or of the cDNA sequence of SEQ ID No 3, or a variant thereof in a test sample, said method comprising the steps of:

a) contacting a test sample suspected of containing the targeted *HKLP* gene sequence comprised in a nucleotide sequence selected from a group consisting of SEQ ID Nos 1-3, or
30 fragments or variants thereof with amplification reaction reagents comprising a pair of amplification primers as described above and located on either side of the polynucleotide region to be amplified; and

b) optionally, detecting the amplification products.

The invention also concerns a kit for the amplification of a human *HKLP* gene sequence,
35 particularly of a portion of the genomic sequences of SEQ ID No 1 and 2 or of the cDNA sequence of SEQ ID No 3, or a variant thereof in a test sample, wherein said kit comprises:

a) a pair of oligonucleotide primers located on either side of the *HKLP* region to be amplified; and

b) Optionally, the reagents necessary for performing the amplification reaction.

In a first preferred embodiment of the above amplification method or kit, the amplification
5 product is detected by hybridization with a labeled probe having a sequence which is complementary to the amplified region.

The primers are more particularly characterized in that they have sufficient complementarity with any sequence of a strand of the genomic sequence close to the region to be amplified, for example with a non-coding sequence adjacent to exons to amplify.

10 In a second preferred embodiment, the nucleic acid primers comprise a sequence which is selected from the group consisting of the nucleotide sequences of B1 to B25, C1 to C25, D1 to D30, and E1 to E30.

HKLP Proteins and Polypeptide Fragments:

The term "HKLP polypeptides" is used herein to embrace all of the proteins and
15 polypeptides of the present invention. Also forming part of the invention are polypeptides encoded by the polynucleotides of the invention, as well as fusion polypeptides comprising such polypeptides. The invention embodies HKLP proteins from humans, including isolated or purified HKLP proteins consisting, consisting essentially, or comprising the sequence of SEQ ID No 4. The HKLP protein has 1816 amino acids in length. The 700 first amino acids of the HKLP protein
20 present 97 % of homology with the murine KIF1B protein (Nangaku et al., 1994). The HKLP protein presents 60-70% of homology with the murine KIF1A protein, and more particularly the 390 first amino acids of the HKLP protein have 85 % of homology therewith.

The present invention embodies isolated, purified, and recombinant polypeptides comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably
25 at least 12, 15, 20, 25, 30, 40, 50, or 100 amino acids of SEQ ID No 4, wherein said contiguous span includes at least 1, 2, 3, 5 or 10 of the amino acid positions 1-478 of the SEQ ID No 4. In other preferred embodiments the contiguous stretch of amino acids comprises the site of a mutation or functional mutation, including a deletion, addition, swap or truncation of the amino acids in the HKLP protein sequence.

30 The invention also encompasses a purified, isolated, or recombinant polynucleotides comprising a nucleotide sequence having at least 70, 75, 80, 85, 90, 95, 98 or 99% nucleotide identity with a nucleotide sequence of SEQ ID No 4 or a complementary sequence thereto or a fragment thereof.

HKLP proteins are preferably isolated from human or mammalian tissue samples or
35 expressed from human or mammalian genes. The HKLP polypeptides of the invention can be made using routine expression methods known in the art. The polynucleotide encoding the desired

polypeptide is ligated into an expression vector suitable for any convenient host. Both eukaryotic and prokaryotic host systems is used in forming recombinant polypeptides, and a summary of some of the more common systems. The polypeptide is then isolated from lysed cells or from the culture medium and purified to the extent needed for its intended use. Purification is by any technique
5 known in the art, for example, differential extraction, salt fractionation, chromatography, centrifugation, and the like. See, for example, Methods in Enzymology for a variety of methods for purifying proteins.

In addition, shorter protein fragments is produced by chemical synthesis. Alternatively the proteins of the invention is extracted from cells or tissues of humans or non-human animals.
10 Methods for purifying proteins are known in the art, and include the use of detergents or chaotropic agents to disrupt particles followed by differential extraction and separation of the polypeptides by ion exchange chromatography, affinity chromatography, sedimentation according to density, and gel electrophoresis.

Any *HKLP* cDNA, including SEQ ID No 3, is used to express *HKLP* proteins and
15 polypeptides. The nucleic acid encoding the *HKLP* protein or polypeptide to be expressed is operably linked to a promoter in an expression vector using conventional cloning technology. The *HKLP* insert in the expression vector may comprise the full coding sequence for the *HKLP* protein or a portion thereof. For example, the *HKLP* derived insert may encode a polypeptide comprising at least 10 consecutive amino acids of the *HKLP* protein of SEQ ID No 4, wherein said contiguous span includes
20 at least 1, 2, 3, 5 or 10 of the amino acid positions 1-478 of the SEQ ID No 4.

The expression vector is any of the mammalian, yeast, insect or bacterial expression systems known in the art. Commercially available vectors and expression systems are available from a variety of suppliers including Genetics Institute (Cambridge, MA), Stratagene (La Jolla, California), Promega (Madison, Wisconsin), and Invitrogen (San Diego, California). If desired, to enhance expression and
25 facilitate proper protein folding, the codon context and codon pairing of the sequence is optimized for the particular expression organism in which the expression vector is introduced, as explained by Hatfield, et al., U.S. Patent No. 5,082,767.

In one embodiment, the entire coding sequence of the *HKLP* cDNA through the poly A signal of the cDNA are operably linked to a promoter in the expression vector. Alternatively, if the nucleic
30 acid encoding a portion of the *HKLP* protein lacks a methionine to serve as the initiation site, an initiating methionine can be introduced next to the first codon of the nucleic acid using conventional techniques. Similarly, if the insert from the *HKLP* cDNA lacks a poly A signal, this sequence can be added to the construct by, for example, splicing out the Poly A signal from pSG5 (Stratagene) using *Bgl*I and *Sall* restriction endonuclease enzymes and incorporating it into the mammalian expression
35 vector pXT1 (Stratagene). pXT1 contains the LTRs and a portion of the *gag* gene from Moloney Murine Leukemia Virus. The position of the LTRs in the construct allow efficient stable transfection. The vector includes the Herpes Simplex Thymidine Kinase promoter and the selectable neomycin gene.

The nucleic acid encoding the HKLP protein or a portion thereof is obtained by PCR from a bacterial vector containing the *HKLP* cDNA of SEQ ID No 3 using oligonucleotide primers complementary to the *HKLP* cDNA or portion thereof and containing restriction endonuclease sequences for Pst I incorporated into the 5' primer and BglII at the 5' end of the corresponding cDNA 3' primer, taking care
5 to ensure that the sequence encoding the HKLP protein or a portion thereof is positioned properly with respect to the poly A signal. The purified fragment obtained from the resulting PCR reaction is digested with PstI, blunt ended with an exonuclease, digested with Bgl II, purified and ligated to pXT1, now containing a poly A signal and digested with BglII.

The ligated product is transfected into mouse NIH 3T3 cells using Lipofectin (Life
10 Technologies, Inc., Grand Island, New York) under conditions outlined in the product specification. Positive transfectants are selected after growing the transfected cells in 600ug/ml G418 (Sigma, St. Louis, Missouri).

Alternatively, the nucleic acids encoding the HKLP protein or a portion thereof is cloned into pED6dpc2 (Genetics Institute, Cambridge, MA). The resulting pED6dpc2 constructs is transfected into
15 a suitable host cell, such as COS 1 cells. Methotrexate resistant cells are selected and expanded.

The above procedures may also be used to express a mutant HKLP protein responsible for a detectable phenotype or a portion thereof.

The expressed proteins are purified using conventional purification techniques such as ammonium sulfate precipitation or chromatographic separation based on size or charge. The protein
20 encoded by the nucleic acid insert may also be purified using standard immunochromatography techniques. In such procedures, a solution containing the expressed HKLP protein or portion thereof, such as a cell extract, is applied to a column having antibodies against the HKLP protein or portion thereof is attached to the chromatography matrix. The expressed protein is allowed to bind the immunochromatography column. Thereafter, the column is washed to remove non-specifically bound
25 proteins. The specifically bound expressed protein is then released from the column and recovered using standard techniques.

To confirm expression of the HKLP protein or a portion thereof, the proteins expressed from host cells containing an expression vector containing an insert encoding the HKLP protein or a portion thereof can be compared to the proteins expressed in host cells containing the expression vector without
30 an insert. The presence of a band in samples from cells containing the expression vector with an insert which is absent in samples from cells containing the expression vector without an insert indicates that the HKLP protein or a portion thereof is being expressed. Generally, the band will have the mobility expected for the HKLP protein or portion thereof. However, the band may have a mobility different than that expected as a result of modifications such as glycosylation, ubiquitination, or enzymatic
35 cleavage.

Antibodies capable of specifically recognizing the expressed HKLP protein or a portion thereof are described below.

If antibody production is not possible, the nucleic acids encoding the HKLP protein or a portion thereof is incorporated into expression vectors designed for use in purification schemes employing chimeric polypeptides. In such strategies the nucleic acid encoding the HKLP protein or a portion thereof is inserted in frame with the gene encoding the other half of the chimera. The other half of the chimera is β -globin or a nickel binding polypeptide encoding sequence. A chromatography matrix having antibody to β -globin or nickel attached thereto is then used to purify the chimeric protein. Protease cleavage sites is engineered between the β -globin gene or the nickel binding polypeptide and the HKLP protein or portion thereof. Thus, the two polypeptides of the chimera is separated from one another by protease digestion.

One useful expression vector for generating β -globin chimerics is pSG5 (Stratagene), which encodes rabbit β -globin. Intron II of the rabbit β -globin gene facilitates splicing of the expressed transcript, and the polyadenylation signal incorporated into the construct increases the level of expression. These techniques are well known to those skilled in the art of molecular biology. Standard methods are published in methods texts such as Davis et al., (1986) and many of the methods are available from Stratagene, Life Technologies, Inc., or Promega. Polypeptide may additionally be produced from the construct using in vitro translation systems such as the In vitro Express™ Translation Kit (Stratagene).

Thus, the present invention also concerns a method for producing one of the polypeptides described herein, and especially a polypeptide of SEQ ID No 4 or a fragment or a variant thereof, wherein said method comprises the steps of:

- a) culturing, in an appropriate culture medium, a cell host previously transformed or transfected with the recombinant vector comprising a nucleic acid encoding a HKLP polypeptide, or a fragment or a variant thereof;
- b) harvesting the culture medium thus conditioned or lyse the cell host, for example by sonication or by an osmotic shock;
- c) separating or purifying, from the said culture medium, or from the pellet of the resultant host cell lysate the thus produced polypeptide of interest.
- d) Optionally characterizing the produced polypeptide of interest.

In a specific embodiment of the above method, step a) is preceded by a step wherein the nucleic acid coding for a HKLP polypeptide, or a fragment or a variant thereof, is inserted in an appropriate vector, optionally after an appropriate cleavage of this amplified nucleic acid with one or several restriction endonucleases. The nucleic acid coding for a HKLP polypeptide or a fragment or a variant thereof may be the resulting product of an amplification reaction using a pair of primers according to the invention (by SDA, TAS, 3SR NASBA, TMA etc.).

Antibodies That Bind HKLP Polypeptides of the Invention

Any HKLP polypeptide or whole protein may be used to generate antibodies capable of specifically binding to expressed HKLP protein or fragments thereof as described. The antibody compositions of the invention are capable of specifically binding or specifically bind to the HKLP protein. For an antibody composition to specifically bind to the HKLP protein it must demonstrate at least a 5%, 10%, 15%, 20%, 25%, 50%, or 100% greater binding affinity for full length HKLP protein than for any full length protein in an ELISA, RIA, or other antibody-based binding assay.

In a preferred embodiment of the invention antibody compositions are capable of selectively binding, or selectively bind to an epitope-containing fragment of a polypeptide comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, or 100 amino acids of SEQ ID No 4, wherein said epitope comprises at least 1, 2, 3, 5 or 10 of the amino acid positions 1-478 of the SEQ ID No 4, wherein said antibody composition is optionally either polyclonal or monoclonal.

The present invention also contemplates the use of polypeptides comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 50, or 100 amino acids of a HKLP polypeptide in the manufacture of antibodies, wherein said contiguous span comprises at least 1, 2, 3, 5 or 10 of the amino acid positions 1-478 of the SEQ ID No 4. In a preferred embodiment such polypeptides are useful in the manufacture of antibodies to detect the presence and absence of the HKLP protein.

Non-human animals or mammals, whether wild-type or transgenic, which express a different species of HKLP than the one to which antibody binding is desired, and animals which do not express HKLP (i.e. a HKLP knock out animal as described in herein) are particularly useful for preparing antibodies. HKLP knock out animals will recognize all or most of the exposed regions of HKLP as foreign antigens, and therefore produce antibodies with a wider array of HKLP epitopes. Moreover, smaller polypeptides with only 10 to 30 amino acids may be useful in obtaining specific binding to the HKLP protein. In addition, the humoral immune system of animals which produce a species of HKLP that resembles the antigenic sequence will preferentially recognize the differences between the animal's native HKLP species and the antigen sequence, and produce antibodies to these unique sites in the antigen sequence. Such a technique will be particularly useful in obtaining antibodies that specifically bind to the HKLP protein.

Antibody preparations prepared according to either protocol are useful in quantitative immunoassays which determine concentrations of antigen-bearing substances in biological samples; they are also used semi-quantitatively or qualitatively to identify the presence of antigen in a biological sample. The antibodies may also be used in therapeutic compositions for killing cells expressing the protein or reducing the levels of the protein in the body.

The antibodies of the invention may be labeled, either by a radioactive, a fluorescent or an enzymatic label.

Consequently, the invention is also directed to a method for detecting specifically the presence of a human HKLP polypeptide according to the invention in a biological sample, said method comprising the following steps :

- a) bringing into contact the biological sample with a polyclonal or monoclonal antibody directed against the HKLP polypeptide of the amino acid sequence of SEQ ID No 4, or to a peptide fragment or variant thereof;
- b) detecting the antigen-antibody complex formed.

The invention also concerns a diagnostic kit for detecting *in vitro* the presence of a human HKLP polypeptide according to the present invention in a biological sample, wherein said kit comprises :

- a) a polyclonal or monoclonal antibody directed against the HKLP polypeptide of the amino acid sequence of SEQ ID No 4, or to a peptide fragment or variant thereof, optionally labeled;
- b) a reagent allowing the detection of the antigen-antibody complexes formed, said reagent carrying optionally a label, or being able to be recognized itself by a labeled reagent, more particularly in the case when the above-mentioned monoclonal or polyclonal antibody is not labeled by itself.

HKLP-related Biallelic Markers

Advantages Of The Biallelic Markers Of The Present Invention

The HKLP-related biallelic markers of the present invention offer a number of important advantages over other genetic markers such as RFLP (Restriction fragment length polymorphism) and VNTR (Variable Number of Tandem Repeats) markers.

The first generation of markers, were RFLPs, which are variations that modify the length of a restriction fragment. But methods used to identify and to type RFLPs are relatively wasteful of materials, effort, and time. The second generation of genetic markers were VNTRs, which can be categorized as either minisatellites or microsatellites. Minisatellites are tandemly repeated DNA sequences present in units of 5-50 repeats which are distributed along regions of the human chromosomes ranging from 0.1 to 20 kilobases in length. Since they present many possible alleles, their informative content is very high. Minisatellites are scored by performing Southern blots to identify the number of tandem repeats present in a nucleic acid sample from the individual being tested. However, there are only 10^4 potential VNTRs that can be typed by Southern blotting. Moreover, both RFLP and VNTR markers are costly and time-consuming to develop and assay in large numbers.

Single nucleotide polymorphism or biallelic markers can be used in the same manner as RFLPs and VNTRs but offer several advantages. SNP are densely spaced in the human genome and represent the most frequent type of variation. An estimated number of more than 10^7 sites are

scattered along the 3×10^9 base pairs of the human genome. Therefore, SNP occur at a greater frequency and with greater uniformity than RFLP or VNTR markers which means that there is a greater probability that such a marker will be found in close proximity to a genetic locus of interest. SNP are less variable than VNTR markers but are mutationally more stable.

5 Also, the different forms of a characterized single nucleotide polymorphism, such as the biallelic markers of the present invention, are often easier to distinguish and can therefore be typed easily on a routine basis. Biallelic markers have single nucleotide based alleles and they have only two common alleles, which allows highly parallel detection and automated scoring. The biallelic markers of the present invention offer the possibility of rapid, high throughput genotyping of a large
10 number of individuals.

Biallelic markers are densely spaced in the genome, sufficiently informative and can be assayed in large numbers. The combined effects of these advantages make biallelic markers extremely valuable in genetic studies. Biallelic markers can be used in linkage studies in families, in allele sharing methods, in linkage disequilibrium studies in populations, in association studies of
15 case-control populations or of trait positive and trait negative populations. An important aspect of the present invention is that biallelic markers allow association studies to be performed to identify genes involved in complex traits. Association studies examine the frequency of marker alleles in unrelated case- and control-populations and are generally employed in the detection of polygenic or sporadic traits. Association studies may be conducted within the general population and are not
20 limited to studies performed on related individuals in affected families (linkage studies). Biallelic markers in different genes can be screened in parallel for direct association with disease or response to a treatment. This multiple gene approach is a powerful tool for a variety of human genetic studies as it provides the necessary statistical power to examine the synergistic effect of multiple genetic factors on a particular phenotype, drug response, sporadic trait, or disease state with a complex
25 genetic etiology.

Candidate Gene Of The Present Invention

Different approaches can be employed to perform association studies: genome-wide association studies, candidate region association studies and candidate gene association studies. Genome-wide association studies rely on the screening of genetic markers evenly spaced and
30 covering the entire genome. The candidate gene approach is based on the study of genetic markers specifically located in genes potentially involved in a biological pathway related to the trait of interest. In the present invention, *HKLP* is the candidate gene. The candidate gene analysis clearly provides a short-cut approach to the identification of genes and gene polymorphisms related to a particular trait when some information concerning the biology of the trait is available. However, it
35 should be noted that all of the biallelic markers disclosed in the instant application can be employed

as part of genome-wide association studies or as part of candidate region association studies and such uses are specifically contemplated in the present invention and claims.

***HKLP*-Related Biallelic Markers And Polynucleotides Related Thereto**

The invention also concerns *HKLP*-related biallelic markers. As used herein the term
5 “*HKLP*-related biallelic marker” relates to a set of biallelic markers in linkage disequilibrium with the *HKLP* gene. The term *HKLP*-related biallelic marker includes the biallelic markers designated A1 to A32.

A portion of the biallelic markers of the present invention are disclosed in Table 2. Their location on the *HKLP* gene is indicated in Table 2 and also as a single base polymorphism in the
10 features of in the related SEQ ID Nos 1-3 and 5-8. The pairs of primers allowing the amplification of a nucleic acid containing the polymorphic base of one *HKLP* biallelic marker are listed in Table 1 of Example 2.

27 *HKLP*-related biallelic markers, A1 to A27, are located in the genomic sequence of *HKLP*. Four of them are located in exonic sequence, namely A1, A23, A24 and A25. The other
15 *HKLP*-related biallelic markers are located in intronic region of *HKLP*. Additionally, 5 biallelic markers are located in intergenic region and are in linkage disequilibrium with the *HKLP* gene.

The primers for amplification or sequencing reaction of a polynucleotide comprising a biallelic marker of the invention may be designed from the disclosed sequences for any method known in the art. A preferred set of primers are fashioned such that the 3' end of the contiguous
20 span of identity with a sequence selected from the group consisting of SEQ ID Nos 1-3 and 5-8 or a sequence complementary thereto or a variant thereof is present at the 3' end of the primer. Such a configuration allows the 3' end of the primer to hybridize to a selected nucleic acid sequence and dramatically increases the efficiency of the primer for amplification or sequencing reactions. Allele specific primers may be designed such that a polymorphic base of a biallelic marker is at the 3' end
25 of the contiguous span and the contiguous span is present at the 3' end of the primer. Such allele specific primers tend to selectively prime an amplification or sequencing reaction so long as they are used with a nucleic acid sample that contains one of the two alleles present at a biallelic marker. The 3' end of the primer of the invention may be located within or at least 2, 4, 6, 8, 10, 12, 15, 18, 20, 25, 50, 100, 250, 500, or 1000 nucleotides upstream of a *HKLP*-related biallelic marker in said
30 sequence or at any other location which is appropriate for their intended use in sequencing, amplification or the location of novel sequences or markers. Thus, another set of preferred amplification primers comprise an isolated polynucleotide consisting essentially of a contiguous span of 8 to 50 nucleotides in a sequence selected from the group consisting of SEQ ID Nos 1-3 and 5-8 or a sequence complementary thereto or a variant thereof, wherein the 3' end of said contiguous
35 span is located at the 3' end of said polynucleotide, and wherein the 3' end of said polynucleotide is located upstream of a *HKLP*-related biallelic marker in said sequence. Preferably, those

amplification primers comprise a sequence selected from the group consisting of the sequences B1 to B25 and C1 to C25. Primers with their 3' ends located 1 nucleotide upstream of a *HKLP*-related biallelic marker have a special utility in microsequencing assays. Preferred microsequencing primers are described in Table 4. Optionally, microsequencing primers are selected from the group
5 consisting of the nucleotide sequences D1 to D30 and E1 to E30.

The probes of the present invention may be designed from the disclosed sequences for any method known in the art, particularly methods which allow for testing if a marker disclosed herein is present. A preferred set of probes may be designed for use in the hybridization assays of the invention in any manner known in the art such that they selectively bind to one allele of a biallelic
10 marker, but not the other allele under any particular set of assay conditions. Preferred hybridization probes comprise the polymorphic base of either allele 1 or allele 2 of the specific biallelic marker. Optionally, said biallelic marker may be within 6, 5, 4, 3, 2, or 1 nucleotides of the center of the hybridization probe or at the center of said probe.

It should be noted that the polynucleotides of the present invention are not limited to having
15 the exact flanking sequences surrounding the polymorphic bases which are enumerated in Sequence Listing. Rather, it will be appreciated that the flanking sequences surrounding the biallelic markers may be lengthened or shortened to any extent compatible with their intended use and the present invention specifically contemplates such sequences. The flanking regions outside of the contiguous span need not be homologous to native flanking sequences which actually occur in human subjects.
20 The addition of any nucleotide sequence which is compatible with the nucleotides intended use is specifically contemplated.

Primers and probes may be labeled or immobilized on a solid support as described in "Oligonucleotide probes and primers".

The polynucleotides of the invention which are attached to a solid support encompass
25 polynucleotides with any further limitation described in this disclosure, or those following, specified alone or in any combination: Optionally, said polynucleotides may be specified as attached individually or in groups of at least 2, 5, 8, 10, 12, 15, 20, or 25 distinct polynucleotides of the invention to a single solid support. Optionally, polynucleotides other than those of the invention may attached to the same solid support as polynucleotides of the invention. Optionally, when
30 multiple polynucleotides are attached to a solid support they may be attached at random locations, or in an ordered array. Optionally, said ordered array may be addressable.

The present invention also encompasses diagnostic kits comprising one or more polynucleotides of the invention with a portion or all of the necessary reagents and instructions for genotyping a test subject by determining the identity of a nucleotide at a *HKLP*-related biallelic
35 marker. The polynucleotides of a kit may optionally be attached to a solid support, or be part of an array or addressable array of polynucleotides. The kit may provide for the determination of the identity of the nucleotide at a marker position by any method known in the art including, but not

limited to, a sequencing assay method, a microsequencing assay method, a hybridization assay method, or an enzyme-based mismatch detection method.

Methods For De Novo Identification Of Biallelic Markers

Any of a variety of methods can be used to screen a genomic fragment for single nucleotide polymorphisms such as differential hybridization with oligonucleotide probes, detection of changes in the mobility measured by gel electrophoresis or direct sequencing of the amplified nucleic acid. A preferred method for identifying biallelic markers involves comparative sequencing of genomic DNA fragments from an appropriate number of unrelated individuals.

In a first embodiment, DNA samples from unrelated individuals are pooled together, following which the genomic DNA of interest is amplified and sequenced. The nucleotide sequences thus obtained are then analyzed to identify significant polymorphisms. One of the major advantages of this method resides in the fact that the pooling of the DNA samples substantially reduces the number of DNA amplification reactions and sequencing reactions, which must be carried out. Moreover, this method is sufficiently sensitive so that a biallelic marker obtained thereby usually demonstrates a sufficient frequency of its less common allele to be useful in conducting association studies.

In a second embodiment, the DNA samples are not pooled and are therefore amplified and sequenced individually. This method is usually preferred when biallelic markers need to be identified in order to perform association studies within candidate genes. Preferably, highly relevant gene regions such as promoter regions or exon regions may be screened for biallelic markers. A biallelic marker obtained using this method may show a lower degree of informativeness for conducting association studies, e.g. if the frequency of its less frequent allele may be less than about 10%. Such a biallelic marker will, however, be sufficiently informative to conduct association studies and it will further be appreciated that including less informative biallelic markers in the genetic analysis studies of the present invention, may allow in some cases the direct identification of causal mutations, which may, depending on their penetrance, be rare mutations.

The following is a description of the various parameters of a preferred method used by the inventors for the identification of the biallelic markers of the present invention.

Genomic DNA Samples

The genomic DNA samples from which the biallelic markers of the present invention are generated are preferably obtained from unrelated individuals corresponding to a heterogeneous population of known ethnic background. The number of individuals from whom DNA samples are obtained can vary substantially, preferably from about 10 to about 1000, preferably from about 50 to about 200 individuals. It is usually preferred to collect DNA samples from at least about 100 individuals in order to have sufficient polymorphic diversity in a given population to identify as many markers as possible and to generate statistically significant results.

As for the source of the genomic DNA to be subjected to analysis, any test sample can be foreseen without any particular limitation. These test samples include biological samples, which can be tested by the methods of the present invention described herein, and include human and animal body fluids such as whole blood, serum, plasma, cerebrospinal fluid, urine, lymph fluids, and
5 various external secretions of the respiratory, intestinal and genitourinary tracts, tears, saliva, milk, white blood cells, myelomas and the like; biological fluids such as cell culture supernatants; fixed tissue specimens including tumor and non-tumor tissue and lymph node tissues; bone marrow aspirates and fixed cell specimens. The preferred source of genomic DNA used in the present invention is from peripheral venous blood of each donor. Techniques to prepare genomic DNA
10 from biological samples are well known to the skilled technician. Details of a preferred embodiment are provided in Example 1. The person skilled in the art can choose to amplify pooled or unpooled DNA samples.

DNA Amplification

The identification of biallelic markers in a sample of genomic DNA may be facilitated
15 through the use of DNA amplification methods. DNA samples can be pooled or unpooled for the amplification step. DNA amplification techniques are well known to those skilled in the art. Various methods to amplify DNA fragments carrying biallelic markers are further described hereinbefore in "Amplification of the *HKLP* gene". The PCR technology is the preferred amplification technique used to identify new biallelic markers. A typical example of a PCR reaction
20 suitable for the purposes of the present invention is provided in Example 2.

In a first embodiment of the present invention, biallelic markers are identified using genomic sequence information generated by the inventors. Sequenced genomic DNA fragments are used to design primers for the amplification of 500 bp fragments. These 500 bp fragments are amplified from genomic DNA and are scanned for biallelic markers. Primers may be designed using the OSP
25 software (Hillier L. and Green P., 1991). All primers may contain, upstream of the specific target bases, a common oligonucleotide tail that serves as a sequencing primer. Those skilled in the art are familiar with primer extensions, which can be used for these purposes.

Preferred primers, useful for the amplification of genomic sequences encoding the candidate genes, focus on promoters, exons and splice sites of the genes. A biallelic marker presents a higher
30 probability to be an eventual causal mutation if it is located in these functional regions of the gene. Preferred amplification primers of the invention include the nucleotide sequences B1 to B25 and C1 to C25, detailed further in Example 2, Table 1.

Sequencing Of Amplified Genomic DNA And Identification Of Single Nucleotide Polymorphisms

35 The amplification products generated as described above, are then sequenced using any method known and available to the skilled technician. Methods for sequencing DNA using either

the dideoxy-mediated method (Sanger method) or the Maxam-Gilbert method are widely known to those of ordinary skill in the art. Such methods are for example disclosed in Sambrook et al.(1989). Alternative approaches include hybridization to high-density DNA probe arrays as described in Chee et al.(1996).

5 Preferably, the amplified DNA is subjected to automated dideoxy terminator sequencing reactions using a dye-primer cycle sequencing protocol. The products of the sequencing reactions are run on sequencing gels and the sequences are determined using gel image analysis. The polymorphism search is based on the presence of superimposed peaks in the electrophoresis pattern resulting from different bases occurring at the same position. Because each dideoxy terminator is
10 labeled with a different fluorescent molecule, the two peaks corresponding to a biallelic site present distinct colors corresponding to two different nucleotides at the same position on the sequence. However, the presence of two peaks can be an artifact due to background noise. To exclude such an artifact, the two DNA strands are sequenced and a comparison between the peaks is carried out. In order to be registered as a polymorphic sequence, the polymorphism has to be detected on both
15 strands.

The above procedure permits those amplification products, which contain biallelic markers to be identified. The detection limit for the frequency of biallelic polymorphisms detected by sequencing pools of 100 individuals is approximately 0.1 for the minor allele, as verified by sequencing pools of known allelic frequencies. However, more than 90% of the biallelic
20 polymorphisms detected by the pooling method have a frequency for the minor allele higher than 0.25. Therefore, the biallelic markers selected by this method have a frequency of at least 0.1 for the minor allele and less than 0.9 for the major allele. Preferably at least 0.2 for the minor allele and less than 0.8 for the major allele, more preferably at least 0.3 for the minor allele and less than 0.7 for the major allele, thus a heterozygosity rate higher than 0.18, preferably higher than 0.32, more
25 preferably higher than 0.42.

In another embodiment, biallelic markers are detected by sequencing individual DNA samples, the frequency of the minor allele of such a biallelic marker may be less than 0.1.

Validation Of The Biallelic Markers Of The Present Invention

The polymorphisms are evaluated for their usefulness as genetic markers by validating that
30 both alleles are present in a population. Validation of the biallelic markers is accomplished by genotyping a group of individuals by a method of the invention and demonstrating that both alleles are present. Microsequencing is a preferred method of genotyping alleles. The validation by genotyping step may be performed on individual samples derived from each individual in the group or by genotyping a pooled sample derived from more than one individual. The group can be as
35 small as one individual if that individual is heterozygous for the allele in question. Preferably the group contains at least three individuals, more preferably the group contains five or six individuals,

so that a single validation test will be more likely to result in the validation of more of the biallelic markers that are being tested. It should be noted, however, that when the validation test is performed on a small group it may result in a false negative result if as a result of sampling error none of the individuals tested carries one of the two alleles. Thus, the validation process is less useful in demonstrating that a particular initial result is an artifact, than it is at demonstrating that there is a *bona fide* biallelic marker at a particular position in a sequence. All of the genotyping, haplotyping, association, and interaction study methods of the invention may optionally be performed solely with validated biallelic markers.

Evaluation Of The Frequency Of The Biallelic Markers Of The Present Invention

The validated biallelic markers are further evaluated for their usefulness as genetic markers by determining the frequency of the least common allele at the biallelic marker site. The higher the frequency of the less common allele the greater the usefulness of the biallelic marker is association and interaction studies. The determination of the least common allele is accomplished by genotyping a group of individuals by a method of the invention and demonstrating that both alleles are present. This determination of frequency by genotyping step may be performed on individual samples derived from each individual in the group or by genotyping a pooled sample derived from more than one individual. The group must be large enough to be representative of the population as a whole. Preferably the group contains at least 20 individuals, more preferably the group contains at least 50 individuals, most preferably the group contains at least 100 individuals. Of course the larger the group the greater the accuracy of the frequency determination because of reduced sampling error. A biallelic marker wherein the frequency of the less common allele is 30% or more is termed a "high quality biallelic marker." All of the genotyping, haplotyping, association, and interaction study methods of the invention may optionally be performed solely with high quality biallelic markers.

The invention also relates to methods of estimating the frequency of an allele in a population comprising: a) genotyping individuals from said population for said biallelic marker according to the method of the present invention; b) determining the proportional representation of said biallelic marker in said population. In addition, the methods of estimating the frequency of an allele in a population of the invention encompass methods with any further limitation described in this disclosure, or those following, specified alone or in any combination; optionally, wherein said *HKLP*-related biallelic marker is selected from the group consisting of A1 to A32, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, wherein said *HKLP*-related biallelic marker is selected from the group consisting of A1 to A22 and A25 to A32, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, wherein said *HKLP*-related biallelic marker is selected from the group consisting of A23 and A24; optionally, determining the frequency of a biallelic marker allele in a population may be accomplished by determining the identity of the nucleotides for both

copies of said biallelic marker present in the genome of each individual in said population and calculating the proportional representation of said nucleotide at said *HKLP*-related biallelic marker for the population; optionally, determining the proportional representation may be accomplished by performing a genotyping method of the invention on a pooled biological sample derived from a
5 representative number of individuals, or each individual, in said population, and calculating the proportional amount of said nucleotide compared with the total.

Methods For Genotyping An Individual For Biallelic Markers

Methods are provided to genotype a biological sample for one or more biallelic markers of the present invention, all of which may be performed *in vitro*. Such methods of genotyping
10 comprise determining the identity of a nucleotide at a *HKLP* biallelic marker site by any method known in the art. These methods find use in genotyping case-control populations in association studies as well as individuals in the context of detection of alleles of biallelic markers which are known to be associated with a given trait, in which case both copies of the biallelic marker present in individual's genome are determined so that an individual may be classified as homozygous or
15 heterozygous for a particular allele.

These genotyping methods can be performed on nucleic acid samples derived from a single individual or pooled DNA samples.

Genotyping can be performed using similar methods as those described above for the identification of the biallelic markers, or using other genotyping methods such as those further
20 described below. In preferred embodiments, the comparison of sequences of amplified genomic fragments from different individuals is used to identify new biallelic markers whereas microsequencing is used for genotyping known biallelic markers in diagnostic and association study applications.

In one embodiment the invention encompasses methods of genotyping comprising
25 determining the identity of a nucleotide at a *HKLP*-related biallelic marker or the complement thereof in a biological sample; Optionally, said *HKLP*-related biallelic marker is selected from the group consisting of A1 to A32, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, said *HKLP*-related biallelic marker is selected from the group consisting of A1 to A17, and A20 to A22, and the complements thereof, or optionally the
30 biallelic markers in linkage disequilibrium therewith; optionally, said *HKLP*-related biallelic marker is selected from the group consisting of A23 and A24, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, wherein said biological sample is derived from a single subject; optionally, wherein the identity of the nucleotides at said biallelic marker is determined for both copies of said biallelic marker present in said individual's genome;
35 optionally, wherein said biological sample is derived from multiple subjects; optionally, further comprising amplifying a portion of said sequence comprising the biallelic marker prior to said

determining step; optionally, wherein said amplifying is performed by PCR; optionally, wherein said determining is performed by a hybridization assay, a sequencing assay, a microsequencing assay, or an enzyme-based mismatch detection assay.

Source of DNA for genotyping

5 Any source of nucleic acids, in purified or non-purified form, can be utilized as the starting nucleic acid, provided it contains or is suspected of containing the specific nucleic acid sequence desired. DNA or RNA may be extracted from cells, tissues, body fluids and the like as described above. While nucleic acids for use in the genotyping methods of the invention can be derived from any mammalian source, the test subjects and individuals from which nucleic acid samples are taken
10 are generally understood to be human.

Amplification Of DNA Fragments Comprising Biallelic Markers

Methods and polynucleotides are provided to amplify a segment of nucleotides comprising one or more biallelic marker of the present invention. It will be appreciated that amplification of DNA fragments comprising biallelic markers may be used in various methods and for various
15 purposes and is not restricted to genotyping. Nevertheless, many genotyping methods, although not all, require the previous amplification of the DNA region carrying the biallelic marker of interest. Such methods specifically increase the concentration or total number of sequences that span the biallelic marker or include that site and sequences located either distal or proximal to it. Diagnostic assays may also rely on amplification of DNA segments carrying a biallelic marker of the present
20 invention. Amplification of DNA may be achieved by any method known in the art. Amplification techniques are described above in the section entitled, "Amplification of the *HKLP* gene".

Some of these amplification methods are particularly suited for the detection of single nucleotide polymorphisms and allow the simultaneous amplification of a target sequence and the identification of the polymorphic nucleotide as it is further described below.

25 The identification of biallelic markers as described above allows the design of appropriate oligonucleotides, which can be used as primers to amplify DNA fragments comprising the biallelic markers of the present invention. Amplification can be performed using the primers initially used to discover new biallelic markers which are described herein or any set of primers allowing the amplification of a DNA fragment comprising a biallelic marker of the present invention.

30 In some embodiments the present invention provides primers for amplifying a DNA fragment containing one or more biallelic markers of the present invention. Preferred amplification primers are listed in Example 2. It will be appreciated that the primers listed are merely exemplary and that any other set of primers which produce amplification products containing one or more biallelic markers of the present invention.

35 The spacing of the primers determines the length of the segment to be amplified. In the context of the present invention, amplified segments carrying biallelic markers can range in size

from at least about 25 bp to 35 kbp. Amplification fragments from 25-3000 bp are typical, fragments from 50-1000 bp are preferred and fragments from 100-600 bp are highly preferred. It will be appreciated that amplification primers for the biallelic markers may be any sequence which allow the specific amplification of any DNA fragment carrying the markers. Amplification primers
5 may be labeled or immobilized on a solid support as described in "Oligonucleotide probes and primers".

Methods of Genotyping DNA samples for Biallelic Markers

Any method known in the art can be used to identify the nucleotide present at a biallelic marker site. Since the biallelic marker allele to be detected has been identified and specified in the
10 present invention, detection will prove simple for one of ordinary skill in the art by employing any of a number of techniques. Many genotyping methods require the previous amplification of the DNA region carrying the biallelic marker of interest. While the amplification of target or signal is often preferred at present, ultrasensitive detection methods which do not require amplification are also encompassed by the present genotyping methods. Methods well-known to those skilled in the
15 art that can be used to detect biallelic polymorphisms include methods such as, conventional dot blot analyzes, single strand conformational polymorphism analysis (SSCP) described by Orita et al.(1989), denaturing gradient gel electrophoresis (DGGE), heteroduplex analysis, mismatch cleavage detection, and other conventional techniques as described in Sheffield et al.(1991), White et al.(1992), Grompe et al.(1989 and 1993). Another method for determining the identity of the
20 nucleotide present at a particular polymorphic site employs a specialized exonuclease-resistant nucleotide derivative as described in US patent 4,656,127.

Preferred methods involve directly determining the identity of the nucleotide present at a biallelic marker site by sequencing assay, enzyme-based mismatch detection assay, or hybridization assay. The following is a description of some preferred methods. A highly preferred method is the
25 microsequencing technique. The term "sequencing" is used herein to refer to polymerase extension of duplex primer/template complexes and includes both traditional sequencing and microsequencing.

1) Sequencing Assays

The nucleotide present at a polymorphic site can be determined by sequencing methods. In a preferred embodiment, DNA samples are subjected to PCR amplification before sequencing as
30 described above. DNA sequencing methods are described in "Sequencing Of Amplified Genomic DNA And Identification Of Single Nucleotide Polymorphisms".

Preferably, the amplified DNA is subjected to automated dideoxy terminator sequencing reactions using a dye-primer cycle sequencing protocol. Sequence analysis allows the identification of the base present at the biallelic marker site.

2) Microsequencing Assays

In microsequencing methods, the nucleotide at a polymorphic site in a target DNA is detected by a single nucleotide primer extension reaction. This method involves appropriate microsequencing primers which, hybridize just upstream of the polymorphic base of interest in the target nucleic acid. A polymerase is used to specifically extend the 3' end of the primer with one single ddNTP (chain terminator) complementary to the nucleotide at the polymorphic site. Next the identity of the incorporated nucleotide is determined in any suitable way.

Typically, microsequencing reactions are carried out using fluorescent ddNTPs and the extended microsequencing primers are analyzed by electrophoresis on ABI 377 sequencing machines to determine the identity of the incorporated nucleotide as described in EP 412 883. Alternatively capillary electrophoresis can be used in order to process a higher number of assays simultaneously. An example of a typical microsequencing procedure that can be used in the context of the present invention is provided in Example 4.

Different approaches can be used for the labeling and detection of ddNTPs. A homogeneous phase detection method based on fluorescence resonance energy transfer has been described by Chen and Kwok (1997) and Chen et al.(1997). In this method, amplified genomic DNA fragments containing polymorphic sites are incubated with a 5'-fluorescein-labeled primer in the presence of allelic dye-labeled dideoxynucleoside triphosphates and a modified Taq polymerase. The dye-labeled primer is extended one base by the dye-terminator specific for the allele present on the template. At the end of the genotyping reaction, the fluorescence intensities of the two dyes in the reaction mixture are analyzed directly without separation or purification. All these steps can be performed in the same tube and the fluorescence changes can be monitored in real time. Alternatively, the extended primer may be analyzed by MALDI-TOF Mass Spectrometry. The base at the polymorphic site is identified by the mass added onto the microsequencing primer (see Haff and Smirnov, 1997).

Microsequencing may be achieved by the established microsequencing method or by developments or derivatives thereof. Alternative methods include several solid-phase microsequencing techniques. The basic microsequencing protocol is the same as described previously, except that the method is conducted as a heterogeneous phase assay, in which the primer or the target molecule is immobilized or captured onto a solid support. To simplify the primer separation and the terminal nucleotide addition analysis, oligonucleotides are attached to solid supports or are modified in such ways that permit affinity separation as well as polymerase extension. The 5' ends and internal nucleotides of synthetic oligonucleotides can be modified in a number of different ways to permit different affinity separation approaches, e.g., biotinylation. If a single affinity group is used on the oligonucleotides, the oligonucleotides can be separated from the incorporated terminator reagent. This eliminates the need of physical or size separation. More than one oligonucleotide can be separated from the terminator reagent and analyzed simultaneously if

more than one affinity group is used. This permits the analysis of several nucleic acid species or more nucleic acid sequence information per extension reaction. The affinity group need not be on the priming oligonucleotide but could alternatively be present on the template. For example, immobilization can be carried out via an interaction between biotinylated DNA and streptavidin-coated microtitration wells or avidin-coated polystyrene particles. In the same manner, oligonucleotides or templates may be attached to a solid support in a high-density format. In such solid phase microsequencing reactions, incorporated ddNTPs can be radiolabeled (Syvänen, 1994) or linked to fluorescein (Livak and Hainer, 1994). The detection of radiolabeled ddNTPs can be achieved through scintillation-based techniques. The detection of fluorescein-linked ddNTPs can be based on the binding of anti-fluorescein antibody conjugated with alkaline phosphatase, followed by incubation with a chromogenic substrate (such as *p*-nitrophenyl phosphate). Other possible reporter-detection pairs include: ddNTP linked to dinitrophenyl (DNP) and anti-DNP alkaline phosphatase conjugate (Harju et al., 1993) or biotinylated ddNTP and horseradish peroxidase-conjugated streptavidin with *o*-phenylenediamine as a substrate (WO 92/15712). As yet another alternative solid-phase microsequencing procedure, Nyren et al. (1993) described a method relying on the detection of DNA polymerase activity by an enzymatic luminometric inorganic pyrophosphate detection assay (ELIDA).

Pastinen et al. (1997) describe a method for multiplex detection of single nucleotide polymorphism in which the solid phase minisequencing principle is applied to an oligonucleotide array format. High-density arrays of DNA probes attached to a solid support (DNA chips) are further described below.

In one aspect the present invention provides polynucleotides and methods to genotype one or more biallelic markers of the present invention by performing a microsequencing assay. Preferred microsequencing primers include the nucleotide sequences D1 to D30 and E1 to E30. It will be appreciated that the microsequencing primers listed in Example 4 are merely exemplary and that, any primer having a 3' end immediately adjacent to the polymorphic nucleotide may be used. Similarly, it will be appreciated that microsequencing analysis may be performed for any biallelic marker or any combination of biallelic markers of the present invention. One aspect of the present invention is a solid support which includes one or more microsequencing primers listed in Example 4, or fragments comprising at least 8, 12, 15, 20, 25, 30, 40, or 50 consecutive nucleotides thereof and having a 3' terminus immediately upstream of the corresponding biallelic marker, for determining the identity of a nucleotide at a biallelic marker site.

3) Mismatch detection assays based on polymerases and ligases

In one aspect the present invention provides polynucleotides and methods to determine the allele of one or more biallelic markers of the present invention in a biological sample, by mismatch detection assays based on polymerases and/or ligases. These assays are based on the specificity of polymerases and ligases. Polymerization reactions places particularly stringent requirements on

correct base pairing of the 3' end of the amplification primer and the joining of two oligonucleotides hybridized to a target DNA sequence is quite sensitive to mismatches close to the ligation site, especially at the 3' end. Methods, primers and various parameters to amplify DNA fragments comprising biallelic markers of the present invention are further described above in "Amplification
5 Of DNA Fragments Comprising Biallelic Markers".

Allele Specific Amplification Primers

Discrimination between the two alleles of a biallelic marker can also be achieved by allele specific amplification, a selective strategy, whereby one of the alleles is amplified without amplification of the other allele. This is accomplished by placing the polymorphic base at the 3' end
10 of one of the amplification primers. Because the extension forms from the 3' end of the primer, a mismatch at or near this position has an inhibitory effect on amplification. Therefore, under appropriate amplification conditions, these primers only direct amplification on their complementary allele. Determining the precise location of the mismatch and the corresponding assay conditions are well with the ordinary skill in the art.

15 Ligation/Amplification Based Methods.

The "Oligonucleotide Ligation Assay" (OLA) uses two oligonucleotides which are designed to be capable of hybridizing to abutting sequences of a single strand of a target molecules. One of the oligonucleotides is biotinylated, and the other is detectably labeled. If the precise complementary sequence is found in a target molecule, the oligonucleotides will hybridize such that
20 their termini abut, and create a ligation substrate that can be captured and detected. OLA is capable of detecting single nucleotide polymorphisms and may be advantageously combined with PCR as described by Nickerson et al.(1990). In this method, PCR is used to achieve the exponential amplification of target DNA, which is then detected using OLA.

Other amplification methods which are particularly suited for the detection of single
25 nucleotide polymorphism include LCR (ligase chain reaction), Gap LCR (GLCR) which are described above in "Amplification of the *HKLP* gene". LCR uses two pairs of probes to exponentially amplify a specific target. The sequences of each pair of oligonucleotides, is selected to permit the pair to hybridize to abutting sequences of the same strand of the target. Such hybridization forms a substrate for a template-dependant ligase. In accordance with the present
30 invention, LCR can be performed with oligonucleotides having the proximal and distal sequences of the same strand of a biallelic marker site. In one embodiment, either oligonucleotide will be designed to include the biallelic marker site. In such an embodiment, the reaction conditions are selected such that the oligonucleotides can be ligated together only if the target molecule either
35 oligonucleotide. In an alternative embodiment, the oligonucleotides will not include the biallelic marker, such that when they hybridize to the target molecule, a "gap" is created as described in WO

90/01069. This gap is then "filled" with complementary dNTPs (as mediated by DNA polymerase), or by an additional pair of oligonucleotides. Thus at the end of each cycle, each single strand has a complement capable of serving as a target during the next cycle and exponential allele-specific amplification of the desired sequence is obtained.

- 5 Ligase/Polymerase-mediated Genetic Bit AnalysisTM is another method for determining the identity of a nucleotide at a preselected site in a nucleic acid molecule (WO 95/21271). This method involves the incorporation of a nucleoside triphosphate that is complementary to the nucleotide present at the preselected site onto the terminus of a primer molecule, and their subsequent ligation to a second oligonucleotide. The reaction is monitored by detecting a specific label attached to the
- 10 reaction's solid phase or by detection in solution.

4) Hybridization Assay Methods

- A preferred method of determining the identity of the nucleotide present at a biallelic marker site involves nucleic acid hybridization. The hybridization probes, which can be conveniently used in such reactions, preferably include the probes defined herein. Any hybridization assay may be
- 15 used including Southern hybridization, Northern hybridization, dot blot hybridization and solid-phase hybridization (see Sambrook et al., 1989).

- Hybridization refers to the formation of a duplex structure by two single stranded nucleic acids due to complementary base pairing. Hybridization can occur between exactly complementary nucleic acid strands or between nucleic acid strands that contain minor regions of mismatch.
- 20 Specific probes can be designed that hybridize to one form of a biallelic marker and not to the other and therefore are able to discriminate between different allelic forms. Allele-specific probes are often used in pairs, one member of a pair showing perfect match to a target sequence containing the original allele and the other showing a perfect match to the target sequence containing the alternative allele. Hybridization conditions should be sufficiently stringent that there is a significant difference
- 25 in hybridization intensity between alleles, and preferably an essentially binary response, whereby a probe hybridizes to only one of the alleles. Stringent, sequence specific hybridization conditions, under which a probe will hybridize only to the exactly complementary target sequence are well known in the art (Sambrook et al., 1989). Stringent conditions are sequence dependent and will be different in different circumstances. Generally, stringent conditions are selected to be about 5°C
- 30 lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. Although such hybridizations can be performed in solution, it is preferred to employ a solid-phase hybridization assay. The target DNA comprising a biallelic marker of the present invention may be amplified prior to the hybridization reaction. The presence of a specific allele in the sample is determined by detecting the presence or the absence of stable hybrid duplexes formed between the
- 35 probe and the target DNA. The detection of hybrid duplexes can be carried out by a number of methods. Various detection assay formats are well known which utilize detectable labels bound to either the target or the probe to enable detection of the hybrid duplexes. Typically, hybridization

duplexes are separated from unhybridized nucleic acids and the labels bound to the duplexes are then detected. Those skilled in the art will recognize that wash steps may be employed to wash away excess target DNA or probe as well as unbound conjugate. Further, standard heterogeneous assay formats are suitable for detecting the hybrids using the labels present on the primers and probes.

5 Two recently developed assays allow hybridization-based allele discrimination with no need for separations or washes (see Landegren U. et al., 1998). The TaqMan assay takes advantage of the 5' nuclease activity of Taq DNA polymerase to digest a DNA probe annealed specifically to the accumulating amplification product. TaqMan probes are labeled with a donor-acceptor dye pair that interacts via fluorescence energy transfer. Cleavage of the TaqMan probe by the advancing
10 polymerase during amplification dissociates the donor dye from the quenching acceptor dye, greatly increasing the donor fluorescence. All reagents necessary to detect two allelic variants can be assembled at the beginning of the reaction and the results are monitored in real time (see Livak et al., 1995). In an alternative homogeneous hybridization based procedure, molecular beacons are used for allele discriminations. Molecular beacons are hairpin-shaped oligonucleotide probes that report
15 the presence of specific nucleic acids in homogeneous solutions. When they bind to their targets they undergo a conformational reorganization that restores the fluorescence of an internally quenched fluorophore (Tyagi et al., 1998).

The polynucleotides provided herein can be used to produce probes which can be used in hybridization assays for the detection of biallelic marker alleles in biological samples. These probes
20 are characterized in that they preferably comprise between 8 and 50 nucleotides, and in that they are sufficiently complementary to a sequence comprising a biallelic marker of the present invention to hybridize thereto and preferably sufficiently specific to be able to discriminate the targeted sequence for only one nucleotide variation. A particularly preferred probe is 25 nucleotides in length. Preferably the biallelic marker is within 4 nucleotides of the center of the polynucleotide probe. In
25 particularly preferred probes, the biallelic marker is at the center of said polynucleotide. Preferred probes comprise a nucleotide sequence selected from the group consisting of amplicons listed in Table 1 and the sequences complementary thereto, or a fragment thereof, said fragment comprising at least about 8 consecutive nucleotides, preferably 10, 15, 20, more preferably 25, 30, 40, 47, or 50 consecutive nucleotides and containing a polymorphic base. In preferred embodiments the
30 polymorphic base is within 5, 4, 3, 2, 1, nucleotides of the center of the said polynucleotide, more preferably at the center of said polynucleotide.

Preferably the probes of the present invention are labeled or immobilized on a solid support. Labels and solid supports are further described in "Oligonucleotide Probes and Primers". The probes can be non-extendable as described in "Oligonucleotide Probes and Primers".

35 By assaying the hybridization to an allele specific probe, one can detect the presence or absence of a biallelic marker allele in a given sample. High-Throughput parallel hybridizations in array format are specifically encompassed within "hybridization assays" and are described below.

5) Hybridization To Addressable Arrays Of Oligonucleotides

Hybridization assays based on oligonucleotide arrays rely on the differences in hybridization stability of short oligonucleotides to perfectly matched and mismatched target sequence variants. Efficient access to polymorphism information is obtained through a basic structure comprising high-density arrays of oligonucleotide probes attached to a solid support (e.g., the chip) at selected positions. Each DNA chip can contain thousands to millions of individual synthetic DNA probes arranged in a grid-like pattern and miniaturized to the size of a dime.

The chip technology has already been applied with success in numerous cases. For example, the screening of mutations has been undertaken in the BRCA1 gene, in *S. cerevisiae* mutant strains, and in the protease gene of HIV-1 virus (Hacia et al., 1996; Shoemaker et al., 1996; Kozal et al., 1996). Chips of various formats for use in detecting biallelic polymorphisms can be produced on a customized basis by Affymetrix (GeneChip™), Hyseq (HyChip and HyGnostics), and Protogene Laboratories.

In general, these methods employ arrays of oligonucleotide probes that are complementary to target nucleic acid sequence segments from an individual which, target sequences include a polymorphic marker. EP 785280 describes a tiling strategy for the detection of single nucleotide polymorphisms. Briefly, arrays may generally be "tiling" for a large number of specific polymorphisms. By "tiling" is generally meant the synthesis of a defined set of oligonucleotide probes which is made up of a sequence complementary to the target sequence of interest, as well as preselected variations of that sequence, e.g., substitution of one or more given positions with one or more members of the basis set of monomers, i.e. nucleotides. Tiling strategies are further described in PCT application No. WO 95/11995. In a particular aspect, arrays are tiled for a number of specific, identified biallelic marker sequences. In particular, the array is tiled to include a number of detection blocks, each detection block being specific for a specific biallelic marker or a set of biallelic markers. For example, a detection block may be tiled to include a number of probes, which span the sequence segment that includes a specific polymorphism. To ensure probes that are complementary to each allele, the probes are synthesized in pairs differing at the biallelic marker. In addition to the probes differing at the polymorphic base, monosubstituted probes are also generally tiled within the detection block. These monosubstituted probes have bases at and up to a certain number of bases in either direction from the polymorphism, substituted with the remaining nucleotides (selected from A, T, G, C and U). Typically the probes in a tiled detection block will include substitutions of the sequence positions up to and including those that are 5 bases away from the biallelic marker. The monosubstituted probes provide internal controls for the tiled array, to distinguish actual hybridization from artefactual cross-hybridization. Upon completion of hybridization with the target sequence and washing of the array, the array is scanned to determine the position on the array to which the target sequence hybridizes. The hybridization data from the scanned array is then analyzed to identify which allele or alleles of the biallelic marker are present in

the sample. Hybridization and scanning may be carried out as described in PCT application No. WO 92/10092 and WO 95/11995 and US patent No. 5,424,186.

Thus, in some embodiments, the chips may comprise an array of nucleic acid sequences of fragments of about 15 nucleotides in length. In further embodiments, the chip may comprise an array including at least one of the sequences selected from the group consisting of amplicons listed in table 1 and the sequences complementary thereto, or a fragment thereof, said fragment comprising at least about 8 consecutive nucleotides, preferably 10, 15, 20, more preferably 25, 30, 40, 47, or 50 consecutive nucleotides and containing a polymorphic base. In preferred embodiments the polymorphic base is within 5, 4, 3, 2, 1, nucleotides of the center of the said polynucleotide, more preferably at the center of said polynucleotide. In some embodiments, the chip may comprise an array of at least 2, 3, 4, 5, 6, 7, 8 or more of these polynucleotides of the invention. Solid supports and polynucleotides of the present invention attached to solid supports are further described in "oligonucleotide probes and primers".

6) Integrated Systems

Another technique, which may be used to analyze polymorphisms, includes multicomponent integrated systems, which miniaturize and compartmentalize processes such as PCR and capillary electrophoresis reactions in a single functional device. An example of such technique is disclosed in US patent 5,589,136, which describes the integration of PCR amplification and capillary electrophoresis in chips.

Integrated systems can be envisaged mainly when microfluidic systems are used. These systems comprise a pattern of microchannels designed onto a glass, silicon, quartz, or plastic wafer included on a microchip. The movements of the samples are controlled by electric, electroosmotic or hydrostatic forces applied across different areas of the microchip to create functional microscopic valves and pumps with no moving parts.

For genotyping biallelic markers, the microfluidic system may integrate nucleic acid amplification, microsequencing, capillary electrophoresis and a detection method such as laser-induced fluorescence detection.

Methods Of Genetic Analysis Using The Biallelic Markers Of The Present Invention

Different methods are available for the genetic analysis of complex traits (see Lander and Schork, 1994). The search for disease-susceptibility genes is conducted using two main methods: the linkage approach in which evidence is sought for cosegregation between a locus and a putative trait locus using family studies, and the association approach in which evidence is sought for a statistically significant association between an allele and a trait or a trait causing allele (Khoury et al., 1993). In general, the biallelic markers of the present invention find use in any method known in the art to demonstrate a statistically significant correlation between a genotype and a phenotype. The biallelic markers may be used in parametric and non-parametric linkage analysis methods.

Preferably, the biallelic markers of the present invention are used to identify genes associated with detectable traits using association studies, an approach which does not require the use of affected families and which permits the identification of genes associated with complex and sporadic traits.

The genetic analysis using the biallelic markers of the present invention may be conducted
5 on any scale. The whole set of biallelic markers of the present invention or any subset of biallelic markers of the present invention corresponding to the candidate gene may be used. Further, any set of genetic markers including a biallelic marker of the present invention may be used. A set of biallelic polymorphisms that could be used as genetic markers in combination with the biallelic markers of the present invention has been described in WO 98/20165. As mentioned above, it
10 should be noted that the biallelic markers of the present invention may be included in any complete or partial genetic map of the human genome. These different uses are specifically contemplated in the present invention and claims.

Linkage Analysis

Linkage analysis is based upon establishing a correlation between the transmission of
15 genetic markers and that of a specific trait throughout generations within a family. Thus, the aim of linkage analysis is to detect marker loci that show cosegregation with a trait of interest in pedigrees.

Parametric Methods

When data are available from successive generations there is the opportunity to study the degree of linkage between pairs of loci. Estimates of the recombination fraction enable loci to be
20 ordered and placed onto a genetic map. With loci that are genetic markers, a genetic map can be established, and then the strength of linkage between markers and traits can be calculated and used to indicate the relative positions of markers and genes affecting those traits (Weir, 1996). The classical method for linkage analysis is the logarithm of odds (lod) score method (see Morton, 1955; Ott, 1991). Calculation of lod scores requires specification of the mode of inheritance for the
25 disease (parametric method). Generally, the length of the candidate region identified using linkage analysis is between 2 and 20Mb. Once a candidate region is identified as described above, analysis of recombinant individuals using additional markers allows further delineation of the candidate region. Linkage analysis studies have generally relied on the use of a maximum of 5,000 microsatellite markers, thus limiting the maximum theoretical attainable resolution of linkage
30 analysis to about 600 kb on average.

Linkage analysis has been successfully applied to map simple genetic traits that show clear Mendelian inheritance patterns and which have a high penetrance (i.e., the ratio between the number of trait positive carriers of allele a and the total number of a carriers in the population). However, parametric linkage analysis suffers from a variety of drawbacks. First, it is limited by its reliance on
35 the choice of a genetic model suitable for each studied trait. Furthermore, as already mentioned, the resolution attainable using linkage analysis is limited, and complementary studies are required to

refine the analysis of the typical 2Mb to 20Mb regions initially identified through linkage analysis. In addition, parametric linkage analysis approaches have proven difficult when applied to complex genetic traits, such as those due to the combined action of multiple genes and/or environmental factors. It is very difficult to model these factors adequately in a lod score analysis. In such cases, too large an effort and cost are needed to recruit the adequate number of affected families required for applying linkage analysis to these situations, as recently discussed by Risch, N. and Merikangas, K. (1996).

Non-Parametric Methods

The advantage of the so-called non-parametric methods for linkage analysis is that they do not require specification of the mode of inheritance for the disease, they tend to be more useful for the analysis of complex traits. In non-parametric methods, one tries to prove that the inheritance pattern of a chromosomal region is not consistent with random Mendelian segregation by showing that affected relatives inherit identical copies of the region more often than expected by chance. Affected relatives should show excess "allele sharing" even in the presence of incomplete penetrance and polygenic inheritance. In non-parametric linkage analysis the degree of agreement at a marker locus in two individuals can be measured either by the number of alleles identical by state (IBS) or by the number of alleles identical by descent (IBD). Affected sib pair analysis is a well-known special case and is the simplest form of these methods.

The biallelic markers of the present invention may be used in both parametric and non-parametric linkage analysis. Preferably biallelic markers may be used in non-parametric methods which allow the mapping of genes involved in complex traits. The biallelic markers of the present invention may be used in both IBD- and IBS- methods to map genes affecting a complex trait. In such studies, taking advantage of the high density of biallelic markers, several adjacent biallelic marker loci may be pooled to achieve the efficiency attained by multi-allelic markers (Zhao et al., 1998).

Population Association Studies

The present invention comprises methods for identifying if the *HKLP* gene is associated with a detectable trait using the biallelic markers of the present invention. In one embodiment the present invention comprises methods to detect an association between a biallelic marker allele or a biallelic marker haplotype and a trait. Further, the invention comprises methods to identify a trait causing allele in linkage disequilibrium with any biallelic marker allele of the present invention.

As described above, alternative approaches can be employed to perform association studies: genome-wide association studies, candidate region association studies and candidate gene association studies. In a preferred embodiment, the biallelic markers of the present invention are used to perform candidate gene association studies. The candidate gene analysis clearly provides a short-cut approach to the identification of genes and gene polymorphisms related to a particular trait

when some information concerning the biology of the trait is available. Further, the biallelic markers of the present invention may be incorporated in any map of genetic markers of the human genome in order to perform genome-wide association studies. Methods to generate a high-density map of biallelic markers has been described in US Provisional Patent application serial number 5 60/082,614. The biallelic markers of the present invention may further be incorporated in any map of a specific candidate region of the genome (a specific chromosome or a specific chromosomal segment for example).

As mentioned above, association studies may be conducted within the general population and are not limited to studies performed on related individuals in affected families. Association 10 studies are extremely valuable as they permit the analysis of sporadic or multifactor traits. Moreover, association studies represent a powerful method for fine-scale mapping enabling much finer mapping of trait causing alleles than linkage studies. Studies based on pedigrees often only narrow the location of the trait causing allele. Association studies using the biallelic markers of the present invention can therefore be used to refine the location of a trait causing allele in a candidate 15 region identified by Linkage Analysis methods. Moreover, once a chromosome segment of interest has been identified, the presence of a candidate gene such as a candidate gene of the present invention, in the region of interest can provide a shortcut to the identification of the trait causing allele. Biallelic markers of the present invention can be used to demonstrate that a candidate gene is associated with a trait. Such uses are specifically contemplated in the present invention.

20 **Determining The Frequency Of A Biallelic Marker Allele Or Of A Biallelic Marker Haplotype In A Population**

Association studies explore the relationships among frequencies for sets of alleles between loci.

Determining The Frequency Of An Allele In A Population

25 Allelic frequencies of the biallelic markers in a populations can be determined using one of the methods described above under the heading "Methods for genotyping an individual for biallelic markers", or any genotyping procedure suitable for this intended purpose. Genotyping pooled samples or individual samples can determine the frequency of a biallelic marker allele in a population. One way to reduce the number of genotypings required is to use pooled samples. A 30 major obstacle in using pooled samples is in terms of accuracy and reproducibility for determining accurate DNA concentrations in setting up the pools. Genotyping individual samples provides higher sensitivity, reproducibility and accuracy and; is the preferred method used in the present invention. Preferably, each individual is genotyped separately and simple gene counting is applied to determine the frequency of an allele of a biallelic marker or of a genotype in a given population.

Determining The Frequency Of A Haplotype In A Population

The gametic phase of haplotypes is unknown when diploid individuals are heterozygous at more than one locus. Using genealogical information in families gametic phase can sometimes be inferred (Perlin et al., 1994). When no genealogical information is available different strategies may be used. One possibility is that the multiple-site heterozygous diploids can be eliminated from the analysis, keeping only the homozygotes and the single-site heterozygote individuals, but this approach might lead to a possible bias in the sample composition and the underestimation of low-frequency haplotypes. Another possibility is that single chromosomes can be studied independently, for example, by asymmetric PCR amplification (see Newton et al, 1989; Wu et al., 1989) or by isolation of single chromosome by limit dilution followed by PCR amplification (see Ruano et al., 1990). Further, a sample may be haplotyped for sufficiently close biallelic markers by double PCR amplification of specific alleles (Sarkar, G. and Sommer S. S., 1991). These approaches are not entirely satisfying either because of their technical complexity, the additional cost they entail, their lack of generalization at a large scale, or the possible biases they introduce. To overcome these difficulties, an algorithm to infer the phase of PCR-amplified DNA genotypes introduced by Clark, A.G.(1990) may be used. Briefly, the principle is to start filling a preliminary list of haplotypes present in the sample by examining unambiguous individuals, that is, the complete homozygotes and the single-site heterozygotes. Then other individuals in the same sample are screened for the possible occurrence of previously recognized haplotypes. For each positive identification, the complementary haplotype is added to the list of recognized haplotypes, until the phase information for all individuals is either resolved or identified as unresolved. This method assigns a single haplotype to each multiheterozygous individual, whereas several haplotypes are possible when there are more than one heterozygous site. Alternatively, one can use methods estimating haplotype frequencies in a population without assigning haplotypes to each individual. Preferably, a method based on an expectation-maximization (EM) algorithm (Dempster et al., 1977) leading to maximum-likelihood estimates of haplotype frequencies under the assumption of Hardy-Weinberg proportions (random mating) is used (see Excoffier L. and Slatkin M., 1995). The EM algorithm is a generalized iterative maximum-likelihood approach to estimation that is useful when data are ambiguous and/or incomplete. The EM algorithm is used to resolve heterozygotes into haplotypes. Haplotype estimations are further described below under the heading "Statistical Methods." Any other method known in the art to determine or to estimate the frequency of a haplotype in a population may be used.

The invention also encompasses methods of estimating the frequency of a haplotype for a set of biallelic markers in a population, comprising the steps of: a) genotyping at least one *HKLP*-related biallelic marker according to a method of the invention for each individual in said population; b) genotyping a second biallelic marker by determining the identity of the nucleotides at said second biallelic marker for both copies of said second biallelic marker present in the genome of

each individual in said population; and c) applying a haplotype determination method to the identities of the nucleotides determined in steps a) and b) to obtain an estimate of said frequency. In addition, the methods of estimating the frequency of a haplotype of the invention encompass methods with any further limitation described in this disclosure, or those following, specified alone
5 or in any combination: Optionally, said *HKLP*-related biallelic marker is selected from the group consisting of A1 to A32, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, said *HKLP*-related biallelic marker is selected from the group consisting of A1 to A17, and A20 to A22, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, said *HKLP*-related biallelic marker is
10 selected from the group consisting of A23 and A24, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; Optionally, said haplotype determination method is performed by asymmetric PCR amplification, double PCR amplification of specific alleles, the Clark algorithm, or an expectation-maximization algorithm.

Linkage Disequilibrium Analysis

15 Linkage disequilibrium is the non-random association of alleles at two or more loci and represents a powerful tool for mapping genes involved in disease traits (see Ajioka R.S. et al., 1997). Biallelic markers, because they are densely spaced in the human genome and can be genotyped in greater numbers than other types of genetic markers (such as RFLP or VNTR markers), are particularly useful in genetic analysis based on linkage disequilibrium.

20 When a disease mutation is first introduced into a population (by a new mutation or the immigration of a mutation carrier), it necessarily resides on a single chromosome and thus on a single "background" or "ancestral" haplotype of linked markers. Consequently, there is complete disequilibrium between these markers and the disease mutation: one finds the disease mutation only in the presence of a specific set of marker alleles. Through subsequent generations recombination
25 events occur between the disease mutation and these marker polymorphisms, and the disequilibrium gradually dissipates. The pace of this dissipation is a function of the recombination frequency, so the markers closest to the disease gene will manifest higher levels of disequilibrium than those that are further away. When not broken up by recombination, "ancestral" haplotypes and linkage disequilibrium between marker alleles at different loci can be tracked not only through pedigrees but
30 also through populations. Linkage disequilibrium is usually seen as an association between one specific allele at one locus and another specific allele at a second locus.

The pattern or curve of disequilibrium between disease and marker loci is expected to exhibit a maximum that occurs at the disease locus. Consequently, the amount of linkage disequilibrium between a disease allele and closely linked genetic markers may yield valuable
35 information regarding the location of the disease gene. For fine-scale mapping of a disease locus, it is useful to have some knowledge of the patterns of linkage disequilibrium that exist between

markers in the studied region. As mentioned above the mapping resolution achieved through the analysis of linkage disequilibrium is much higher than that of linkage studies. The high density of biallelic markers combined with linkage disequilibrium analysis provides powerful tools for fine-scale mapping. Different methods to calculate linkage disequilibrium are described below under the heading "Statistical Methods".

Population-Based Case-Control Studies Of Trait-Marker Associations

As mentioned above, the occurrence of pairs of specific alleles at different loci on the same chromosome is not random and the deviation from random is called linkage disequilibrium. Association studies focus on population frequencies and rely on the phenomenon of linkage disequilibrium. If a specific allele in a given gene is directly involved in causing a particular trait, its frequency will be statistically increased in an affected (trait positive) population, when compared to the frequency in a trait negative population or in a random control population. As a consequence of the existence of linkage disequilibrium, the frequency of all other alleles present in the haplotype carrying the trait-causing allele will also be increased in trait positive individuals compared to trait negative individuals or random controls. Therefore, association between the trait and any allele (specifically a biallelic marker allele) in linkage disequilibrium with the trait-causing allele will suffice to suggest the presence of a trait-related gene in that particular region. Case-control populations can be genotyped for biallelic markers to identify associations that narrowly locate a trait causing allele. As any marker in linkage disequilibrium with one given marker associated with a trait will be associated with the trait. Linkage disequilibrium allows the relative frequencies in case-control populations of a limited number of genetic polymorphisms (specifically biallelic markers) to be analyzed as an alternative to screening all possible functional polymorphisms in order to find trait-causing alleles. Association studies compare the frequency of marker alleles in unrelated case-control populations, and represent powerful tools for the dissection of complex traits.

Case-Control Populations (Inclusion Criteria)

Population-based association studies do not concern familial inheritance but compare the prevalence of a particular genetic marker, or a set of markers, in case-control populations. They are case-control studies based on comparison of unrelated case (affected or trait positive) individuals and unrelated control (unaffected, trait negative or random) individuals. Preferably the control group is composed of unaffected or trait negative individuals. Further, the control group is ethnically matched to the case population. Moreover, the control group is preferably matched to the case-population for the main known confusion factor for the trait under study (for example age-matched for an age-dependent trait). Ideally, individuals in the two samples are paired in such a way that they are expected to differ only in their disease status. The terms "trait positive population", "case population" and "affected population" are used interchangeably herein.

An important step in the dissection of complex traits using association studies is the choice of case-control populations (see Lander and Schork, 1994). A major step in the choice of case-control populations is the clinical definition of a given trait or phenotype. Any genetic trait may be analyzed by the association method proposed here by carefully selecting the individuals to be included in the trait positive and trait negative phenotypic groups. Four criteria are often useful: clinical phenotype, age at onset, family history and severity. The selection procedure for continuous or quantitative traits (such as blood pressure for example) involves selecting individuals at opposite ends of the phenotype distribution of the trait under study, so as to include in these trait positive and trait negative populations individuals with non-overlapping phenotypes. Preferably, case-control populations consist of phenotypically homogeneous populations. Trait positive and trait negative populations consist of phenotypically uniform populations of individuals representing each between 1 and 98%, preferably between 1 and 80%, more preferably between 1 and 50%, and more preferably between 1 and 30%, most preferably between 1 and 20% of the total population under study, and preferably selected among individuals exhibiting non-overlapping phenotypes. The clearer the difference between the two trait phenotypes, the greater the probability of detecting an association with biallelic markers. The selection of those drastically different but relatively uniform phenotypes enables efficient comparisons in association studies and the possible detection of marked differences at the genetic level, provided that the sample sizes of the populations under study are significant enough.

In preferred embodiments, a first group of between 50 and 300 trait positive individuals, preferably about 100 individuals, are recruited according to their phenotypes. A similar number of control individuals are included in such studies.

Association Analysis

The invention also comprises methods of detecting an association between a genotype and a phenotype, comprising the steps of: a) determining the frequency of at least one *HKLP*-related biallelic marker in a trait positive population according to a genotyping method of the invention; b) determining the frequency of said *HKLP*-related biallelic marker in a control population according to a genotyping method of the invention; and c) determining whether a statistically significant association exists between said genotype and said phenotype. In addition, the methods of detecting an association between a genotype and a phenotype of the invention encompass methods with any further limitation described in this disclosure, or those following, specified alone or in any combination: Optionally, said *HKLP*-related biallelic marker is selected from the group consisting of A1 to A32, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, said *HKLP*-related biallelic marker is selected from the group consisting of A1 to A17, and A20 to A22, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, said *HKLP*-related biallelic marker is selected from the group consisting of A23 and A24, and the complements thereof, or optionally the

biallelic markers in linkage disequilibrium therewith; Optionally, said control population may be a trait negative population, or a random population; Optionally, each of said genotyping steps a) and b) may be performed on a pooled biological sample derived from each of said populations; Optionally, each of said genotyping of steps a) and b) is performed separately on biological samples
5 derived from each individual in said population or a subsample thereof.

The general strategy to perform association studies using biallelic markers derived from a region carrying a candidate gene is to scan two groups of individuals (case-control populations) in order to measure and statistically compare the allele frequencies of the biallelic markers of the present invention in both groups.

10 If a statistically significant association with a trait is identified for at least one or more of the analyzed biallelic markers, one can assume that: either the associated allele is directly responsible for causing the trait (i.e. the associated allele is the trait causing allele), or more likely the associated allele is in linkage disequilibrium with the trait causing allele. The specific characteristics of the associated allele with respect to the candidate gene function usually give further insight into the
15 relationship between the associated allele and the trait (causal or in linkage disequilibrium). If the evidence indicates that the associated allele within the candidate gene is most probably not the trait causing allele but is in linkage disequilibrium with the real trait causing allele, then the trait causing allele can be found by sequencing the vicinity of the associated marker, and performing further association studies with the polymorphisms that are revealed in an iterative manner.

20 Association studies are usually run in two successive steps. In a first phase, the frequencies of a reduced number of biallelic markers from the candidate gene are determined in the trait positive and control populations. In a second phase of the analysis, the position of the genetic loci responsible for the given trait is further refined using a higher density of markers from the relevant region. However, if the candidate gene under study is relatively small in length, as is the case for
25 *HKLP*, a single phase may be sufficient to establish significant associations.

Haplotype Analysis

As described above, when a chromosome carrying a disease allele first appears in a population as a result of either mutation or migration, the mutant allele necessarily resides on a chromosome having a set of linked markers: the ancestral haplotype. This haplotype can be tracked
30 through populations and its statistical association with a given trait can be analyzed.

Complementing single point (allelic) association studies with multi-point association studies also called haplotype studies increases the statistical power of association studies. Thus, a haplotype association study allows one to define the frequency and the type of the ancestral carrier haplotype. A haplotype analysis is important in that it increases the statistical power of an analysis involving
35 individual markers.

In a first stage of a haplotype frequency analysis, the frequency of the possible haplotypes based on various combinations of the identified biallelic markers of the invention is determined.

The haplotype frequency is then compared for distinct populations of trait positive and control individuals. The number of trait positive individuals, which should be, subjected to this analysis to obtain statistically significant results usually ranges between 30 and 300, with a preferred number of individuals ranging between 50 and 150. The same considerations apply to the number of

5 unaffected individuals (or random control) used in the study. The results of this first analysis provide haplotype frequencies in case-control populations, for each evaluated haplotype frequency a p-value and an odd ratio are calculated. If a statistically significant association is found the relative risk for an individual carrying the given haplotype of being affected with the trait under study can be approximated.

10 An additional embodiment of the present invention encompasses methods of detecting an association between a haplotype and a phenotype, comprising the steps of: a) estimating the frequency of at least one haplotype in a trait positive population, according to a method of the invention for estimating the frequency of a haplotype; b) estimating the frequency of said haplotype in a control population, according to a method of the invention for estimating the frequency of a
15 haplotype; and c) determining whether a statistically significant association exists between said haplotype and said phenotype. In addition, the methods of detecting an association between a haplotype and a phenotype of the invention encompass methods with any further limitation described in this disclosure, or those following: Optionally, said *HKLP*-related biallelic marker is selected from the group consisting of A1 to A32, and the complements thereof, or optionally the
20 biallelic markers in linkage disequilibrium therewith; optionally, said *HKLP*-related biallelic marker is selected from the group consisting of A1 to A17, and A20 to A22, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, said *HKLP*-related biallelic marker is selected from the group consisting of A23 and A24, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; Optionally, said
25 control population is a trait negative population, or a random population. Optionally, said method comprises the additional steps of determining the phenotype in said trait positive and said control populations prior to step c).

Interaction Analysis

The biallelic markers of the present invention may also be used to identify patterns of
30 biallelic markers associated with detectable traits resulting from polygenic interactions. The analysis of genetic interaction between alleles at unlinked loci requires individual genotyping using the techniques described herein. The analysis of allelic interaction among a selected set of biallelic markers with appropriate level of statistical significance can be considered as a haplotype analysis. Interaction analysis consists in stratifying the case-control populations with respect to a given
35 haplotype for the first loci and performing a haplotype analysis with the second loci with each subpopulation.

Statistical methods used in association studies are further described below.

Testing For Linkage In The Presence Of Association

The biallelic markers of the present invention may further be used in TDT (transmission/disequilibrium test). TDT tests for both linkage and association and is not affected by population stratification. TDT requires data for affected individuals and their parents or data from unaffected sibs instead of from parents (see Spielmann S. et al., 1993; Schaid D.J. et al., 1996, 5 Spielmann S. and Ewens W.J., 1998). Such combined tests generally reduce the false – positive errors produced by separate analyses.

Statistical methods

In general, any method known in the art to test whether a trait and a genotype show a 10 statistically significant correlation may be used.

1) Methods In Linkage Analysis

Statistical methods and computer programs useful for linkage analysis are well-known to those skilled in the art (see Terwilliger J.D. and Ott J., 1994; Ott J., 1991).

2) Methods To Estimate Haplotype Frequencies In A Population

15 As described above, when genotypes are scored, it is often not possible to distinguish heterozygotes so that haplotype frequencies cannot be easily inferred. When the gametic phase is not known, haplotype frequencies can be estimated from the multilocus genotypic data. Any method known to person skilled in the art can be used to estimate haplotype frequencies (see Lange K., 1997; Weir, B.S., 1996) Preferably, maximum-likelihood haplotype frequencies are computed using 20 an Expectation- Maximization (EM) algorithm (see Dempster et al., 1977; Excoffier L. and Slatkin M., 1995). This procedure is an iterative process aiming at obtaining maximum-likelihood estimates of haplotype frequencies from multi-locus genotype data when the gametic phase is unknown. Haplotype estimations are usually performed by applying the EM algorithm using for example the EM-HAPLO program (Hawley M. E. et al., 1994) or the Arlequin program (Schneider et al., 1997). 25 The EM algorithm is a generalized iterative maximum likelihood approach to estimation and is briefly described below.

Please note that in the present section, “Methods To Estimate Haplotype Frequencies In A Population,” of this text, phenotypes will refer to multi-locus genotypes with unknown phase. Genotypes will refer to known-phase multi-locus genotypes.

30 A sample of N unrelated individuals is typed for K markers. The data observed are the unknown-phase K-locus phenotypes that can be categorized in F different phenotypes. Suppose that we have H underlying possible haplotypes (in case of K biallelic markers, $H=2^K$).

For phenotype j, suppose that c_j genotypes are possible. We thus have the following equation

$$P_j = \sum_{i=1}^{c_j} pr(genotype_i) = \sum_{i=1}^{c_j} pr(h_k, h_l) \quad \text{Equation 1}$$

where P_j is the probability of the phenotype j , h_k and h_l are the two haplotypes constituent the genotype i . Under the Hardy-Weinberg equilibrium, $pr(h_k, h_l)$ becomes:

$$pr(h_k, h_l) = pr(h_k)^2 \text{ if } h_k = h_l, pr(h_k, h_l) = 2 pr(h_k) \cdot pr(h_l) \text{ if } h_k \neq h_l. \quad \text{Equation 2}$$

5 The successive steps of the E-M algorithm can be described as follows:

Starting with initial values of the of haplotypes frequencies, noted $p_1^{(0)}, p_2^{(0)}, \dots, p_H^{(0)}$, these initial values serve to estimate the genotype frequencies (Expectation step) and then estimate another set of haplotype frequencies (Maximization step), noted $p_1^{(1)}, p_2^{(1)}, \dots, p_H^{(1)}$, these two steps are iterated until changes in the sets of haplotypes frequency are very small.

10 A stop criterion can be that the maximum difference between haplotype frequencies between two iterations is less than 10^{-7} . These values can be adjusted according to the desired precision of estimations.

At a given iteration s , the Expectation step consists in calculating the genotypes frequencies by the following equation:

$$pr(genotype_i)^{(s)} = pr(phenotype_j) \cdot pr(genotype_i | phenotype_j)^{(s)} \\ = \frac{n_j}{N} \cdot \frac{pr(h_k, h_l)^{(s)}}{P_j^{(s)}} \quad \text{Equation 3}$$

where genotype i occurs in phenotype j , and where h_k and h_l constitute genotype i . Each probability is derived according to eq. 1, and eq. 2 described above.

Then the Maximization step simply estimates another set of haplotype frequencies given the 20 genotypes frequencies. This approach is also known as the gene-counting method (Smith, 1957).

$$p_t^{(s+1)} = \frac{1}{2} \sum_{j=1}^F \sum_{i=1}^{c_j} \delta_{it} \cdot pr(genotype_i)^{(s)} \quad \text{Equation 4}$$

Where δ_{it} is an indicator variable which count the number of time haplotype t in genotype i . It takes the values of 0, 1 or 2.

To ensure that the estimation finally obtained is the maximum-likelihood estimation several 25 values of departures are required. The estimations obtained are compared and if they are different the estimations leading to the best likelihood are kept.

3) Methods To Calculate Linkage Disequilibrium Between Markers

A number of methods can be used to calculate linkage disequilibrium between any two genetic positions, in practice linkage disequilibrium is measured by applying a statistical association test to haplotype data taken from a population.

- 5 Linkage disequilibrium between any pair of biallelic markers comprising at least one of the biallelic markers of the present invention (M_i, M_j) having alleles (a_i/b_i) at marker M_i and alleles (a_j/b_j) at marker M_j can be calculated for every allele combination ($a_i, a_j, a_i, b_j, b_i, a_j$ and b_i, b_j), according to the Piazza formula:

$$\Delta_{aij} = \sqrt{\theta_4} - \sqrt{(\theta_4 + \theta_3)(\theta_4 + \theta_2)}, \text{ where:}$$

- 10 $\theta_4 = - - =$ frequency of genotypes not having allele a_i at M_i and not having allele a_j at M_j
 $\theta_3 = - + =$ frequency of genotypes not having allele a_i at M_i and having allele a_j at M_j
 $\theta_2 = + - =$ frequency of genotypes having allele a_i at M_i and not having allele a_j at M_j

- Linkage disequilibrium (LD) between pairs of biallelic markers (M_i, M_j) can also be
 15 calculated for every allele combination ($a_i, a_j, a_i, b_j, b_i, a_j$ and b_i, b_j), according to the maximum-likelihood estimate (MLE) for delta (the composite genotypic disequilibrium coefficient), as described by Weir (Weir B. S., 1996). The MLE for the composite linkage disequilibrium is:

$$D_{aij} = (2n_1 + n_2 + n_3 + n_4/2)/N - 2(pr(a_i) \cdot pr(a_j))$$

- Where $n_1 = \Sigma$ phenotype ($a_i/a_i, a_j/a_j$), $n_2 = \Sigma$ phenotype ($a_i/a_i, a_j/b_j$), $n_3 = \Sigma$ phenotype ($a_i/b_i, a_j/a_j$), $n_4 = \Sigma$ phenotype ($a_i/b_i, a_j/b_j$) and N is the number of individuals in the sample.
 20

This formula allows linkage disequilibrium between alleles to be estimated when only genotype, and not haplotype, data are available.

- Another means of calculating the linkage disequilibrium between markers is as follows. For
 25 a couple of biallelic markers, $M_i (a_i/b_i)$ and $M_j (a_j/b_j)$, fitting the Hardy-Weinberg equilibrium, one can estimate the four possible haplotype frequencies in a given population according to the approach described above.

The estimation of gametic disequilibrium between a_i and a_j is simply:

$$D_{aij} = pr(haplotype(a_i, a_j)) - pr(a_i) \cdot pr(a_j).$$

- 30 Where $pr(a_i)$ is the probability of allele a_i and $pr(a_j)$ is the probability of allele a_j and where $pr(haplotype(a_i, a_j))$ is estimated as in Equation 3 above.

For a couple of biallelic marker only one measure of disequilibrium is necessary to describe the association between M_i and M_j .

Then a normalized value of the above is calculated as follows:

- 35 $D'_{aij} = D_{aij} / \max(-pr(a_i) \cdot pr(a_j), -pr(b_i) \cdot pr(b_j))$ with $D_{aij} < 0$
 $D'_{aij} = D_{aij} / \max(pr(b_i) \cdot pr(a_j), pr(a_i) \cdot pr(b_j))$ with $D_{aij} > 0$

The skilled person will readily appreciate that other linkage disequilibrium calculation methods can be used.

Linkage disequilibrium among a set of biallelic markers having an adequate heterozygosity rate can be determined by genotyping between 50 and 1000 unrelated individuals, preferably
5 between 75 and 200, more preferably around 100.

4) Testing For Association

Methods for determining the statistical significance of a correlation between a phenotype and a genotype, in this case an allele at a biallelic marker or a haplotype made up of such alleles, may be determined by any statistical test known in the art and with any accepted threshold of
10 statistical significance being required. The application of particular methods and thresholds of significance are well within the skill of the ordinary practitioner of the art.

Testing for association is performed by determining the frequency of a biallelic marker allele in case and control populations and comparing these frequencies with a statistical test to determine if there is a statistically significant difference in frequency which would indicate a correlation between
15 the trait and the biallelic marker allele under study. Similarly, a haplotype analysis is performed by estimating the frequencies of all possible haplotypes for a given set of biallelic markers in case and control populations, and comparing these frequencies with a statistical test to determine if there is a statistically significant correlation between the haplotype and the phenotype (trait) under study. Any statistical tool useful to test for a statistically significant association between a genotype and a
20 phenotype may be used. Preferably the statistical test employed is a chi-square test with one degree of freedom. A P-value is calculated (the P-value is the probability that a statistic as large or larger than the observed one would occur by chance).

Statistical Significance

In preferred embodiments, significance for diagnosis purposes, either as a positive basis for
25 further diagnostic tests or as a preliminary starting point for early preventive therapy, the p value related to a biallelic marker association is preferably about 1×10^{-2} or less, more preferably about 1×10^{-4} or less, for a single biallelic marker analysis and about 1×10^{-3} or less, still more preferably 1×10^{-6} or less and most preferably of about 1×10^{-8} or less, for a haplotype analysis involving two or more markers. These values are believed to be applicable to any association studies involving single
30 or multiple marker combinations.

The skilled person can use the range of values set forth above as a starting point in order to carry out association studies with biallelic markers of the present invention. In doing so, significant associations between the biallelic markers of the present invention and a trait can be revealed and used for diagnosis and drug screening purposes.

Phenotypic Permutation

In order to confirm the statistical significance of the first stage haplotype analysis described above, it might be suitable to perform further analyses in which genotyping data from case-control individuals are pooled and randomized with respect to the trait phenotype. Each individual

5 genotyping data is randomly allocated to two groups, which contain the same number of individuals as the case-control populations used to compile the data obtained in the first stage. A second stage haplotype analysis is preferably run on these artificial groups, preferably for the markers included in the haplotype of the first stage analysis showing the highest relative risk coefficient. This experiment is reiterated preferably at least between 100 and 10000 times. The repeated iterations

10 allow the determination of the probability to obtain the tested haplotype by chance.

Assessment Of Statistical Association

To address the problem of false positives similar analysis may be performed with the same case-control populations in random genomic regions. Results in random regions and the candidate region are compared as described in a co-pending US Provisional Patent Application entitled

15 "Methods, Software And Apparati For Identifying Genomic Regions Harboring A Gene Associated With A Detectable Trait," U.S. Serial Number 60/107,986, filed November 10, 1998, the contents of which are incorporated herein by reference.

5) Evaluation Of Risk Factors

The association between a risk factor (in genetic epidemiology the risk factor is the presence

20 or the absence of a certain allele or haplotype at marker loci) and a disease is measured by the odds ratio (OR) and by the relative risk (RR). If $P(R^+)$ is the probability of developing the disease for individuals with R and $P(R^-)$ is the probability for individuals without the risk factor, then the relative risk is simply the ratio of the two probabilities, that is:

$$RR = P(R^+)/P(R^-)$$

25 In case-control studies, direct measures of the relative risk cannot be obtained because of the sampling design. However, the odds ratio allows a good approximation of the relative risk for low-incidence diseases and can be calculated:

$$OR = \left[\frac{F^+}{1 - F^+} \right] / \left[\frac{F^-}{(1 - F^-)} \right]$$

$$OR = (F^+ / (1 - F^+)) / (F^- / (1 - F^-))$$

F^+ is the frequency of the exposure to the risk factor in cases and F^- is the frequency of the

30 exposure to the risk factor in controls. F^+ and F^- are calculated using the allelic or haplotype frequencies of the study and further depend on the underlying genetic model (dominant, recessive, additive...).

One can further estimate the attributable risk (AR) which describes the proportion of individuals in a population exhibiting a trait due to a given risk factor. This measure is important in

quantifying the role of a specific factor in disease etiology and in terms of the public health impact of a risk factor. The public health relevance of this measure lies in estimating the proportion of cases of disease in the population that could be prevented if the exposure of interest were absent.

AR is determined as follows:

$$5 \quad AR = P_E (RR-1) / (P_E (RR-1)+1)$$

AR is the risk attributable to a biallelic marker allele or a biallelic marker haplotype. P_E is the frequency of exposure to an allele or a haplotype within the population at large; and RR is the relative risk which, is approximated with the odds ratio when the trait under study has a relatively low incidence in the general population.

10 **Identification Of Biallelic Markers In Linkage Disequilibrium With The Biallelic Markers of the Invention**

Once a first biallelic marker has been identified in a genomic region of interest, the practitioner of ordinary skill in the art, using the teachings of the present invention, can easily identify additional biallelic markers in linkage disequilibrium with this first marker. As mentioned
 15 before any marker in linkage disequilibrium with a first marker associated with a trait will be associated with the trait. Therefore, once an association has been demonstrated between a given biallelic marker and a trait, the discovery of additional biallelic markers associated with this trait is of great interest in order to increase the density of biallelic markers in this particular region. The causal gene or mutation will be found in the vicinity of the marker or set of markers showing the
 20 highest correlation with the trait.

Identification of additional markers in linkage disequilibrium with a given marker involves:
 (a) amplifying a genomic fragment comprising a first biallelic marker from a plurality of individuals;
 (b) identifying of second biallelic markers in the genomic region harboring said first biallelic marker; (c) conducting a linkage disequilibrium analysis between said first biallelic marker and
 25 second biallelic markers; and (d) selecting said second biallelic markers as being in linkage disequilibrium with said first marker. Subcombinations comprising steps (b) and (c) are also contemplated.

Methods to identify biallelic markers and to conduct linkage disequilibrium analysis are described herein and can be carried out by the skilled person without undue experimentation. The
 30 present invention then also concerns biallelic markers which are in linkage disequilibrium with the specific biallelic markers A1 to A32 and which are expected to present similar characteristics in terms of their respective association with a given trait. In a preferred embodiment, the invention concerns biallelic markers which are in linkage disequilibrium with the specific biallelic markers

Identification Of Functional Mutations

35 Mutations in the *HKLP* gene which are responsible for a detectable phenotype or trait may be identified by comparing the sequences of the *HKLP* gene from trait positive and control

individuals. Once a positive association is confirmed with a biallelic marker of the present invention, the identified locus can be scanned for mutations. In a preferred embodiment, functional regions such as exons and splice sites, promoters and other regulatory regions of the *HKLP* gene are scanned for mutations. In a preferred embodiment the sequence of the *HKLP* gene is compared in
5 trait positive and control individuals. Preferably, trait positive individuals carry the haplotype shown to be associated with the trait and trait negative individuals do not carry the haplotype or allele associated with the trait. The detectable trait or phenotype may comprise a variety of manifestations of altered *HKLP* function.

The mutation detection procedure is essentially similar to that used for biallelic marker
10 identification. The method used to detect such mutations generally comprises the following steps:

- amplification of a region of the *HKLP* gene comprising a biallelic marker or a group of biallelic markers associated with the trait from DNA samples of trait positive patients and trait-negative controls;
- sequencing of the amplified region;
- 15 - comparison of DNA sequences from trait positive and control individuals;
- determination of mutations specific to trait-positive patients.

In one embodiment, said biallelic marker is selected from the group consisting of A1 to A32, and the complements thereof. It is preferred that candidate polymorphisms be then verified by screening a larger population of cases and controls by means of any genotyping procedure such as
20 those described herein, preferably using a microsequencing technique in an individual test format. Polymorphisms are considered as candidate mutations when present in cases and controls at frequencies compatible with the expected association results. Polymorphisms are considered as candidate "trait-causing" mutations when they exhibit a statistically significant correlation with the detectable phenotype.

25

Recombinant Vectors

The term "vector" is used herein to designate either a circular or a linear DNA or RNA molecule, which is either double-stranded or single-stranded, and which comprise at least one polynucleotide of interest that is sought to be transferred in a cell host or in a unicellular or multicellular host organism.

30 The present invention encompasses a family of recombinant vectors that comprise a regulatory polynucleotide derived from the *HKLP* genomic sequence, and/or a coding polynucleotide from either the *HKLP* genomic sequence or the cDNA sequence.

Generally, a recombinant vector of the invention may comprise any of the polynucleotides described herein, including regulatory sequences, coding sequences and polynucleotide constructs,
35 as well as any *HKLP* primer or probe as defined above. More particularly, the recombinant vectors of the present invention can comprise any of the polynucleotides described in the "Genomic

Sequences Of tThe *HKLP* Gene” section, the “*HKLP* cDNA Sequences” section, the “Coding Regions” section, the “Polynucleotide constructs” section, and the “Oligonucleotide Probes And Primers” section.

In a first preferred embodiment, a recombinant vector of the invention is used to amplify the
5 inserted polynucleotide derived from a *HKLP* genomic sequence of SEQ ID Nos 1 and 2 or a *HKLP* cDNA, for example the cDNA of SEQ ID No 3 in a suitable cell host, this polynucleotide being amplified at every time that the recombinant vector replicates.

A second preferred embodiment of the recombinant vectors according to the invention consists of expression vectors comprising either a regulatory polynucleotide or a coding nucleic acid
10 of the invention, or both. Within certain embodiments, expression vectors are employed to express the *HKLP* polypeptide which can be then purified and, for example be used in ligand screening assays or as an immunogen in order to raise specific antibodies directed against the *HKLP* protein. In other embodiments, the expression vectors are used for constructing transgenic animals and also for gene therapy. Expression requires that appropriate signals are provided in the vectors, said
15 signals including various regulatory elements, such as enhancers/promoters from both viral and mammalian sources that drive expression of the genes of interest in host cells. Dominant drug selection markers for establishing permanent, stable cell clones expressing the products are generally included in the expression vectors of the invention, as they are elements that link expression of the drug selection markers to expression of the polypeptide.

20 More particularly, the present invention relates to expression vectors which include nucleic acids encoding a *HKLP* protein, preferably the *HKLP* protein of the amino acid sequence of SEQ ID No 4 or variants or fragments thereof.

The invention also pertains to a recombinant expression vector useful for the expression of the *HKLP* coding sequence, wherein said vector comprises a nucleic acid of SEQ ID No 3.

25 Recombinant vectors comprising a nucleic acid containing a *HKLP*-related biallelic marker is also part of the invention. In a preferred embodiment, said biallelic marker is selected from the group consisting of A1 to A32, and the complements thereof.

Some of the elements which can be found in the vectors of the present invention are described in further detail in the following sections.

30 1. General features of the expression vectors of the invention

A recombinant vector according to the invention comprises, but is not limited to, a YAC (Yeast Artificial Chromosome), a BAC (Bacterial Artificial Chromosome), a phage, a phagemid, a cosmid, a plasmid or even a linear DNA molecule which may consist of a chromosomal, non-chromosomal, semi-synthetic and synthetic DNA. Such a recombinant vector can comprise a
35 transcriptional unit comprising an assembly of:

(1) a genetic element or elements having a regulatory role in gene expression, for example promoters or enhancers. Enhancers are cis-acting elements of DNA, usually from about 10 to 300 bp in length that act on the promoter to increase the transcription.

(2) a structural or coding sequence which is transcribed into mRNA and eventually translated into a polypeptide, said structural or coding sequence being operably linked to the regulatory elements described in (1); and

(3) appropriate transcription initiation and termination sequences. Structural units intended for use in yeast or eukaryotic expression systems preferably include a leader sequence enabling extracellular secretion of translated protein by a host cell. Alternatively, when a recombinant protein is expressed without a leader or transport sequence, it may include a N-terminal residue. This residue may or may not be subsequently cleaved from the expressed recombinant protein to provide a final product.

Generally, recombinant expression vectors will include origins of replication, selectable markers permitting transformation of the host cell, and a promoter derived from a highly expressed gene to direct transcription of a downstream structural sequence. The heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences, and preferably a leader sequence capable of directing secretion of the translated protein into the periplasmic space or the extracellular medium. In a specific embodiment wherein the vector is adapted for transfecting and expressing desired sequences in mammalian host cells, preferred vectors will comprise an origin of replication in the desired host, a suitable promoter and enhancer, and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5'-flanking non-transcribed sequences. DNA sequences derived from the SV40 viral genome, for example SV40 origin, early promoter, enhancer, splice and polyadenylation sites may be used to provide the required non-transcribed genetic elements.

The *in vivo* expression of a HKLP polypeptide of SEQ ID No 4 or fragments or variants thereof may be useful in order to correct a genetic defect related to the expression of the native gene in a host organism or to the production of a biologically inactive HKLP protein.

Consequently, the present invention also deals with recombinant expression vectors mainly designed for the *in vivo* production of the HKLP polypeptide of SEQ ID No 4 or fragments or variants thereof by the introduction of the appropriate genetic material in the organism of the patient to be treated. This genetic material may be introduced *in vitro* in a cell that has been previously extracted from the organism, the modified cell being subsequently reintroduced in the said organism, directly *in vivo* into the appropriate tissue.

2. Regulatory Elements

Promoters

The suitable promoter regions used in the expression vectors according to the present invention are chosen taking into account the cell host in which the heterologous gene has to be expressed. The particular promoter employed to control the expression of a nucleic acid sequence of interest is not believed to be important, so long as it is capable of directing the expression of the nucleic acid in the targeted cell. Thus, where a human cell is targeted, it is preferable to position the nucleic acid coding region adjacent to and under the control of a promoter that is capable of being expressed in a human cell, such as, for example, a human or a viral promoter.

A suitable promoter may be heterologous with respect to the nucleic acid for which it controls the expression or alternatively can be endogenous to the native polynucleotide containing the coding sequence to be expressed. Additionally, the promoter is generally heterologous with respect to the recombinant vector sequences within which the construct promoter/coding sequence has been inserted.

Promoter regions can be selected from any desired gene using, for example, CAT (chloramphenicol transferase) vectors and more preferably pKK232-8 and pCM7 vectors.

Preferred bacterial promoters are the LacI, LacZ, the T3 or T7 bacteriophage RNA polymerase promoters, the gpt, lambda PR, PL and trp promoters (EP 0036776), the polyhedrin promoter, or the p10 protein promoter from baculovirus (Kit Novagen) (Smith et al., 1983; O'Reilly et al., 1992), the lambda PR promoter or also the trc promoter.

Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-L. Selection of a convenient vector and promoter is well within the level of ordinary skill in the art.

The choice of a promoter is well within the ability of a person skilled in the field of genetic engineering. For example, one may refer to the book of Sambrook et al.(1989) or also to the procedures described by Fuller et al.(1996).

Other regulatory elements

Where a cDNA insert is employed, one will typically desire to include a polyadenylation signal to effect proper polyadenylation of the gene transcript. The nature of the polyadenylation signal is not believed to be crucial to the successful practice of the invention, and any such sequence may be employed such as human growth hormone and SV40 polyadenylation signals. Also contemplated as an element of the expression cassette is a terminator. These elements can serve to enhance message levels and to minimize read through from the cassette into other sequences.

3. Selectable Markers

Such markers would confer an identifiable change to the cell permitting easy identification of cells containing the expression construct. The selectable marker genes for selection of

transformed host cells are preferably dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, TRP1 for *S. cerevisiae* or tetracycline, rifampicin or ampicillin resistance in *E. coli*, or levan saccharase for mycobacteria, this latter marker being a negative selection marker.

4. Preferred Vectors.

5 Bacterial vectors

As a representative but non-limiting example, useful expression vectors for bacterial use can comprise a selectable marker and a bacterial origin of replication derived from commercially available plasmids comprising genetic elements of pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia, Uppsala, Sweden), and GEM1 (Promega
10 Biotec, Madison, WI, USA).

Large numbers of other suitable vectors are known to those of skill in the art, and commercially available, such as the following bacterial vectors: pQE70, pQE60, pQE-9 (Qiagen), pbs, pD10, phagescript, psiX174, pbluescript SK, pbsks, pNH8A, pNH16A, pNH18A, pNH46A (Stratagene); ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 (Pharmacia); pWLNEO, pSV2CAT,
15 pOG44, pXT1, pSG (Stratagene); pSVK3, pBPV, pMSG, pSVL (Pharmacia); pQE-30 (QIAexpress).

Bacteriophage vectors

The P1 bacteriophage vector may contain large inserts ranging from about 80 to about 100 kb.

20 The construction of P1 bacteriophage vectors such as p158 or p158/neo8 are notably described by Sternberg (1992, 1994). Recombinant P1 clones comprising *HKLP* nucleotide sequences may be designed for inserting large polynucleotides of more than 40 kb (Linton et al., 1993). To generate P1 DNA for transgenic experiments, a preferred protocol is the protocol described by McCormick et al.(1994). Briefly, *E. coli* (preferably strain NS3529) harboring the P1
25 plasmid are grown overnight in a suitable broth medium containing 25 µg/ml of kanamycin. The P1 DNA is prepared from the *E. coli* by alkaline lysis using the Qiagen Plasmid Maxi kit (Qiagen, Chatsworth, CA, USA), according to the manufacturer's instructions. The P1 DNA is purified from the bacterial lysate on two Qiagen-tip 500 columns, using the washing and elution buffers contained in the kit. A phenol/chloroform extraction is then performed before precipitating the DNA with 70%
30 ethanol. After solubilizing the DNA in TE (10 mM Tris-HCl, pH 7.4, 1 mM EDTA), the concentration of the DNA is assessed by spectrophotometry.

When the goal is to express a P1 clone comprising *HKLP* nucleotide sequences in a transgenic animal, typically in transgenic mice, it is desirable to remove vector sequences from the P1 DNA fragment, for example by cleaving the P1 DNA at rare-cutting sites within the P1
35 polylinker (*Sfi*I, *Not*I or *Sal*I). The P1 insert is then purified from vector sequences on a pulsed-field agarose gel, using methods similar using methods similar to those originally reported for the

isolation of DNA from YACs (Schedl et al., 1993a; Peterson et al., 1993). At this stage, the resulting purified insert DNA can be concentrated, if necessary, on a Millipore Ultrafree-MC Filter Unit (Millipore, Bedford, MA, USA – 30,000 molecular weight limit) and then dialyzed against microinjection buffer (10 mM Tris-HCl, pH 7.4; 250 μ M EDTA) containing 100 mM NaCl, 30 μ M spermine, 70 μ M spermidine on a microdialysis membrane (type VS, 0.025 μ M from Millipore). The intactness of the purified P1 DNA insert is assessed by electrophoresis on 1% agarose (Sea Kem GTG; FMC Bio-products) pulse-field gel and staining with ethidium bromide.

Baculovirus vectors

A suitable vector for the expression of the HKLP polypeptide of SEQ ID No 3 or fragments or variants thereof is a baculovirus vector that can be propagated in insect cells and in insect cell lines. A specific suitable host vector system is the pVL1392/1393 baculovirus transfer vector (Pharmingen) that is used to transfect the SF9 cell line (ATCC N^oCRL 1711) which is derived from *Spodoptera frugiperda*.

Other suitable vectors for the expression of the HKLP polypeptide of SEQ ID No 3 or fragments or variants thereof in a baculovirus expression system include those described by Chai et al.(1993), Vlasak et al.(1983) and Lenhard et al.(1996).

Viral vectors

In one specific embodiment, the vector is derived from an adenovirus. Preferred adenovirus vectors according to the invention are those described by Feldman and Steg (1996) or Ohno et al.(1994). Another preferred recombinant adenovirus according to this specific embodiment of the present invention is the human adenovirus type 2 or 5 (Ad 2 or Ad 5) or an adenovirus of animal origin (French patent application N^o FR-93.05954).

Retrovirus vectors and adeno-associated virus vectors are generally understood to be the recombinant gene delivery systems of choice for the transfer of exogenous polynucleotides *in vivo* , particularly to mammals, including humans. These vectors provide efficient delivery of genes into cells, and the transferred nucleic acids are stably integrated into the chromosomal DNA of the host.

Particularly preferred retroviruses for the preparation or construction of retroviral *in vitro* or *in vitro* gene delivery vehicles of the present invention include retroviruses selected from the group consisting of Mink-Cell Focus Inducing Virus, Murine Sarcoma Virus, Reticuloendotheliosis virus and Rous Sarcoma virus. Particularly preferred Murine Leukemia Viruses include the 4070A and the 1504A viruses, Abelson (ATCC No VR-999), Friend (ATCC No VR-245), Gross (ATCC No VR-590), Rauscher (ATCC No VR-998) and Moloney Murine Leukemia Virus (ATCC No VR-190; PCT Application No WO 94/24298). Particularly preferred Rous Sarcoma Viruses include Bryan high titer (ATCC Nos VR-334, VR-657, VR-726, VR-659 and VR-728). Other preferred retroviral vectors are those described in Roth et al.(1996), PCT Application No WO 93/25234, PCT Application No WO 94/ 06920, Roux et al., 1989, Julan et al., 1992 and Neda et al., 1991.

Yet another viral vector system that is contemplated by the invention consists in the adeno-associated virus (AAV). The adeno-associated virus is a naturally occurring defective virus that requires another virus, such as an adenovirus or a herpes virus, as a helper virus for efficient replication and a productive life cycle (Muzyczka et al., 1992). It is also one of the few viruses that may integrate its DNA into non-dividing cells, and exhibits a high frequency of stable integration (Flotte et al., 1992; Samulski et al., 1989; McLaughlin et al., 1989). One advantageous feature of AAV derives from its reduced efficacy for transducing primary cells relative to transformed cells.

BAC vectors

The bacterial artificial chromosome (BAC) cloning system (Shizuya et al., 1992) has been developed to stably maintain large fragments of genomic DNA (100-300 kb) in *E. coli*. A preferred BAC vector consists of pBeloBAC11 vector that has been described by Kim et al. (1996). BAC libraries are prepared with this vector using size-selected genomic DNA that has been partially digested using enzymes that permit ligation into either the *Bam* HI or *Hind*III sites in the vector. Flanking these cloning sites are T7 and SP6 RNA polymerase transcription initiation sites that can be used to generate end probes by either RNA transcription or PCR methods. After the construction of a BAC library in *E. coli*, BAC DNA is purified from the host cell as a supercoiled circle. Converting these circular molecules into a linear form precedes both size determination and introduction of the BACs into recipient cells. The cloning site is flanked by two *Not* I sites, permitting cloned segments to be excised from the vector by *Not* I digestion. Alternatively, the DNA insert contained in the pBeloBAC11 vector may be linearized by treatment of the BAC vector with the commercially available enzyme lambda terminase that leads to the cleavage at the unique *cos*N site, but this cleavage method results in a full length BAC clone containing both the insert DNA and the BAC sequences.

5. Delivery Of The Recombinant Vectors

In order to effect expression of the polynucleotides and polynucleotide constructs of the invention, these constructs must be delivered into a cell. This delivery may be accomplished *in vitro*, as in laboratory procedures for transforming cell lines, or *in vivo* or *ex vivo*, as in the treatment of certain diseases states.

One mechanism is viral infection where the expression construct is encapsulated in an infectious viral particle.

Several non-viral methods for the transfer of polynucleotides into cultured mammalian cells are also contemplated by the present invention, and include, without being limited to, calcium phosphate precipitation (Graham et al., 1973; Chen et al., 1987;), DEAE-dextran (Gopal, 1985), electroporation (Tur-Kaspa et al., 1986; Potter et al., 1984), direct microinjection (Harland et al., 1985), DNA-loaded liposomes (Nicolau et al., 1982; Fraley et al., 1979), and receptor-mediated

transfection (Wu and Wu, 1987; 1988). Some of these techniques may be successfully adapted for *in vivo* or *ex vivo* use.

Once the expression polynucleotide has been delivered into the cell, it may be stably integrated into the genome of the recipient cell. This integration may be in the cognate location and orientation via homologous recombination (gene replacement) or it may be integrated in a random, non specific location (gene augmentation). In yet further embodiments, the nucleic acid may be stably maintained in the cell as a separate, episomal segment of DNA. Such nucleic acid segments or "episomes" encode sequences sufficient to permit maintenance and replication independent of or in synchronization with the host cell cycle.

One specific embodiment for a method for delivering a protein or peptide to the interior of a cell of a vertebrate *in vivo* comprises the step of introducing a preparation comprising a physiologically acceptable carrier and a naked polynucleotide operatively coding for the polypeptide of interest into the interstitial space of a tissue comprising the cell, whereby the naked polynucleotide is taken up into the interior of the cell and has a physiological effect. This is particularly applicable for transfer *in vitro* but it may be applied to *in vivo* as well.

Compositions for use *in vitro* and *in vivo* comprising a "naked" polynucleotide are described in PCT application N° WO 90/11092 (Vical Inc.) and also in PCT application No. WO 95/11307 (Institut Pasteur, INSERM, Université d'Ottawa) as well as in the articles of Tacson et al.(1996) and of Huygen et al.(1996).

In still another embodiment of the invention, the transfer of a naked polynucleotide of the invention, including a polynucleotide construct of the invention, into cells may be proceeded with a particle bombardment (biolistic), said particles being DNA-coated microprojectiles accelerated to a high velocity allowing them to pierce cell membranes and enter cells without killing them, such as described by Klein et al.(1987).

In a further embodiment, the polynucleotide of the invention may be entrapped in a liposome (Ghosh and Bacchawat, 1991; Wong et al., 1980; Nicolau et al., 1987)

In a specific embodiment, the invention provides a composition for the *in vivo* production of the HKLP protein or polypeptide described herein. It comprises a naked polynucleotide operatively coding for this polypeptide, in solution in a physiologically acceptable carrier, and suitable for introduction into a tissue to cause cells of the tissue to express the said protein or polypeptide.

The amount of vector to be injected to the desired host organism varies according to the site of injection. As an indicative dose, it will be injected between 0,1 and 100 µg of the vector in an animal body, preferably a mammal body, for example a mouse body.

In another embodiment of the vector according to the invention, it may be introduced *in vitro* in a host cell, preferably in a host cell previously harvested from the animal to be treated and more preferably a somatic cell such as a muscle cell. In a subsequent step, the cell that has been transformed with the vector coding for the desired HKLP polypeptide or the desired fragment

thereof is reintroduced into the animal body in order to deliver the recombinant protein within the body either locally or systemically.

Cell Hosts

Another object of the invention consists of a host cell that has been transformed or
 5 transfected with one of the polynucleotides described herein, and in particular a polynucleotide either comprising a *HKLP* regulatory polynucleotide or the coding sequence of the *HKLP* polypeptide selected from the group consisting of SEQ ID Nos 1-3 or a fragment or a variant thereof. Also included are host cells that are transformed (prokaryotic cells) or that are transfected (eukaryotic cells) with a recombinant vector such as one of those described above. More
 10 particularly, the cell hosts of the present invention can comprise any of the polynucleotides described in the "Genomic Sequences Of the *HKLP* Gene" section, the "*HKLP* cDNA Sequences" section, the "Coding Regions" section, the "Polynucleotide constructs" section, the "Oligonucleotide Probes And Primers" section and the "Recombinant Vectors" section.

A further recombinant cell host according to the invention comprises a polynucleotide
 15 containing a biallelic marker selected from the group consisting of A1 to A32, and the complements thereof.

Preferred host cells used as recipients for the expression vectors of the invention are the following:

- a) Prokaryotic host cells: *Escherichia coli* strains (I.E.DH5- α strain), *Bacillus subtilis*,
 20 *Salmonella typhimurium*, and strains from species like *Pseudomonas*, *Streptomyces* and *Staphylococcus*.
- b) Eukaryotic host cells: HeLa cells (ATCC N^oCCL2; N^oCCL2.1; N^oCCL2.2), Cv 1 cells (ATCC N^oCCL70), COS cells (ATCC N^oCRL1650; N^oCRL1651), Sf-9 cells (ATCC N^oCRL1711), C127 cells (ATCC N^o CRL-1804), 3T3 (ATCC N^o CRL-6361), CHO (ATCC N^o CCL-61), human
 25 kidney 293. (ATCC N^o 45504; N^o CRL-1573) and BHK (ECACC N^o 84100501; N^o 84111301).
- c) Other mammalian host cells.

The *HKLP* gene expression in mammalian, and typically human, cells may be rendered defective, or alternatively it may be proceeded with the insertion of a *HKLP* genomic or cDNA sequence with the replacement of the *HKLP* gene counterpart in the genome of an animal cell by a
 30 *HKLP* polynucleotide according to the invention. These genetic alterations may be generated by homologous recombination events using specific DNA constructs that have been previously described.

One kind of cell hosts that may be used are mammal zygotes, such as murine zygotes. For example, murine zygotes may undergo microinjection with a purified DNA molecule of interest, for
 35 example a purified DNA molecule that has previously been adjusted to a concentration range from 1 ng/ml –for BAC inserts- 3 ng/ μ l –for P1 bacteriophage inserts- in 10 mM Tris-HCl, pH 7.4, 250 μ M

EDTA containing 100 mM NaCl, 30 μ M spermine, and 70 μ M spermidine. When the DNA to be microinjected has a large size, polyamines and high salt concentrations can be used in order to avoid mechanical breakage of this DNA, as described by Schedl et al (1993b).

Anyone of the polynucleotides of the invention, including the DNA constructs described
5 herein, may be introduced in an embryonic stem (ES) cell line, preferably a mouse ES cell line. ES cell lines are derived from pluripotent, uncommitted cells of the inner cell mass of pre-implantation blastocysts. Preferred ES cell lines are the following: ES-E14TG2a (ATCC n° CRL-1821), ES-D3 (ATCC n° CRL1934 and n° CRL-11632), YS001 (ATCC n° CRL-11776), 36.5 (ATCC n° CRL-11116). To maintain ES cells in an uncommitted state, they are cultured in the presence of growth
10 inhibited feeder cells which provide the appropriate signals to preserve this embryonic phenotype and serve as a matrix for ES cell adherence. Preferred feeder cells consist of primary embryonic fibroblasts that are established from tissue of day 13- day 14 embryos of virtually any mouse strain, that are maintained in culture, such as described by Abbondanzo et al.(1993) and are inhibited in growth by irradiation, such as described by Robertson (1987), or by the presence of an inhibitory
15 concentration of LIF, such as described by Pease and Williams (1990).

The constructs in the host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence.

Following transformation of a suitable host and growth of the host to an appropriate cell density, the selected promoter is induced by appropriate means, such as temperature shift or
20 chemical induction, and cells are cultivated for an additional period.

Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

Microbial cells employed in the expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing
25 agents. Such methods are well known by the skill artisan.

Transgenic Animals

The terms "transgenic animals" or "host animals" are used herein designate animals that have their genome genetically and artificially manipulated so as to include one of the nucleic acids according to the invention. Preferred animals are non-human mammals and include those belonging
30 to a genus selected from *Mus* (e.g. mice), *Rattus* (e.g. rats) and *Oryctogalus* (e.g. rabbits) which have their genome artificially and genetically altered by the insertion of a nucleic acid according to the invention. In one embodiment, the invention encompasses non-human host mammals and animals comprising a recombinant vector of the invention or a HKLP gene disrupted by homologous recombination with a knock out vector.

35 The transgenic animals of the invention all include within a plurality of their cells a cloned recombinant or synthetic DNA sequence, more specifically one of the purified or isolated nucleic

acids comprising a *HKLP* coding sequence, a *HKLP* regulatory polynucleotide, a polynucleotide construct, or a DNA sequence encoding an antisense polynucleotide such as described in the present specification.

Generally, a transgenic animal according to the present invention comprises any one of the polynucleotides, the recombinant vectors and the cell hosts described in the present invention. More particularly, the transgenic animals of the present invention can comprise any of the polynucleotides described in the "Genomic Sequences Of the *HKLP* Gene" section, the "*HKLP* cDNA Sequences" section, the "Coding Regions" section, the "Polynucleotide constructs" section, the "Oligonucleotide Probes And Primers" section, the "Recombinant Vectors" section and the "Cell Hosts" section.

A further transgenic animal according to the invention contains in their somatic cells and/or in their germ line cells a polynucleotide comprising a biallelic marker selected from the group consisting of A1 to A32, and the complements thereof.

In a first preferred embodiment, these transgenic animals may be good experimental models in order to study the diverse pathologies related to cell differentiation, in particular concerning the transgenic animals within the genome of which has been inserted one or several copies of a polynucleotide encoding a native *HKLP* protein, or alternatively a mutant *HKLP* protein.

In a second preferred embodiment, these transgenic animals may express a desired polypeptide of interest under the control of the regulatory polynucleotides of the *HKLP* gene, leading to good yields in the synthesis of this protein of interest, and eventually a tissue specific expression of this protein of interest.

The design of the transgenic animals of the invention may be made according to the conventional techniques well known from the one skilled in the art. For more details regarding the production of transgenic animals, and specifically transgenic mice, it may be referred to US Patents Nos 4,873,191, issued Oct. 10, 1989; 5,464,764 issued Nov 7, 1995; and 5,789,215, issued Aug 4, 1998; these documents being herein incorporated by reference to disclose methods producing transgenic mice.

Transgenic animals of the present invention are produced by the application of procedures which result in an animal with a genome that has incorporated exogenous genetic material. The procedure involves obtaining the genetic material, or a portion thereof, which encodes either a *HKLP* coding sequence, a *HKLP* regulatory polynucleotide or a DNA sequence encoding a *HKLP* antisense polynucleotide such as described in the present specification.

A recombinant polynucleotide of the invention is inserted into an embryonic or ES stem cell line. The insertion is preferably made using electroporation, such as described by Thomas et al.(1987). The cells subjected to electroporation are screened (e.g. by selection via selectable markers, by PCR or by Southern blot analysis) to find positive cells which have integrated the exogenous recombinant polynucleotide into their genome, preferably via an homologous

recombination event. An illustrative positive-negative selection procedure that may be used according to the invention is described by Mansour et al.(1988).

Then, the positive cells are isolated, cloned and injected into 3.5 days old blastocysts from mice, such as described by Bradley (1987). The blastocysts are then inserted into a female host animal and allowed to grow to term.

Alternatively, the positive ES cells are brought into contact with embryos at the 2.5 days old 8-16 cell stage (morulae) such as described by Wood et al.(1993) or by Nagy et al.(1993), the ES cells being internalized to colonize extensively the blastocyst including the cells which will give rise to the germ line.

The offspring of the female host are tested to determine which animals are transgenic e.g. include the inserted exogenous DNA sequence and which are wild-type.

Thus, the present invention also concerns a transgenic animal containing a nucleic acid, a recombinant expression vector or a recombinant host cell according to the invention.

Recombinant Cell Lines Derived From The Transgenic Animals Of The Invention.

A further object of the invention consists of recombinant host cells obtained from a transgenic animal described herein. In one embodiment the invention encompasses cells derived from non-human host mammals and animals comprising a recombinant vector of the invention or a *HKLP* gene disrupted by homologous recombination with a knock out vector.

Recombinant cell lines may be established *in vitro* from cells obtained from any tissue of a transgenic animal according to the invention, for example by transfection of primary cell cultures with vectors expressing *onc*-genes such as SV40 large T antigen, as described by Chou (1989) and Shay et al.(1991).

Methods for screening substances interacting with a HKLP polypeptide

For the purpose of the present invention, a ligand means a molecule, such as a protein, a peptide, an antibody or any synthetic chemical compound capable of binding to the HKLP protein or one of its fragments or variants or to modulate the expression of the polynucleotide coding for HKLP or a fragment or variant thereof.

In the ligand screening method according to the present invention, a biological sample or a defined molecule to be tested as a putative ligand of the HKLP protein is brought into contact with the corresponding purified HKLP protein, for example the corresponding purified recombinant HKLP protein produced by a recombinant cell host as described hereinbefore, in order to form a complex between this protein and the putative ligand molecule to be tested.

As an illustrative example, to study the interaction of the HKLP protein, or a fragment comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, or 100 amino acids of SEQ ID No 4, wherein said contiguous span includes at least 1, 2, 3, 5 or 10 of the amino acid positions 1-478 of the SEQ ID No

4, with drugs or small molecules, such as molecules generated through combinatorial chemistry approaches, the microdialysis coupled to HPLC method described by Wang et al. (1997) or the affinity capillary electrophoresis method described by Bush et al. (1997), the disclosures of which are incorporated by reference, can be used.

- 5 In further methods, peptides, drugs, fatty acids, lipoproteins, or small molecules which interact with the HKLP protein, or a fragment comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, or 100 amino acids of SEQ ID No 4, wherein said contiguous span includes at least 1, 2, 3, 5 or 10 of the amino acid positions 1-478 of the SEQ ID No 4 may be identified using assays such as the
- 10 following. The molecule to be tested for binding is labeled with a detectable label, such as a fluorescent, radioactive, or enzymatic tag and placed in contact with immobilized HKLP protein, or a fragment thereof under conditions which permit specific binding to occur. After removal of non-specifically bound molecules, bound molecules are detected using appropriate means.

Another object of the present invention consists of methods and kits for the screening of

15 candidate substances that interact with HKLP polypeptide.

The present invention pertains to methods for screening substances of interest that interact with a HKLP protein or one fragment or variant thereof. By their capacity to bind covalently or non-covalently to a HKLP protein or to a fragment or variant thereof, these substances or molecules may be advantageously used both *in vitro* and *in vivo*.

- 20 *In vitro*, said interacting molecules may be used as detection means in order to identify the presence of a HKLP protein in a sample, preferably a biological sample.

A method for the screening of a candidate substance comprises the following steps :

- a) providing a polypeptide consisting of a HKLP protein or a fragment comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at
- 25 least 12, 15, 20, 25, 30, 40, 50, or 100 amino acids of SEQ ID No 4, wherein said contiguous span includes at least 1, 2, 3, 5 or 10 of the amino acid positions 1-478 of the SEQ ID No 4 or a variant thereof;
- b) obtaining a candidate substance;
- c) bringing into contact said polypeptide with said candidate substance;
- 30 d) detecting the complexes formed between said polypeptide and said candidate substance.

The invention further concerns a kit for the screening of a candidate substance interacting with the HKLP polypeptide, wherein said kit comprises :

- a) a HKLP protein having an amino acid sequence selected from the group consisting of the amino acid sequences of SEQ ID No 4 or a peptide fragment comprising a contiguous span of at
- 35 least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, or 100 amino acids of SEQ ID No 4, wherein said contiguous span includes at least 1, 2, 3, 5 or 10 of the amino acid positions 1-478 of the SEQ ID No 4 or a variant thereof ;

b) optionally means useful to detect the complex formed between the HKLP protein or a peptide fragment or a variant thereof and the candidate substance.

In a preferred embodiment of the kit described above, the detection means consist in monoclonal or polyclonal antibodies directed against the HKLP protein or a peptide fragment or a
5 variant thereof.

Various candidate substances or molecules can be assayed for interaction with a HKLP polypeptide. These substances or molecules include, without being limited to, natural or synthetic organic compounds or molecules of biological origin such as polypeptides. When the candidate substance or molecule consists of a polypeptide, this polypeptide may be the resulting expression
10 product of a phage clone belonging to a phage-based random peptide library, or alternatively the polypeptide may be the resulting expression product of a cDNA library cloned in a vector suitable for performing a two-hybrid screening assay.

The invention also pertains to kits useful for performing the hereinbefore described screening method. Preferably, such kits comprise a HKLP polypeptide or a fragment or a variant
15 thereof, and optionally means useful to detect the complex formed between the HKLP polypeptide or its fragment or variant and the candidate substance. In a preferred embodiment the detection means consist in monoclonal or polyclonal antibodies directed against the corresponding HKLP polypeptide or a fragment or a variant thereof.

A. Candidate ligands obtained from random peptide libraries

20 In a particular embodiment of the screening method, the putative ligand is the expression product of a DNA insert contained in a phage vector (Parmley and Smith, 1988). Specifically, random peptide phages libraries are used. The random DNA inserts encode for peptides of 8 to 20 amino acids in length (Oldenburg K.R. et al., 1992; Valadon P., et al., 1996; Lucas A.H., 1994; Westerink M.A.J., 1995; Felici F. et al., 1991). According to this particular embodiment, the
25 recombinant phages expressing a protein that binds to the immobilized HKLP protein is retained and the complex formed between the HKLP protein and the recombinant phage may be subsequently immunoprecipitated by a polyclonal or a monoclonal antibody directed against the HKLP protein.

Once the ligand library in recombinant phages has been constructed, the phage population is brought into contact with the immobilized HKLP protein. Then the preparation of complexes is
30 washed in order to remove the non-specifically bound recombinant phages. The phages that bind specifically to the HKLP protein are then eluted by a buffer (acid pH) or immunoprecipitated by the monoclonal antibody produced by the hybridoma anti-HKLP, and this phage population is subsequently amplified by an over-infection of bacteria (for example E. coli). The selection step may be repeated several times, preferably 2-4 times, in order to select the more specific recombinant
35 phage clones. The last step consists in characterizing the peptide produced by the selected recombinant phage clones either by expression in infected bacteria and isolation, expressing the

phage insert in another host-vector system, or sequencing the insert contained in the selected recombinant phages.

B. Candidate ligands obtained by competition experiments.

Alternatively, peptides, drugs or small molecules which bind to the HKLP protein, or a
5 fragment comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, or 100 amino acids of SEQ ID No 4, wherein said contiguous span includes at least 1, 2, 3, 5 or 10 of the amino acid positions 1-478 of the SEQ ID No 4, may be identified in competition experiments. In such assays, the HKLP protein, or a fragment thereof, is immobilized to a surface, such as a plastic plate. Increasing amounts of the
10 peptides, drugs or small molecules are placed in contact with the immobilized HKLP protein, or a fragment thereof, in the presence of a detectable labeled known HKLP protein ligand. For example, the HKLP ligand may be detectably labeled with a fluorescent, radioactive, or enzymatic tag. The ability of the test molecule to bind the HKLP protein, or a fragment thereof, is determined by measuring the amount of detectably labeled known ligand bound in the presence of the test
15 molecule. A decrease in the amount of known ligand bound to the HKLP protein, or a fragment thereof, when the test molecule is present indicated that the test molecule is able to bind to the HKLP protein, or a fragment thereof.

C. Candidate ligands obtained by affinity chromatography.

Proteins or other molecules interacting with the HKLP protein, or a fragment comprising a
20 contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, or 100 amino acids of SEQ ID No 4, wherein said contiguous span includes at least 1, 2, 3, 5 or 10 of the amino acid positions 1-478 of the SEQ ID No 4, can also be found using affinity columns which contain the HKLP protein, or a fragment thereof. The HKLP protein, or a fragment thereof, may be attached to the column using conventional techniques
25 including chemical coupling to a suitable column matrix such as agarose, Affi Gel®, or other matrices familiar to those of skill in art. In some embodiments of this method, the affinity column contains chimeric proteins in which the HKLP protein, or a fragment thereof, is fused to glutathion S transferase (GST). A mixture of cellular proteins or pool of expressed proteins as described above is applied to the affinity column. Proteins or other molecules interacting with the HKLP protein, or a
30 fragment thereof, attached to the column can then be isolated and analyzed on 2-D electrophoresis gel as described in Ramunsen et al. (1997), the disclosure of which is incorporated by reference. Alternatively, the proteins retained on the affinity column can be purified by electrophoresis based methods and sequenced. The same method can be used to isolate antibodies, to screen phage display products, or to screen phage display human antibodies.

D. Candidate ligands obtained by optical biosensor methods

Proteins interacting with the HKLP protein, or a fragment comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, or 100 amino acids of SEQ ID No 4, wherein said contiguous span includes at least 1, 2, 3, 5 or 10 of the amino acid positions 1-478 of the SEQ ID No 4, can also be screened by using an Optical Biosensor as described in Edwards and Leatherbarrow (1997) and also in Szabo et al. (1995), the disclosure of which is incorporated by reference. This technique permits the detection of interactions between molecules in real time, without the need of labeled molecules. This technique is based on the surface plasmon resonance (SPR) phenomenon. Briefly, the candidate ligand molecule to be tested is attached to a surface (such as a carboxymethyl dextran matrix). A light beam is directed towards the side of the surface that does not contain the sample to be tested and is reflected by said surface. The SPR phenomenon causes a decrease in the intensity of the reflected light with a specific association of angle and wavelength. The binding of candidate ligand molecules cause a change in the refraction index on the surface, which change is detected as a change in the SPR signal. For screening of candidate ligand molecules or substances that are able to interact with the HKLP protein, or a fragment thereof, the HKLP protein, or a fragment thereof, is immobilized onto a surface. This surface consists of one side of a cell through which flows the candidate molecule to be assayed. The binding of the candidate molecule on the HKLP protein, or a fragment thereof, is detected as a change of the SPR signal. The candidate molecules tested may be proteins, peptides, carbohydrates, lipids, or small molecules generated by combinatorial chemistry. This technique may also be performed by immobilizing eukaryotic or prokaryotic cells or lipid vesicles exhibiting an endogenous or a recombinantly expressed HKLP protein at their surface.

The main advantage of the method is that it allows the determination of the association rate between the HKLP protein and molecules interacting with the HKLP protein. It is thus possible to select specifically ligand molecules interacting with the HKLP protein, or a fragment thereof, through strong or conversely weak association constants.

E. Candidate ligands obtained through a two-hybrid screening assay.

The yeast two-hybrid system is designed to study protein-protein interactions *in vivo* (Fields and Song, 1989), and relies upon the fusion of a bait protein to the DNA binding domain of the yeast Gal4 protein. This technique is also described in the US Patent N° US 5,667,973 and the US Patent N° 5,283,173 (Fields et al.) the technical teachings of both patents being herein incorporated by reference.

The general procedure of library screening by the two-hybrid assay may be performed as described by Harper et al. (1993) or as described by Cho et al. (1998) or also Fromont-Racine et al. (1997).

The bait protein or polypeptide consists of a HKLP polypeptide or a fragment comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, or 100 amino acids of SEQ ID No 4, wherein said contiguous span includes at least 1, 2, 3, 5 or 10 of the amino acid positions 1-478 of the SEQ ID No 4 or a variant thereof.

More precisely, the nucleotide sequence encoding the HKLP polypeptide or a fragment or variant thereof is fused to a polynucleotide encoding the DNA binding domain of the GAL4 protein, the fused nucleotide sequence being inserted in a suitable expression vector, for example pAS2 or pM3.

Then, a human cDNA library is constructed in a specially designed vector, such that the human cDNA insert is fused to a nucleotide sequence in the vector that encodes the transcriptional domain of the GAL4 protein. Preferably, the vector used is the pACT vector. The polypeptides encoded by the nucleotide inserts of the human cDNA library are termed "pray" polypeptides.

A third vector contains a detectable marker gene, such as beta galactosidase gene or CAT gene that is placed under the control of a regulation sequence that is responsive to the binding of a complete Gal4 protein containing both the transcriptional activation domain and the DNA binding domain. For example, the vector pG5EC may be used.

Two different yeast strains are also used. As an illustrative but non limiting example the two different yeast strains may be the followings :

- Y190, the phenotype of which is (*MATa, Leu2-3, 112 ura3-12, trp1-901, his3-D200, ade2-101, gal4Dgal180D URA3 GAL-LacZ, LYS GAL-HIS3, cyh^r*);
- Y187, the phenotype of which is (*MATa gal4 gal80 his3 trp1-901 ade2-101 ura3-52 leu2-3, -112 URA3 GAL-lacZmet^r*), which is the opposite mating type of Y190.

Briefly, 20 µg of pAS2/HKLP and 20 µg of pACT-cDNA library are co-transformed into yeast strain Y190. The transformants are selected for growth on minimal media lacking histidine, leucine and tryptophan, but containing the histidine synthesis inhibitor 3-AT (50 mM). Positive colonies are screened for beta galactosidase by filter lift assay. The double positive colonies (*His⁺, beta-gal⁺*) are then grown on plates lacking histidine, leucine, but containing tryptophan and cycloheximide (10 mg/ml) to select for loss of pAS2/HKLP plasmids but retention of pACT-cDNA library plasmids. The resulting Y190 strains are mated with Y187 strains expressing HKLP or non-related control proteins; such as cyclophilin B, lamin, or SNF1, as *Gal4* fusions as described by Harper et al. (1993) and by Bram et al. (1993), and screened for beta galactosidase by filter lift assay. Yeast clones that are *beta gal-* after mating with the control *Gal4* fusions are considered false positives.

In another embodiment of the two-hybrid method according to the invention, interaction between the HKLP or a fragment or variant thereof with cellular proteins may be assessed using the Matchmaker Two Hybrid System 2 (Catalog No. K1604-1, Clontech). As described in the manual

accompanying the Matchmaker Two Hybrid System 2 (Catalog No. K1604-1, Clontech), the disclosure of which is incorporated herein by reference, nucleic acids encoding the HKLP protein or a portion thereof, are inserted into an expression vector such that they are in frame with DNA encoding the DNA binding domain of the yeast transcriptional activator GAL4. A desired cDNA, preferably human
5 cDNA, is inserted into a second expression vector such that they are in frame with DNA encoding the activation domain of GAL4. The two expression plasmids are transformed into yeast and the yeast are plated on selection medium which selects for expression of selectable markers on each of the expression vectors as well as GAL4 dependent expression of the HIS3 gene. Transformants capable of growing on medium lacking histidine are screened for GAL4 dependent lacZ expression. Those cells which are
10 positive in both the histidine selection and the lacZ assay contain interaction between HKLP and the protein or peptide encoded by the initially selected cDNA insert.

Methods For Screening Substances Modulating The Activity Of The HKLP protein

The invention also concerns a method for screening new agents, or candidate substances which modulate the activity of the HKLP protein or a fragment thereof. Preferably, the HKLP
15 protein or a fragment thereof is a polypeptide code comprising a contiguous span of at least 6 amino acids of SEQ ID No 4, wherein said contiguous span includes at least 1 of the amino acid positions 1-478 of the SEQ ID No 4. Preferably, the candidate substance is mixed with the HKLP protein and the activity of the HKLP protein is measured. Candidate substances include, without being limited to, natural or synthetic organic compounds or molecules of biological origin such as polypeptides.

20 Various assays for biological activity of motor proteins are known (Sato-Yoshitake et al, 1992 and Scholey, 1993). In vitro motility assays to characterize specific KLPs, for example, include microtubule gliding assays demonstrating translocation of microtubules, organelle movement assays to visualize the movement of a cargo of interest, and single molecule motility assays (Howard 1989, Block 1990) to characterize structural elements.

25 In short, microtubule gliding assays can be performed by applying a sample containing the HKLP protein to a glass surface without any treatment and incubating with microtubules reconstituted from polymerized tubulin. Microtubule translocation activity and the direction of movement can be determined as in Nangaku 1994 by observing the movement of axonemes on the glass surface. Organelle movement assays can be performed by applying a composition containing
30 the cargo of interest with a solution containing vesicles and the HKLP protein to a glass surface. Movement of the organelle can be observed, for example, by using a cargo-specific fluorescent probe to stain vesicles before incubation with the HKLP protein.

Methods For Inhibiting The Expression Of A HKLP Gen

Other therapeutic compositions according to the present invention comprise advantageously
35 an oligonucleotide fragment of the nucleic sequence of *HKLP* as an antisense tool or a triple helix

tool that inhibits the expression of the corresponding *HKLP* gene. A preferred fragment of the nucleic sequence of *HKLP* comprises an allele of at least one of the biallelic markers A1 to A32.

Antisense Approach

Preferred methods using antisense polynucleotide according to the present invention are the
5 procedures described by Sczakiel et al.(1995).

Preferably, the antisense tools are chosen among the polynucleotides (15-200 bp long) that are complementary to the 5' end of the *HKLP* mRNA. In another embodiment, a combination of different antisense polynucleotides complementary to different parts of the desired targeted gene are used.

10 Preferred antisense polynucleotides according to the present invention are complementary to a sequence of the mRNAs of *HKLP* that contains either the translation initiation codon ATG or a splicing donor or acceptor site.

The antisense nucleic acids should have a length and melting temperature sufficient to permit formation of an intracellular duplex having sufficient stability to inhibit the expression of the
15 *HKLP* mRNA in the duplex. Strategies for designing antisense nucleic acids suitable for use in gene therapy are disclosed in Green et al., (1986) and Izant and Weintraub, (1984), the disclosures of which are incorporated herein by reference.

In some strategies, antisense molecules are obtained by reversing the orientation of the *HKLP* coding region with respect to a promoter so as to transcribe the opposite strand from that
20 which is normally transcribed in the cell. The antisense molecules may be transcribed using in vitro transcription systems such as those which employ T7 or SP6 polymerase to generate the transcript. Another approach involves transcription of *HKLP* antisense nucleic acids in vivo by operably linking DNA containing the antisense sequence to a promoter in a suitable expression vector.

Alternatively, suitable antisense strategies are those described by Rossi et al.(1991), in the
25 International Applications Nos. WO 94/23026, WO 95/04141, WO 92/18522 and in the European Patent Application No. EP 0 572 287 A2

An alternative to the antisense technology that is used according to the present invention consists in using ribozymes that will bind to a target sequence via their complementary polynucleotide tail and that will cleave the corresponding RNA by hydrolyzing its target site
30 (namely "hammerhead ribozymes"). Briefly, the simplified cycle of a hammerhead ribozyme consists of (1) sequence specific binding to the target RNA via complementary antisense sequences; (2) site-specific hydrolysis of the cleavable motif of the target strand; and (3) release of cleavage products, which gives rise to another catalytic cycle. Indeed, the use of long-chain antisense polynucleotide (at least 30 bases long) or ribozymes with long antisense arms are advantageous. A
35 preferred delivery system for antisense ribozyme is achieved by covalently linking these antisense ribozymes to lipophilic groups or to use liposomes as a convenient vector. Preferred antisense

ribozymes according to the present invention are prepared as described by Sczakiel et al.(1995), the specific preparation procedures being referred to in said article being herein incorporated by reference.

Triple Helix Approach

5 The *HKLP* genomic DNA may also be used to inhibit the expression of the *HKLP* gene based on intracellular triple helix formation.

Triple helix oligonucleotides are used to inhibit transcription from a genome. They are particularly useful for studying alterations in cell activity when it is associated with a particular gene.

10 Similarly, a portion of the *HKLP* genomic DNA can be used to study the effect of inhibiting *HKLP* transcription within a cell. Traditionally, homopurine sequences were considered the most useful for triple helix strategies. However, homopyrimidine sequences can also inhibit gene expression. Such homopyrimidine oligonucleotides bind to the major groove at homopurine:homopyrimidine sequences. Thus, both types of sequences from the *HKLP* genomic
15 DNA are contemplated within the scope of this invention.

To carry out gene therapy strategies using the triple helix approach, the sequences of the *HKLP* genomic DNA are first scanned to identify 10-mer to 20-mer homopyrimidine or homopurine stretches which could be used in triple-helix based strategies for inhibiting *HKLP* expression. Following identification of candidate homopyrimidine or homopurine stretches, their efficiency in
20 inhibiting *HKLP* expression is assessed by introducing varying amounts of oligonucleotides containing the candidate sequences into tissue culture cells which express the *HKLP* gene.

The oligonucleotides can be introduced into the cells using a variety of methods known to those skilled in the art, including but not limited to calcium phosphate precipitation, DEAE-Dextran, electroporation, liposome-mediated transfection or native uptake.

25 Treated cells are monitored for altered cell function or reduced *HKLP* expression using techniques such as Northern blotting, RNase protection assays, or PCR based strategies to monitor the transcription levels of the *HKLP* gene in cells which have been treated with the oligonucleotide.

The oligonucleotides which are effective in inhibiting gene expression in tissue culture cells may then be introduced in vivo using the techniques described above in the antisense approach at a
30 dosage calculated based on the in vitro results, as described in antisense approach.

In some embodiments, the natural (beta) anomers of the oligonucleotide units can be replaced with alpha anomers to render the oligonucleotide more resistant to nucleases. Further, an intercalating agent such as ethidium bromide, or the like, can be attached to the 3' end of the alpha oligonucleotide to stabilize the triple helix. For information on the generation of oligonucleotides
35 suitable for triple helix formation see Griffin et al.(1989), which is hereby incorporated by this reference.

Computer-Related Embodiments

As used herein the term "nucleic acid codes of the invention" encompass the nucleotide sequences comprising, consisting essentially of, or consisting of any one of the following: a) a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 1, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 1: 1-39624, 39705-40589, 40666-43629, 43710-44203, 44311-45125, 45210-45440, 45622-45717, 45791-68580, 68675-70246, 70396-72421, 72601-73295, 73434-74648, 74898-83055, 83175-85192, 85279-85609, 85740-85906, 86070-88304, 88396-90585, 90705-91767, 91824-94380, 94490-96296, 96364-97184, 97270-101167, 101274-109465, 109581-110228, 110363-111819, 111882-113636, 113783-113945, 114186-117002, 117075-119676, and 119677-121162; b) a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 2 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 2: 1-1600, 1751-2138, 2332-2539, 2659-3829 and 8885-10884; c) a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 3 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 3: 391-1619 and 6988-10682; d) a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 1, 2 or 3, or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 nucleotide positions of any one of the following ranges of nucleotide positions of: (1) SEQ ID No 1: 1-1000, 1001-2000, 2001-3000, 3001-4000, 4001-5000, 5001-6000, 6001-7000, 7001-8000, 8001-9000, 9001-10000, 10001-11000, 11001-12000, 12001-13000, 13001-14000, 14001-15000, 15001-16000, 16001-17000, 17001-18000, 18001-19000, 19001-20000, 20001-21000, 21001-22000, 22001-23000, 23001-24000, 24001-25000, 25001-26000, 26001-27000, 27001-28000, 28001-29000, 29001-30000, 30001-31000, 31001-32000, 32001-33000, 33001-34000, 34001-35000, 35001-36000, 36001-37000, 37001-38000, 38001-39000, 39001-39624, 39705-40589, 40666-43629, 43710-44203, 44311-45125, 45210-45440, 45622-45717, 45791-68580, 68675-70246, 70396-72421, 72601-73295, 73434-74648, 74898-83055, 83175-85192, 85279-85609, 85740-85906, 86070-88304, 88396-90585, 90705-91767, 91824-94380, 94490-96296, 96364-97184, 97270-101167, 101274-109465, 109581-110228, 110363-111819, 111882-113636, 113783-113945, 114186-117002, 117075-119676, and 119677-121162; and (2) SEQ ID No 2: 1-1600, 1751-2138, 2332-2539, 2659-3829 and 8885-10884; e) a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 1 or the complements thereof, wherein said contiguous span comprises a G at position 7159 of SEQ ID No 1; f) a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 2 or the complements thereof, wherein said contiguous span comprises a C either at

- position 2551 or 4500 of SEQ ID No 2; g) a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 3 or the complements thereof, wherein said contiguous span comprises a nucleotide selected in the group consisting of a C at position 5487, and a C at position 6265 of SEQ ID No 3; and, j) a nucleotide sequence
- 5 complementary to any one of the preceding nucleotide sequences.

- The "nucleic acid codes of the invention" further encompass nucleotide sequences homologous to: a) a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 1, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 1: 1-39624, 39705-40589,
- 10 40666-43629, 43710-44203, 44311-45125, 45210-45440, 45622-45717, 45791-68580, 68675-70246, 70396-72421, 72601-73295, 73434-74648, 74898-83055, 83175-85192, 85279-85609, 85740-85906, 86070-88304, 88396-90585, 90705-91767, 91824-94380, 94490-96296, 96364-97184, 97270-101167, 101274-109465, 109581-110228, 110363-111819, 111882-113636, 113783-113945, 114186-117002, 117075-119676, and 119677-121162; b) a contiguous span of at least 12,
- 15 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 2 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 2: 1-1600, 1751-2138, 2332-2539, 2659-3829 and 8885-10884 c) a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 3 or the complements thereof, wherein said
- 20 contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 3: 391-1619 and 6988-10682; and d) sequences complementary to all of the preceding sequences. Homologous sequences refer to a sequence having at least 99%, 98%, 97%, 96%, 95%, 90%, 85%, 80%, or 75% homology to these contiguous spans. Homology may be determined using any method described herein, including BLAST2N with the default parameters or with any modified parameters.
- 25 Homologous sequences also may include RNA sequences in which uridines replace the thymines in the nucleic acid codes of the invention. It will be appreciated that the nucleic acid codes of the invention can be represented in the traditional single character format (See the inside back cover of Stryer, Lubert. *Biochemistry*, 3rd edition. W. H Freeman & Co., New York.) or in any other format or code which records the identity of the nucleotides in a sequence.

- 30 As used herein the term "polypeptide codes of the invention" encompass the polypeptide sequences comprising a contiguous span of at least 6, 8, 10, 12, 15, 20, 25, 30, 40, 50, or 100 amino acids of SEQ ID No 4, wherein said contiguous span includes at least 1, 2, 3, 5 or 10 of the amino acid positions 1-478 of the SEQ ID No 4. It will be appreciated that the polypeptide codes of the invention can be represented in the traditional single character format or three letter format (See the
- 35 inside back cover of Stryer, Lubert. *Biochemistry*, 3rd edition. W. H Freeman & Co., New York.) or in any other format or code which records the identity of the polypeptides in a sequence.

It will be appreciated by those skilled in the art that the nucleic acid codes of the invention and polypeptide codes of the invention can be stored, recorded, and manipulated on any medium which can be read and accessed by a computer. As used herein, the words "recorded" and "stored" refer to a process for storing information on a computer medium. A skilled artisan can readily adopt any of the presently known methods for recording information on a computer readable medium to generate manufactures comprising one or more of the nucleic acid codes of the invention, or one or more of the polypeptide codes of the invention. Another aspect of the present invention is a computer readable medium having recorded thereon at least 2, 5, 10, 15, 20, 25, 30, or 50 nucleic acid codes of the invention. Another aspect of the present invention is a computer readable medium having recorded thereon at least 2, 5, 10, 15, 20, 25, 30, or 50 polypeptide codes of the invention.

Computer readable media include magnetically readable media, optically readable media, electronically readable media and magnetic/optical media. For example, the computer readable media may be a hard disk, a floppy disk, a magnetic tape, CD-ROM, Digital Versatile Disk (DVD), Random Access Memory (RAM), or Read Only Memory (ROM) as well as other types of other media known to those skilled in the art.

Embodiments of the present invention include systems, particularly computer systems which store and manipulate the sequence information described herein. One example of a computer system 100 is illustrated in block diagram form in Figure 1. As used herein, "a computer system" refers to the hardware components, software components, and data storage components used to analyze the nucleotide sequences of the nucleic acid codes of the invention or the amino acid sequences of the polypeptide codes of the invention. In one embodiment, the computer system 100 is a Sun Enterprise 1000 server (Sun Microsystems, Palo Alto, CA). The computer system 100 preferably includes a processor for processing, accessing and manipulating the sequence data. The processor 105 can be any well-known type of central processing unit, such as the Pentium III from Intel Corporation, or similar processor from Sun, Motorola, Compaq or International Business Machines.

Preferably, the computer system 100 is a general purpose system that comprises the processor 105 and one or more internal data storage components 110 for storing data, and one or more data retrieving devices for retrieving the data stored on the data storage components. A skilled artisan can readily appreciate that any one of the currently available computer systems are suitable.

In one particular embodiment, the computer system 100 includes a processor 105 connected to a bus which is connected to a main memory 115 (preferably implemented as RAM) and one or more internal data storage devices 110, such as a hard drive and/or other computer readable media having data recorded thereon. In some embodiments, the computer system 100 further includes one or more data retrieving device 118 for reading the data stored on the internal data storage devices 110.

The data retrieving device 118 may represent, for example, a floppy disk drive, a compact disk drive, a magnetic tape drive, etc. In some embodiments, the internal data storage device 110 is a removable computer readable medium such as a floppy disk, a compact disk, a magnetic tape, etc.

containing control logic and/or data recorded thereon. The computer system 100 may advantageously include or be programmed by appropriate software for reading the control logic and/or the data from the data storage component once inserted in the data retrieving device.

The computer system 100 includes a display 120 which is used to display output to a computer user. It should also be noted that the computer system 100 can be linked to other computer systems 125a-c in a network or wide area network to provide centralized access to the computer system 100.

Software for accessing and processing the nucleotide sequences of the nucleic acid codes of the invention or the amino acid sequences of the polypeptide codes of the invention (such as search tools, compare tools, and modeling tools etc.) may reside in main memory 115 during execution.

10 In some embodiments, the computer system 100 may further comprise a sequence comparer for comparing the above-described nucleic acid codes of the invention or the polypeptide codes of the invention stored on a computer readable medium to reference nucleotide or polypeptide sequences stored on a computer readable medium. A "sequence comparer" refers to one or more programs which are implemented on the computer system 100 to compare a nucleotide or polypeptide sequence with
15 other nucleotide or polypeptide sequences and/or compounds including but not limited to peptides, peptidomimetics, and chemicals stored within the data storage means. For example, the sequence comparer may compare the nucleotide sequences of nucleic acid codes of the invention or the amino acid sequences of the polypeptide codes of the invention stored on a computer readable medium to reference sequences stored on a computer readable medium to identify homologies, motifs implicated in
20 biological function, or structural motifs. The various sequence comparer programs identified elsewhere in this patent specification are particularly contemplated for use in this aspect of the invention.

Figure 2 is a flow diagram illustrating one embodiment of a process 200 for comparing a new nucleotide or protein sequence with a database of sequences in order to determine the homology levels between the new sequence and the sequences in the database. The database of sequences can be a
25 private database stored within the computer system 100, or a public database such as GENBANK, PIR OR SWISSPROT that is available through the Internet.

The process 200 begins at a start state 201 and then moves to a state 202 wherein the new sequence to be compared is stored to a memory in a computer system 100. As discussed above, the memory could be any type of memory, including RAM or an internal storage device.

30 The process 200 then moves to a state 204 wherein a database of sequences is opened for analysis and comparison. The process 200 then moves to a state 206 wherein the first sequence stored in the database is read into a memory on the computer. A comparison is then performed at a state 210 to determine if the first sequence is the same as the second sequence. It is important to note that this step is not limited to performing an exact comparison between the new sequence and the first sequence
35 in the database. Well-known methods are known to those of skill in the art for comparing two nucleotide or protein sequences, even if they are not identical. For example, gaps can be introduced into one sequence in order to raise the homology level between the two tested sequences. The parameters

that control whether gaps or other features are introduced into a sequence during comparison are normally entered by the user of the computer system.

Once a comparison of the two sequences has been performed at the state 210, a determination is made at a decision state 210 whether the two sequences are the same. Of course, the term "same" is not limited to sequences that are absolutely identical. Sequences that are within the homology parameters entered by the user will be marked as "same" in the process 200.

If a determination is made that the two sequences are the same, the process 200 moves to a state 214 wherein the name of the sequence from the database is displayed to the user. This state notifies the user that the sequence with the displayed name fulfills the homology constraints that were entered. Once the name of the stored sequence is displayed to the user, the process 200 moves to a decision state 218 wherein a determination is made whether more sequences exist in the database. If no more sequences exist in the database, then the process 200 terminates at an end state 220. However, if more sequences do exist in the database, then the process 200 moves to a state 224 wherein a pointer is moved to the next sequence in the database so that it can be compared to the new sequence. In this manner, the new sequence is aligned and compared with every sequence in the database.

It should be noted that if a determination had been made at the decision state 212 that the sequences were not homologous, then the process 200 would move immediately to the decision state 218 in order to determine if any other sequences were available in the database for comparison.

Accordingly, one aspect of the present invention is a computer system comprising a processor, a data storage device having stored thereon a nucleic acid code of the invention or a polypeptide code of the invention, a data storage device having retrievably stored thereon reference nucleotide sequences or polypeptide sequences to be compared to the nucleic acid code of the invention or polypeptide code of the invention and a sequence comparer for conducting the comparison. The sequence comparer may indicate a homology level between the sequences compared or identify structural motifs in the nucleic acid code of the invention and polypeptide codes of the invention or it may identify structural motifs in sequences which are compared to these nucleic acid codes and polypeptide codes. In some embodiments, the data storage device may have stored thereon the sequences of at least 2, 5, 10, 15, 20, 25, 30, or 50 of the nucleic acid codes of the invention or polypeptide codes of the invention.

Another aspect of the present invention is a method for determining the level of homology between a nucleic acid code of the invention and a reference nucleotide sequence, comprising the steps of reading the nucleic acid code and the reference nucleotide sequence through the use of a computer program which determines homology levels and determining homology between the nucleic acid code and the reference nucleotide sequence with the computer program. The computer program may be any of a number of computer programs for determining homology levels, including those specifically enumerated herein, including BLAST2N with the default parameters or with any modified parameters. The method may be implemented using the computer systems described above. The

method may also be performed by reading 2, 5, 10, 15, 20, 25, 30, or 50 of the above described nucleic acid codes of the invention through the use of the computer program and determining homology between the nucleic acid codes and reference nucleotide sequences.

Figure 3 is a flow diagram illustrating one embodiment of a process 250 in a computer for determining whether two sequences are homologous. The process 250 begins at a start state 252 and then moves to a state 254 wherein a first sequence to be compared is stored to a memory. The second sequence to be compared is then stored to a memory at a state 256. The process 250 then moves to a state 260 wherein the first character in the first sequence is read and then to a state 262 wherein the first character of the second sequence is read. It should be understood that if the sequence is a nucleotide sequence, then the character would normally be either A, T, C, G or U. If the sequence is a protein sequence, then it should be in the single letter amino acid code so that the first and second sequences can be easily compared.

A determination is then made at a decision state 264 whether the two characters are the same. If they are the same, then the process 250 moves to a state 268 wherein the next characters in the first and second sequences are read. A determination is then made whether the next characters are the same. If they are, then the process 250 continues this loop until two characters are not the same. If a determination is made that the next two characters are not the same, the process 250 moves to a decision state 274 to determine whether there are any more characters either sequence to read.

If there aren't any more characters to read, then the process 250 moves to a state 276 wherein the level of homology between the first and second sequences is displayed to the user. The level of homology is determined by calculating the proportion of characters between the sequences that were the same out of the total number of sequences in the first sequence. Thus, if every character in a first 100 nucleotide sequence aligned with a every character in a second sequence, the homology level would be 100%.

Alternatively, the computer program may be a computer program which compares the nucleotide sequences of the nucleic acid codes of the present invention, to reference nucleotide sequences in order to determine whether the nucleic acid code of the invention differs from a reference nucleic acid sequence at one or more positions. Optionally such a program records the length and identity of inserted, deleted or substituted nucleotides with respect to the sequence of either the reference polynucleotide or the nucleic acid code of the invention. In one embodiment, the computer program may be a program which determines whether the nucleotide sequences of the nucleic acid codes of the invention contain one or more single nucleotide polymorphisms (SNP) with respect to a reference nucleotide sequence. These single nucleotide polymorphisms may each comprise a single base substitution, insertion, or deletion.

Another aspect of the present invention is a method for determining the level of homology between a polypeptide code of the invention and a reference polypeptide sequence, comprising the

steps of reading the polypeptide code of the invention and the reference polypeptide sequence through use of a computer program which determines homology levels and determining homology between the polypeptide code and the reference polypeptide sequence using the computer program.

Accordingly, another aspect of the present invention is a method for determining whether a
5 nucleic acid code of the invention differs at one or more nucleotides from a reference nucleotide sequence comprising the steps of reading the nucleic acid code and the reference nucleotide sequence through use of a computer program which identifies differences between nucleic acid sequences and identifying differences between the nucleic acid code and the reference nucleotide sequence with the computer program. In some embodiments, the computer program is a program which identifies single
10 nucleotide polymorphisms. The method may be implemented by the computer systems described above and the method illustrated in Figure 3. The method may also be performed by reading at least 2, 5, 10, 15, 20, 25, 30, or 50 of the nucleic acid codes of the invention and the reference nucleotide sequences through the use of the computer program and identifying differences between the nucleic acid codes and the reference nucleotide sequences with the computer program.

15 In other embodiments the computer based system may further comprise an identifier for identifying features within the nucleotide sequences of the nucleic acid codes of the invention or the amino acid sequences of the polypeptide codes of the invention.

An "identifier" refers to one or more programs which identifies certain features within the above-described nucleotide sequences of the nucleic acid codes of the invention or the amino acid
20 sequences of the polypeptide codes of the invention. In one embodiment, the identifier may comprise a program which identifies an open reading frame in the cDNAs codes of the invention.

Figure 4 is a flow diagram illustrating one embodiment of an identifier process 300 for detecting the presence of a feature in a sequence. The process 300 begins at a start state 302 and then moves to a state 304 wherein a first sequence that is to be checked for features is stored to a
25 memory 115 in the computer system 100. The process 300 then moves to a state 306 wherein a database of sequence features is opened. Such a database would include a list of each feature's attributes along with the name of the feature. For example, a feature name could be "Initiation Codon" and the attribute would be "ATG". Another example would be the feature name "TAATAA Box" and the feature attribute would be "TAATAA". An example of such a database is produced by
30 the University of Wisconsin Genetics Computer Group (www.gcg.com).

Once the database of features is opened at the state 306, the process 300 moves to a state 308 wherein the first feature is read from the database. A comparison of the attribute of the first feature with the first sequence is then made at a state 310. A determination is then made at a decision state 316 whether the attribute of the feature was found in the first sequence. If the attribute
35 was found, then the process 300 moves to a state 318 wherein the name of the found feature is displayed to the user.

The process 300 then moves to a decision state 320 wherein a determination is made whether move features exist in the database. If no more features do exist, then the process 300 terminates at an end state 324. However, if more features do exist in the database, then the process 300 reads the next sequence feature at a state 326 and loops back to the state 310 wherein the
5 attribute of the next feature is compared against the first sequence.

It should be noted, that if the feature attribute is not found in the first sequence at the decision state 316, the process 300 moves directly to the decision state 320 in order to determine if any more features exist in the database.

In another embodiment, the identifier may comprise a molecular modeling program which
10 determines the 3-dimensional structure of the polypeptides codes of the invention. In some embodiments, the molecular modeling program identifies target sequences that are most compatible with profiles representing the structural environments of the residues in known three-dimensional protein structures. (See, e.g., Eisenberg et al., U.S. Patent No. 5,436,850 issued July 25, 1995). In another technique, the known three-dimensional structures of proteins in a given family are
15 superimposed to define the structurally conserved regions in that family. This protein modeling technique also uses the known three-dimensional structure of a homologous protein to approximate the structure of the polypeptide codes of the invention. (See e.g., Srinivasan, et al., U.S. Patent No. 5,557,535 issued September 17, 1996). Conventional homology modeling techniques have been used routinely to build models of proteases and antibodies. (Sowdhamini et al., 1997). Comparative
20 approaches can also be used to develop three-dimensional protein models when the protein of interest has poor sequence identity to template proteins. In some cases, proteins fold into similar three-dimensional structures despite having very weak sequence identities. For example, the three-dimensional structures of a number of helical cytokines fold in similar three-dimensional topology in spite of weak sequence homology.

25 The recent development of threading methods now enables the identification of likely folding patterns in a number of situations where the structural relatedness between target and template(s) is not detectable at the sequence level. Hybrid methods, in which fold recognition is performed using Multiple Sequence Threading (MST), structural equivalencies are deduced from the threading output using a distance geometry program DRAGON to construct a low resolution model,
30 and a full-atom representation is constructed using a molecular modeling package such as QUANTA.

According to this 3-step approach, candidate templates are first identified by using the novel fold recognition algorithm MST, which is capable of performing simultaneous threading of multiple aligned sequences onto one or more 3-D structures. In a second step, the structural equivalencies
35 obtained from the MST output are converted into interresidue distance restraints and fed into the distance geometry program DRAGON, together with auxiliary information obtained from secondary structure predictions. The program combines the restraints in an unbiased manner and rapidly

generates a large number of low resolution model confirmations. In a third step, these low resolution model confirmations are converted into full-atom models and subjected to energy minimization using the molecular modeling package QUANTA. (See e.g., Aszódi et al., 1997).

The results of the molecular modeling analysis may then be used in rational drug design techniques to identify agents which modulate the activity of the polypeptide codes of the invention.

Accordingly, another aspect of the present invention is a method of identifying a feature within the nucleic acid codes of the invention or the polypeptide codes of the invention comprising reading the nucleic acid code(s) or the polypeptide code(s) through the use of a computer program which identifies features therein and identifying features within the nucleic acid code(s) or polypeptide code(s) with the computer program. In one embodiment, computer program comprises a computer program which identifies open reading frames. In a further embodiment, the computer program identifies structural motifs in a polypeptide sequence. In another embodiment, the computer program comprises a molecular modeling program. The method may be performed by reading a single sequence or at least 2, 5, 10, 15, 20, 25, 30, or 50 of the nucleic acid codes of the invention or the polypeptide codes of the invention through the use of the computer program and identifying features within the nucleic acid codes or polypeptide codes with the computer program.

The nucleic acid codes of the invention or the polypeptide codes of the invention may be stored and manipulated in a variety of data processor programs in a variety of formats. For example, they may be stored as text in a word processing file, such as MicrosoftWORD or WORDPERFECT or as an ASCII file in a variety of database programs familiar to those of skill in the art, such as DB2, SYBASE, or ORACLE. In addition, many computer programs and databases may be used as sequence comparers, identifiers, or sources of reference nucleotide or polypeptide sequences to be compared to the nucleic acid codes of the invention or the polypeptide codes of the invention. The following list is intended not to limit the invention but to provide guidance to programs and databases which are useful with the nucleic acid codes of the invention or the polypeptide codes of the invention. The programs and databases which may be used include, but are not limited to: MacPattern (EMBL), DiscoveryBase (Molecular Applications Group), GeneMine (Molecular Applications Group), Look (Molecular Applications Group), MacLook (Molecular Applications Group), BLAST and BLAST2 (NCBI), BLASTN and BLASTX (Altschul et al, 1990), FASTA (Pearson and Lipman, 1988), FASTDB (Brutlag et al., 1990), Catalyst (Molecular Simulations Inc.), Catalyst/SHAPE (Molecular Simulations Inc.), Cerius².DBAccess (Molecular Simulations Inc.), HypoGen (Molecular Simulations Inc.), Insight II, (Molecular Simulations Inc.), Discover (Molecular Simulations Inc.), CHARMm (Molecular Simulations Inc.), Felix (Molecular Simulations Inc.), DelPhi, (Molecular Simulations Inc.), QuanteMM, (Molecular Simulations Inc.), Homology (Molecular Simulations Inc.), Modeler (Molecular Simulations Inc.), ISIS (Molecular Simulations Inc.), Quanta/Protein Design (Molecular Simulations Inc.), WebLab (Molecular Simulations Inc.), WebLab Diversity Explorer (Molecular Simulations Inc.), Gene Explorer (Molecular Simulations Inc.), SeqFold (Molecular Simulations Inc.),

the EMBL/Swissprotein database, the MDL Available Chemicals Directory database, the MDL Drug Data Report data base, the Comprehensive Medicinal Chemistry database, Derwent's World Drug Index database, the BioByteMasterFile database, the Genbank database, and the Genseqn database.

Many other programs and data bases would be apparent to one of skill in the art given the present
5 disclosure.

Motifs which may be detected using the above programs include sequences encoding leucine zippers, helix-turn-helix motifs, glycosylation sites, ubiquitination sites, alpha helices, and beta sheets, signal sequences encoding signal peptides which direct the secretion of the encoded proteins, sequences implicated in transcription regulation such as homeoboxes, acidic stretches,
10 enzymatic active sites, substrate binding sites, and enzymatic cleavage sites.

Throughout this application, various publications, patents and published patent applications are cited. The disclosures of these publications, patents and published patent specification referenced in this application are hereby incorporated by reference into the present disclosure to
15 more fully describe the state of the art to which this invention pertains.

EXAMPLES

Example 1

Identification Of Biallelic Markers - DNA Extraction

Donors were unrelated and healthy. They presented a sufficient diversity for being
20 representative of a French heterogeneous population. The DNA from 100 individuals was extracted and tested for the detection of the biallelic markers.

30 ml of peripheral venous blood were taken from each donor in the presence of EDTA. Cells (pellet) were collected after centrifugation for 10 minutes at 2000 rpm. Red cells were lysed by a lysis solution (50 ml final volume: 10 mM Tris pH7.6; 5 mM MgCl₂; 10 mM NaCl). The
25 solution was centrifuged (10 minutes, 2000 rpm) as many times as necessary to eliminate the residual red cells present in the supernatant, after resuspension of the pellet in the lysis solution.

The pellet of white cells was lysed overnight at 42°C with 3.7 ml of lysis solution composed of:

- 3 ml TE 10-2 (Tris-HCl 10 mM, EDTA 2 mM) / NaCl 0.4 M
- 30 - 200 µl SDS 10%
- 500 µl K-proteinase (2 mg K-proteinase in TE 10-2 / NaCl 0.4 M).

For the extraction of proteins, 1 ml saturated NaCl (6M) (1/3.5 v/v) was added. After vigorous agitation, the solution was centrifuged for 20 minutes at 10000 rpm.

For the precipitation of DNA, 2 to 3 volumes of 100% ethanol were added to the previous supernatant, and the solution was centrifuged for 30 minutes at 2000 rpm. The DNA solution was rinsed three times with 70% ethanol to eliminate salts, and centrifuged for 20 minutes at 2000 rpm. The pellet was dried at 37°C, and resuspended in 1 ml TE 10-1 or 1 ml water. The DNA

5 concentration was evaluated by measuring the OD at 260 nm (1 unit OD = 50 µg/ml DNA).

To determine the presence of proteins in the DNA solution, the OD 260 / OD 280 ratio was determined. Only DNA preparations having a OD 260 / OD 280 ratio between 1.8 and 2 were used in the subsequent examples described below.

The pool was constituted by mixing equivalent quantities of DNA from each individual.

10

Example 2

Identification Of Biallelic Markers: Amplification Of Genomic DNA By PCR

The amplification of specific genomic sequences of the DNA samples of example 1 was carried out on the pool of DNA obtained previously. In addition, 50 individual samples were similarly amplified.

15 PCR assays were performed using the following protocol:

	Final volume	25 µl
	DNA	2 ng/µl
	MgCl ₂	2 mM
	dNTP (each)	200 µM
20	primer (each)	2.9 ng/µl
	Ampli Taq Gold DNA polymerase	0.05 unit/µl
	PCR buffer (10x = 0.1 M TrisHCl pH8.3 0.5M KCl)	1x

Each pair of first primers was designed using the sequence information of the *HKLP* gene
25 disclosed herein and the OSP software (Hillier & Green, 1991). This first pair of primers was about 20 nucleotides in length and had the sequences disclosed in Table 1 in the columns labeled PU and RP.

Preferably, the primers contained a common oligonucleotide tail upstream of the specific bases targeted for amplification which was useful for sequencing.

30 Primers PU contain the following additional PU 5' sequence:

TGTAAAACGACGGCCAGT; primers RP contain the following RP 5' sequence:

CAGGAAACAGCTATGACC. The primer containing the additional PU 5' sequence is listed in SEQ ID No 9. The primer containing the additional RP 5' sequence is listed in SEQ ID No 10.

The synthesis of these primers was performed following the phosphoramidite method, on a
35 GENSET UFPS 24.1 synthesizer.

DNA amplification was performed on a Genius II thermocycler. After heating at 95°C for 10 min, 40 cycles were performed. Each cycle comprised: 30 sec at 95°C, 54°C for 1 min, and 30 sec at 72°C. For final elongation, 10 min at 72°C ended the amplification. The quantities of the amplification products obtained were determined on 96-well microtiter plates, using a fluorometer 5 and Picogreen as intercalant agent (Molecular Probes).

Table 1

<u>Amplicon</u>	Position range of the amplicon in SEQ ID 1		PU Primer name	Position range of amplification primer in SEQ ID No 1		RP Primer name	Complementary position range of amplification primer in SEQ ID No 1	
12-809	7041	7565	B1	7041	7060	C1	7545	7565
12-805	16004	16483	B2	16004	16024	C2	16464	16483
12-790	26232	26705	B3	26232	26252	C3	26685	26705
12-791	30902	31301	B4	30902	30922	C4	31282	31301
12-803	33476	33932	B5	33476	33496	C5	33912	33932
99-33040	33934	34383	B6	33934	33953	C6	34364	34383
12-810	34918	35369	B7	34918	34938	C7	35350	35369
12-787	45465	45993	B8	45465	45485	C8	45974	45993
12-793	54361	54879	B9	54361	54381	C9	54861	54879
12-792	56365	56813	B10	56365	56385	C10	56793	56813
99-41009	60111	60580	B11	60111	60130	C11	60562	60580
12-593	78815	79349	B12	78815	78835	C12	79332	79349
12-589	79706	80167	B13	79706	79726	C13	80149	80167
12-785	87264	87773	B14	87264	87284	C14	87753	87773
12-588	101161	101704	B15	101161	101178	C15	101686	101704
12-603	104131	104578	B16	104131	104151	C16	104558	104578
12-586	117017	117501	B17	117017	117036	C17	117481	117501
<u>Amplicon</u>	Position range of the amplicon in SEQ ID 2		PU Primer name	Position range of amplification primer in SEQ ID No 2		RP Primer name	Complementary position range of amplification primer in SEQ ID No 2	
12-602	2203	2620	B18	2203	2221	C18	2600	2620
12-587	4479	4878	B19	4479	4499	C19	4858	4878
12-596	5996	6443	B20	5996	6015	C20	6423	6443
12-808	10079	10543	B21	10079	10098	C21	10523	10543
Amplicon	Position range of the amplicon		Primer name	Position range of amplification primer		Primer name	Complementary position range of amplification primer	
10-265	1	357	B22	1	18	C22	338	357
	SEQ ID No 5			SEQ ID No 5			SEQ ID No 5	
10-266	1	420	B23	1	20	C23	401	420
	SEQ ID No 6			SEQ ID No 6			SEQ ID No 6	
12-592	1	465	B24	1	19	C24	448	465
	SEQ ID No 7			SEQ ID No 7			SEQ ID No 7	
12-783	1	449	B25	1	19	C25	429	449
	SEQ ID No 8			SEQ ID No 8			SEQ ID No 8	

Example 3**Identification Of Biallelic Markers - Sequencing Of Amplified Genomic DNA And Identification Of Polymorphisms**

The sequencing of the amplified DNA obtained in example 2 was carried out on ABI 377
 5 sequencers. The sequences of the amplification products were determined using automated dideoxy terminator sequencing reactions with a dye terminator cycle sequencing protocol. The products of the sequencing reactions were run on sequencing gels and the sequences were determined using gel image analysis (ABI Prism DNA Sequencing Analysis software (2.1.2 version)).

The sequence data were further evaluated to detect the presence of biallelic markers within
 10 the amplified fragments. The polymorphism search was based on the presence of superimposed peaks in the electrophoresis pattern resulting from different bases occurring at the same position as described previously.

In the 25 fragments of amplification, 32 biallelic markers were detected. The localization of these biallelic markers are as shown in Table 2.

15

Example 4**Validation Of The Polymorphisms Through Microsequencing**

The biallelic markers identified in example 3 were further confirmed and their respective frequencies were determined through microsequencing. Microsequencing was carried out for each individual DNA sample described in Example 1.

20 Amplification from genomic DNA of individuals was performed by PCR as described above for the detection of the biallelic markers with the same set of PCR primers (Table 1).

The preferred primers used in microsequencing were about 19 nucleotides in length and hybridized just upstream of the considered polymorphic base. According to the invention, the primers used in microsequencing are detailed in Table 4.

25 The microsequencing reaction was performed as follows :

After purification of the amplification products, the microsequencing reaction mixture was prepared by adding, in a 20µl final volume: 10 pmol microsequencing oligonucleotide, 1 U Thermosequenase (Amersham E79000G), 1.25 µl Thermosequenase buffer (260 mM Tris HCl pH 9.5, 65 mM MgCl₂), and the two appropriate fluorescent ddNTPs (Perkin Elmer, Dye Terminator Set 30 401095) complementary to the nucleotides at the polymorphic site of each biallelic marker tested, following the manufacturer's recommendations. After 4 minutes at 94°C, 20 PCR cycles of 15 sec at 55°C, 5 sec at 72°C, and 10 sec at 94°C were carried out in a Tetrad PTC-225 thermocycler (MJ Research). The unincorporated dye terminators were then removed by ethanol precipitation. Samples were finally resuspended in formamide-EDTA loading buffer and heated for 2 min at 95°C

before being loaded on a polyacrylamide sequencing gel. The data were collected by an ABI PRISM 377 DNA sequencer and processed using the GENESCAN software (Perkin Elmer).

Following gel analysis, data were automatically processed with software that allows the determination of the alleles of biallelic markers present in each amplified fragment.

- 5 The software evaluates such factors as whether the intensities of the signals resulting from the above microsequencing procedures are weak, normal, or saturated, or whether the signals are ambiguous. In addition, the software identifies significant peaks (according to shape and height criteria). Among the significant peaks, peaks corresponding to the targeted site are identified based on their position. When two significant peaks are detected for the same position, each sample is
- 10 categorized classification as homozygous or heterozygous type based on the height ratio.

Table 2

Amplicon	BM	Marker Name	Localizati n in <i>HKLP</i> gene	P lymor- phism		BM p siti n in SEQ ID	
				all1	all2	No 1	No 3
12-809	A1	12-809-119	Exon 3	G	C	7159	471
12-805	A2	12-805-115	Intron 5	A	G	16369	
12-790	A3	12-790-396	Intron 11	A	G	26310	
12-791	A4	12-791-211	Intron 14	A	G	31112	
12-803	A5	12-803-125	Intron 14	A	T	33808	
99-33040	A6	99-33040-321	Intron 14	C	T	34255	
12-810	A7	12-810-77	Intron 14	A	G	35293	
12-787	A8	12-787-103	Intron 21	A	G	45892	
12-793	A9	12-793-383	Intron 21	G	T	54497	
12-792	A10	12-792-233	Intron 21	A	G	56582	
99-41009	A11	99-41009-244	Intron 21	A	G	60336	
99-41009	A12	99-41009-111	Intron 21	C	T	60469	
12-593	A13	12-593-287	Intron 26	-	TAAAT	79063	
12-593	A14	12-593-174	Intron 26	C	T	79176	
12-589	A15	12-589-152	Intron 26	G	T	80016	
12-785	A16	12-785-200	Intron 30	C	T	87463	
12-785	A17	12-785-393	Intron 30	A	G	87656	
12-588	A18	12-588-103	Intron 36	A	G	101602	
12-603	A19	12-603-191	Intron 37	C	T	104391	
12-586	A20	12-586-414	Intron 43	A	G	117430	
12-586	A21	12-586-443	Intron 43	-	C	117459	
Amplicon	BM	Marker Name	Localization in <i>HKLP</i> gene	Polymor- phism		BM position in SEQ ID	
				all1	all2	No 2	No 3
12-602	A22	12-602-196	Intron 46	C	T	2397	
12-602	A23	12-602-350	Exon 47	A	C	2551	5487
12-587	A24	12-587-379	Exon 48	A	C	4500	6265
12-596	A25	12-596-124	Exon 48	A	G	6119	7887
12-808	A26	12-808-52	3' regulatory	A	G	10130	
12-808	A27	12-808-75	3' regulatory	G	C	10153	
Amplicon	BM	Marker Name	Localization	Polymor- phism		BM position In SEQ ID No	
				all1	all2		
10-265	A28	10-265-178	intergenic	A	G	178	5
10-266	A29	10-266-203	intergenic	C	T	203	6
12-592	A30	12-592-118	intergenic	A	T	118	7
12-783	A31	12-783-421	intergenic	C	T	420	8
12-783	A32	12-783-73	intergenic	G	C	72	8

BM refers to "biallelic marker". All1 and all2 refer respectively to allele 1 and allele 2 of the biallelic marker.

Table 3

BM	Marker Name	Position range f probes in SEQ ID No 1		Probes
A1	12-809-119	7147	7171	P1
A2	12-805-115	16357	16381	P2
A3	12-790-396	26298	26322	P3
A4	12-791-211	31100	31124	P4
A5	12-803-125	33796	33820	P5
A6	99-33040-321	34243	34267	P6
A7	12-810-77	35281	35305	P7
A8	12-787-103	45880	45904	P8
A9	12-793-383	54485	54509	P9
A10	12-792-233	56570	56594	P10
A11	99-41009-244	60324	60348	P11
A12	99-41009-111	60457	60481	P12
A14	12-593-174	79164	79188	P13
A15	12-589-152	80004	80028	P14
A16	12-785-200	87451	87475	P15
A17	12-785-393	87644	87668	P16
A18	12-588-103	101590	101614	P17
A19	12-603-191	104379	104403	P18
A20	12-586-414	117418	117442	P19
BM	Marker Name	Position range of probes in SEQ ID No 2		Probes
A22	12-602-196	2385	2409	P20
A23	12-602-350	2539	2563	P21
A24	12-587-379	4488	4512	P22
A25	12-596-124	6107	6131	P23
A26	12-808-52	10118	10142	P24
A27	12-808-75	10141	10165	P25
BM	Marker Name	Position range of probes		Probes
A28	10-265-178	166	190	P26
		In SEQ ID No 5		
A29	10-266-203	191	215	P27
		In SEQ ID No 6		
A30	12-592-118	106	130	P28
		In SEQ ID No 7		
A31	12-783-421	408	432	P29
		In SEQ ID No 8		
A32	12-783-73	60	84	P30
		In SEQ ID No 8		

Table 4

Marker Name	Biallelic Marker	Mis. 1	Position range of microsequencing primer in SEQ ID No 1		Mis. 2	Complementary position range of microsequencing primer in SEQ ID No 1	
12-809-119	A1	D1	7140	7158	E1	7160	7178
12-805-115	A2	D2	16350	16368	E2	16370	16388
12-790-396	A3	D3	26291	26309	E3	26311	26329
12-791-211	A4	D4	31093	31111	E4	31113	31131
12-803-125	A5	D5	33789	33807	E5	33809	33827
99-33040-321	A6	D6	34236	34254	E6	34256	34274
12-810-77	A7	D7	35274	35292	E7	35294	35312
12-787-103	A8	D8	45873	45891	E8	45893	45911
12-793-383	A9	D9	54478	54496	E9	54498	54516
12-792-233	A10	D10	56563	56581	E10	56583	56601
99-41009-244	A11	D11	60317	60335	E11	60337	60355
99-41009-111	A12	D12	60450	60468	E12	60470	60488
12-593-174	A14	D13	79157	79175	E13	79177	79195
12-589-152	A15	D14	79997	80015	E14	80017	80035
12-785-200	A16	D15	87444	87462	E15	87464	87482
12-785-393	A17	D16	87637	87655	E16	87657	87675
12-588-103	A18	D17	101583	101601	E17	101603	101621
12-603-191	A19	D18	104372	104390	E18	104392	104410
12-586-414	A20	D19	117411	117429	E19	117431	117449
Marker Name	Biallelic Marker	Mis. 1	Position range of microsequencing primer in SEQ ID No 2		Mis. 2	Complementary position range of microsequencing primer in SEQ ID No 2	
12-602-196	A22	D20	2378	2396	E20	2398	2416
12-602-350	A23	D21	2532	2550	E21	2552	2570
12-587-379	A24	D22	4481	4499	E22	4501	4519
12-596-124	A25	D23	6100	6118	E23	6120	6138
12-808-52	A26	D24	10111	10129	E24	10131	10149
12-808-75	A27	D25	10134	10152	E25	10154	10172
Marker Name	Biallelic Marker	Mis. 1	Position range of microsequencing primer		Mis. 2	Complementary position range of microsequencing primer	
10-265-178	A28	D26	159	177	E26	179	197
			In SEQ ID No 5			In SEQ ID No 5	
10-266-203	A29	D27	184	202	E27	204	222
			In SEQ ID No 6			In SEQ ID No 6	
12-592-118	A30	D28	99	117	E28	119	137
			In SEQ ID No 7			In SEQ ID No 7	
12-783-421	A31	D29	401	419	E29	421	439
			In SEQ ID No 8			In SEQ ID No 8	
12-783-73	A32	D30	53	71	E30	73	91
			In SEQ ID No 8			In SEQ ID No 8	

Mis 1 and Mis 2 respectively refer to microsequencing primers which hybridized with the non-coding strand of the *HKLP* gene or with the coding strand of the *HKLP* gene.

Example 5**Preparation of Antibody Compositions to the HKLP protein**

Substantially pure protein or polypeptide is isolated from transfected or transformed cells containing an expression vector encoding the HKLP protein or a portion thereof. The concentration of protein in the final preparation is adjusted, for example, by concentration on an Amicon filter device, to the level of a few micrograms/ml. Monoclonal or polyclonal antibody to the protein can then be prepared as follows:

A. Monoclonal Antibody Production by Hybridoma Fusion

Monoclonal antibody to epitopes in the HKLP protein or a portion thereof can be prepared from murine hybridomas according to the classical method of Kohler, G. and Milstein, C., (1975) or derivative methods thereof. Also see Harlow, E., and D. Lane. 1988.

Briefly, a mouse is repetitively inoculated with a few micrograms of the HKLP protein or a portion thereof over a period of a few weeks. The mouse is then sacrificed, and the antibody producing cells of the spleen isolated. The spleen cells are fused by means of polyethylene glycol with mouse myeloma cells, and the excess unfused cells destroyed by growth of the system on selective media comprising aminopterin (HAT media). The successfully fused cells are diluted and aliquots of the dilution placed in wells of a microtiter plate where growth of the culture is continued. Antibody-producing clones are identified by detection of antibody in the supernatant fluid of the wells by immunoassay procedures, such as ELISA, as originally described by Engvall, (1980), and derivative methods thereof. Selected positive clones can be expanded and their monoclonal antibody product harvested for use. Detailed procedures for monoclonal antibody production are described in Davis, L. et al. Basic Methods in Molecular Biology Elsevier, New York. Section 21-2.

B. Polyclonal Antibody Production by Immunization

Polyclonal antiserum containing antibodies to heterogeneous epitopes in the HKLP protein or a portion thereof can be prepared by immunizing suitable non-human animal with the HKLP protein or a portion thereof, which can be unmodified or modified to enhance immunogenicity. A suitable non-human animal is preferably a non-human mammal is selected, usually a mouse, rat, rabbit, goat, or horse. Alternatively, a crude preparation which has been enriched for HKLP concentration can be used to generate antibodies. Such proteins, fragments or preparations are introduced into the non-human mammal in the presence of an appropriate adjuvant (e.g. aluminum hydroxide, RIBI, etc.) which is known in the art. In addition the protein, fragment or preparation can be pretreated with an agent which will increase antigenicity, such agents are known in the art and include, for example, methylated bovine serum albumin (mBSA), bovine serum albumin (BSA), Hepatitis B surface antigen, and keyhole limpet hemocyanin (KLH). Serum from the immunized animal is collected, treated and tested according to known procedures. If the serum contains polyclonal antibodies to undesired epitopes, the polyclonal antibodies can be purified by immunoaffinity chromatography.

Effective polyclonal antibody production is affected by many factors related both to the antigen and the host species. Also, host animals vary in response to site of inoculations and dose, with both inadequate or excessive doses of antigen resulting in low titer antisera. Small doses (ng level) of antigen administered at multiple intradermal sites appears to be most reliable. Techniques
 5 for producing and processing polyclonal antisera are known in the art, see for example, Mayer and Walker (1987). An effective immunization protocol for rabbits can be found in Vaitukaitis, J. et al. (1971).

Booster injections can be given at regular intervals, and antiserum harvested when antibody titer thereof, as determined semi-quantitatively, for example, by double immunodiffusion in agar against
 10 known concentrations of the antigen, begins to fall. See, for example, Ouchterlony, O. et al., (1973). Plateau concentration of antibody is usually in the range of 0.1 to 0.2 mg/ml of serum (about 12 μ M). Affinity of the antisera for the antigen is determined by preparing competitive binding curves, as described, for example, by Fisher, D., (1980).

Antibody preparations prepared according to either the monoclonal or the polyclonal protocol
 15 are useful in quantitative immunoassays which determine concentrations of antigen-bearing substances in biological samples; they are also used semi-quantitatively or qualitatively to identify the presence of antigen in a biological sample. The antibodies may also be used in therapeutic compositions for killing cells expressing the protein or reducing the levels of the protein in the body.

20 While the preferred embodiment of the invention has been illustrated and described, it will be appreciated that various changes can be made therein by the one skilled in the art without departing from the spirit and scope of the invention.

FREE TEXT OF THE SEQUENCE LISTING

The following free text appears in the accompanying Sequence Listing :
 25 3'regulatory region
 polymorphic base
 or
 complement
 probe
 30 deletion of
 insertion of
 sequencing oligonucleotide Primer
 Artificial Sequence

REFERENCES

- The following references cited herein are incorporated herein by reference in their entireties
- Abbondanzo SJ et al., 1993, *Methods in Enzymology*, Academic Press, New York, pp. 803-823 / Ajioka R.S. et al., *Am. J. Hum. Genet.*, 60:1439-1447, 1997 / Altschul et al., 1990, *J. Mol. Biol.* 215(3):403-410 / Altschul et al., 1993, *Nature Genetics* 3:266-272 / Altschul et al., 1997, *Nuc. Acids Res.* 25:3389-3402 / Anton M. et al., 1995, *J. Virol.*, 69 : 4600-4606 / Araki K et al. (1995) *Proc. Natl. Acad. Sci. USA*. 92(1):160-4. / Aszódi et al., *Proteins:Structure, Function, and Genetics*, Supplement 1:38-42 (1997) / Baubonis W. (1993) *Nucleic Acids Res.* 21(9):2025-9. / Beaucage et al., *Tetrahedron Lett* 1981, 22: 1859-1862 / Block et al, 1990, *Nature* 348, 348-352 / Bradley A., 1987, Production and analysis of chimaeric mice. In: E.J. Robertson (Ed.), *Teratocarcinomas and embryonic stem cells: A practical approach*. IRL Press, Oxford, pp.113. / Bram RJ et al., 1993, *Mol. Cell Biol.*, 13 : 4760-4769 / Brown EL, et al., *Methods Enzymol* 1979;68:109-151 / Brutlag et al. *Comp. App. Biosci.* 6:237-245, 1990 / Bush et al., 1997, *J. Chromatogr.*, 777 : 311-328. / Chai H. et al. (1993) *Biotechnol. Appl. Biochem.*18:259-273. / Chee et al. (1996) *Science*. 274:610-614. / Chen et al. (1987) *Mol. Cell. Biol.* 7:2745-2752. / Chen et al. *Proc. Natl. Acad. Sci. USA* 94/20 10756-10761,1997 / Chen and Kwok *Nucleic Acids Research* 25:347-353 1997 / Cho RJ et al., 1998, *Proc. Natl. Acad. Sci. USA*, 95(7) : 3752-3757. / Chou J.Y., 1989, *Mol. Endocrinol.*, 3: 1511-1514. / Clark A.G. (1990) *Mol. Biol. Evol.* 7:111-122. / Compton J. (1991) *Nature*. 350(6313):91-92. / Davis L.G., et al., *Basic Methods in Molecular Biology*, ed., Elsevier Press, NY, 1986 / Dempster et al., (1977) *J. R. Stat. Soc.*, 39B:1-38. / Eckner R. et al. (1991) *EMBO J.* 10:3513-3522. / Edwards et Leatherbarrow, *Analytical Biochemistry*, 246, 1-6 (1997) / Engvall, E., *Meth. Enzymol.* 70:419 (1980) / Excoffier L. and Slatkin M. (1995) *Mol. Biol. Evol.*, 12(5): 921-927.
- 25 / Feldman and Steg, 1996, *Medecine/Sciences, synthese*, 12:47-55 / Felici F., 1991, *J. Mol. Biol.*, Vol. 222:301-310 / Fields and Song, 1989, *Nature*, 340 : 245-246 / Fisher, D., Chap. 42 in: *Manual of Clinical Immunology*, 2d Ed. (Rose and Friedman, Eds.) Amer. Soc. For Microbiol., Washington, D.C. (1980) / Flotte et al. (1992) *Am. J. Respir. Cell Mol. Biol.* 7:349-356. / Fodor et al. (1991) *Science* 251:767-777. / Fraley et al. (1979) *Proc. Natl. Acad. Sci. USA*. 76:3348-3352.
- 30 / Fromont-Racine M. et al., 1997, *Nature Genetics*, 16(3) : 277-282. / Fuller S. A. et al. (1996) *Immunology in Current Protocols in Molecular Biology*, Ausubel et al. Eds, John Wiley & Sons, Inc., USA. / Furth P.A. et al. (1994) *Proc. Natl. Acad. Sci USA*. 91:9302-9306. / Geysen H. Mario et al. 1984. *Proc. Natl. Acad. Sci. U.S.A.* 81:3998-4002 / Ghosh and Bacchawat, 1991, *Targeting of liposomes to hepatocytes*, IN: *Liver Diseases, Targeted diagnosis and therapy using specific receptors and ligands*. Wu et al. Eds., Marcel Dekker, New York, pp. 87-104. / Gonnet et al., 1992, *Science* 256:1443-1445 / Gopal (1985) *Mol. Cell. Biol.*, 5:1188-1190. / Gossen

- M. et al. (1992) *Proc. Natl. Acad. Sci. USA*. 89:5547-5551. / Gossen M. et al. (1995) *Science*. 268:1766-1769. / Graham et al. (1973) *Virology* 52:456-457. / Green et al., *Ann. Rev. Biochem.* 55:569-597 (1986) / Griffin et al. *Science* 245:967-971 (1989) / Grompe, M. (1993) *Nature Genetics*. 5:111-117. / Grompe, M. et al. (1989) *Proc. Natl. Acad. Sci. U.S.A.* 86:5855-5892.
- 5 / Gu H. et al. (1993) *Cell* 73:1155-1164. / Gu H. et al. (1994) *Science* 265:103-106. / Guatelli J C et al. *Proc. Natl. Acad. Sci. USA*. 35:273-286. / Hacia JG, et al., *Nat Genet* 1996;14(4):441-447 / Haff L. A. and Smirnov I. P. (1997) *Genome Research*, 7:378-388. / Hames B.D. and Higgins S.J. (1985) *Nucleic Acid Hybridization: A Practical Approach*. Hames and Higgins Ed., IRL Press, Oxford. / Harju L, et al., *Clin Chem* 1993;39(11Pt 1):2282-2287 / Harland et al.
- 10 (1985) *J. Cell. Biol.* 101:1094-1095. / Harlow, E., and D. Lane. 1988. *Antibodies A Laboratory Manual*. Cold Spring Harbor Laboratory. pp. 53-242 / Harper JW et al., 1993, *Cell*, 75 : 805-816 / Hawley M.E. et al. (1994) *Am. J. Phys. Anthropol.* 18:104. / Henikoff and Henikoff, 1993, *Proteins* 17:49-61 / Higgins et al., 1996, *Methods Enzymol.* 266:383-402 / Hillier L. and Green P. *Methods Appl.*, 1991, 1: 124-8. / Hoess et al. (1986) *Nucleic Acids Res.*
- 15 14:2287-2300. / Howard et al (1989) *Nature* 342, 154-158 / Huang L. et al. (1996) *Cancer Res* 56(5):1137-1141. / Huygen et al. (1996) *Nature Medicine*. 2(8):893-898. / Izant JG, Weintraub H, *Cell* 1984 Apr;36(4):1007-15 / Julan et al. (1992) *J. Gen. Virol.* 73:3251-3255. / Kanegae Y. et al., 1995 *Nucl. Acids Res.* 23:3816-3821. / Karlin and Altschul, 1990, *Proc. Natl. Acad. Sci. USA* 87:2267-2268 / Khoury J. et al., *Fundamentals of Genetic Epidemiology*, Oxford
- 20 *University Press, NY*, 1993 / Kim U-J. et al. (1996) *Genomics* 34:213-218. / Klein et al. (1987) *Nature*. 327:70-73. / Kohler, G. and Milstein, C., *Nature* 256:495 (1975) / Koller et al. (1992) *Annu. Rev. Immunol.* 10:705-730. / Kozal MJ, et al., *Nat Med* 1996;2(7):753-759 / Landegren U. et al. (1998) *Genome Research*, 8:769-776. / Lander and Schork, *Science*, 265, 2037-2048, 1994 / Lange K. (1997) *Mathematical and Statistical Methods for Genetic Analysis*. Springer,
- 25 New York. / Lenhard T. et al. (1996) *Gene*. 169:187-190. / Linton M.F. et al. (1993) *J. Clin. Invest.* 92:3029-3037. / Liu Z. et al. (1994) *Proc. Natl. Acad. Sci. USA*. 91: 4528-4262. / Livak KJ, Hainer JW, *Hum Mutat* 1994;3(4):379-385 / Livak et al., *Nature Genetics*, 9:341-342, 1995 / Lucas A.H., 1994, In : *Development and Clinical Uses of Haemophilus b Conjugate*; / Mackey K, et al., 1998, *Mol Biotechnol*, 9(1):1-5 / Mansour S.L. et al. (1988) *Nature*. 336:348-
- 30 352. / Marshall R. L. et al. (1994) *PCR Methods and Applications*. 4:80-84. / McCormick et al. (1994) *Genet. Anal. Tech. Appl.* 11:158-164. / McLaughlin B.A. et al. (1996) *Am. J. Hum. Genet.* 59:561-569. / Morton N.E., *Am.J. Hum. Genet.*, 7:277-318, 1955 / Muzyczka et al. (1992) *Curr. Topics in Micro. and Immunol.* 158:97-129. / Nada S. et al. (1993) *Cell* 73:1125-1135. / Nagy A. et al., 1993, *Proc. Natl. Acad. Sci. USA*, 90: 8424-8428. / Nangaku M.
- 35 (1994) *Cell* 79, 1209-1220 / Narang SA, et al., *Methods Enzymol* 1979;68:90-98 / Neda et al. (1991) *J. Biol. Chem.* 266:14143-14146. / Newton et al. (1989) *Nucleic Acids Res.* 17:2503-2516. / Nickerson D.A. et al. (1990) *Proc. Natl. Acad. Sci. U.S.A.* 87:8923-8927. / Nicolau

- C. et al., 1987, *Methods Enzymol.*, **149**:157-76. / Nicolau et al. (1982) *Biochim. Biophys. Acta.* **721**:185-190. / Nyren P, Pettersson B, Uhlen M, *Anal Biochem* 1993;**208**(1):171-175 / O'Reilly et al. (1992) *Baculovirus Expression Vectors: A Laboratory Manual*. W. H. Freeman and Co., New York. / Ohno et al. (1994) *Science*. **265**:781-784. / Oldenburg K.R. et al., 1992, *Proc. Natl. Acad. Sci.*, **89**:5393-5397. / Orita et al. (1989) *Proc. Natl. Acad. Sci. U.S.A.* **86**: 2776-2770. / Ott J., *Analysis of Human Genetic Linkage*, John Hopkins University Press, Baltimore, 1991 / Ouchterlony, O. et al., Chap. 19 in: Handbook of Experimental Immunology D. Wier (ed) Blackwell (1973) / Parmley and Smith, *Gene*, 1988, **73**:305-318 / Pastinen et al., *Genome Research* 1997; **7**:606-614 / Pearson and Lipman, 1988, *Proc. Natl. Acad. Sci. USA* **85**(8):2444-2448 / Pease S. and William R.S., 1990, *Exp. Cell. Res.*, **190**: 209-211. / Perlin et al. (1994) *Am. J. Hum. Genet.* **55**:777-787. / Peterson et al., 1993, *Proc. Natl. Acad. Sci. USA*, **90** : 7593-7597. / Potter et al. (1984) *Proc. Natl. Acad. Sci. U.S.A.* **81**(22):7161-7165. / Ramunsen et al., 1997, *Electrophoresis*, **18** : 588-598. / Reid L.H. et al. (1990) *Proc. Natl. Acad. Sci. U.S.A.* **87**:4299-4303. / Risch, N. and Merikangas, K. (*Science*, **273**:1516-1517, 1996 /
- 15 Robertson E., 1987, Embryo-derived stem cell lines. In: E.J. Robertson Ed. *Teratocarcinomas and embryonic stem cells: a practical approach*. IRL Press, Oxford, pp. 71. / Rossi et al., *Pharmacol. Ther.* **50**:245-254, (1991) / Roth J.A. et al. (1996) *Nature Medicine*. **2**(9):985-991. / Roux et al. (1989) *Proc. Natl. Acad. Sci. U.S.A.* **86**:9079-9083. / Ruano et al. (1990) *Proc. Natl. Acad. Sci. U.S.A.* **87**:6296-6300. / Sambrook, J., et al. (1989) *Molecular Cloning: A Laboratory*
- 20 *Manual*. 2ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York. / Samson M, et al. (1996) *Nature*, **382**(6593):722-725. / Samulski et al. (1989) *J. Virol.* **63**:3822-3828. / Sanchez-Pescador R. (1988) *J. Clin. Microbiol.* **26**(10):1934-1938. / Sarkar, G. and Sommer S.S. (1991) *Biotechniques*. / Sato-Yoshitake et al., *J Biol Chem* 1992 Nov **25**;267(33):23930-6 / Sauer B. et al. (1988) *Proc. Natl. Acad. Sci. U.S.A.* **85**:5166-5170. / Schedl A. et al., 1993a,
- 25 *Nature*, **362**: 258-261. / Schedl et al., 1993b, *Nucleic Acids Res.*, **21**: 4783-4787. / Schneider et al. (1997) *Arlequin: A Software For Population Genetics Data Analysis*. University of Geneva. / Scholey, 1993, *Motility assays for Motor Proteins*, (San Diego, California : Academic Press) / Schwartz and Dayhoff, eds., 1978, *Matrices for Detecting Distance Relationships: Atlas of Protein Sequence and Structure*, Washington: National Biomedical Research Foundation / Schaid D.J. et
- 30 al., *Genet. Epidemiol.*, **13**:423-450, 1996 / Sczakiel G. et al. (1995) *Trends Microbiol.* **3**(6):213-217. / Shay J.W. et al., 1991, *Biochem. Biophys. Acta*, **1072**: 1-7. / Sheffield, V.C. et al. (1991) *Proc. Natl. Acad. Sci. U.S.A.* **49**:699-706. / Shizuya et al. (1992) *Proc. Natl. Acad. Sci. U.S.A.* **89**:8794-8797. / Shoemaker DD, Lashkari DA, Morris D, Mittmann M, Davis RW, *Nat Genet* 1996;**14**(4):450-456 / Spielmann S. et al., *Am. J. Hum. Genet.*, **52**:506-516, 1993 /
- 35 Spielmann S. and Ewens W.J., *Am. J. Hum. Genet.*, **62**:450-458, 1998 / Smith (1957) *Ann. Hum. Genet.* **21**:254-276. / Smith et al. (1983) *Mol. Cell. Biol.* **3**:2156-2165. / Sowdhamini et al., *Protein Engineering* **10**:207, 215 (1997) / Sternberg N.L. (1992) *Trends Genet.* **8**:1-16. /

- Sternberg N.L. (1994) *Mamm. Genome*. 5:397-404. / Syvanen AC, *Clin Chim Acta* 1994;226(2):225-236 / Szabo A. et al. *Curr Opin Struct Biol* 5, 699-705 (1995) / Tacson et al. (1996) *Nature Medicine*. 2(8):888-892. / Te Riele et al. (1990) *Nature*. 348:649-651. / Terwilliger J.D. and Ott J., *Handbook of Human Genetic Linkage*, John Hopkins University Press, London, 1994 / Thomas K.R. et al. (1986) *Cell*. 44:419-428. / Thomas K.R. et al. (1987) *Cell*. 51:503-512. / Thompson et al., 1994, *Nucleic Acids Res*. 22(2):4673-4680 / Tur-Kaspa et al. (1986) *Mol. Cell. Biol*. 6:716-718. / Tyagi et al. (1998) *Nature Biotechnology*. 16:49-53. / Urdea M.S. (1988) *Nucleic Acids Research*. 11:4937-4957. / Urdea M.S. et al. (1991) *Nucleic Acids Symp. Ser.* 24:197-200. / Vaitukaitis, J. et al. *J. Clin. Endocrinol. Metab.* 33:988-991 (1971) / Valadon P., et al., 1996, *J. Mol. Biol.*, 261:11-22. / Van der Lugt et al. (1991) *Gene*. 105:263-267. / Vlasak R. et al. (1983) *Eur. J. Biochem*. 135:123-126. / Wabiko et al. (1986) *DNA*. 5(4):305-314. / Walker et al. (1996) *Clin. Chem*. 42:9-13. / Wang et al., 1997, *Chromatographia*, 44 : 205-208. / Weir, B.S. (1996) *Genetic data Analysis II: Methods for Discrete population genetic Data*, Sinauer Assoc., Inc., Sunderland, MA, U.S.A. / Westerink M.A.J., 1995, *Proc. Natl. Acad. Sci.*, 92:4021-4025 / White, M.B. et al. (1992) *Genomics*. 12:301-306. / White, M.B. et al. (1997) *Genomics*. 12:301-306. / Wong et al. (1980) *Gene*. 10:87-94. / Wood S.A. et al., 1993, *Proc. Natl. Acad. Sci. USA*, 90: 4582-4585. / Wu and Wu (1987) *J. Biol. Chem*. 262:4429-4432. / Wu and Wu (1988) *Biochemistry*. 27:887-892. / Wu et al. (1989) *Proc. Natl. Acad. Sci. U.S.A.* 86:2757. / Yagi T. et al. (1990) *Proc. Natl. Acad. Sci. U.S.A.* 87:9918-9922. / Yonekawa Y. et al. (1998) *J. Cell. Biol*. 141, 431-441 / Zhao et al., *Am. J. Hum. Genet.*, 63:225-240, 1998 / Zou Y. R. et al. (1994) *Curr. Biol*. 4:1099-1103.

CLAIMS

1. An isolated, purified or recombinant polynucleotide comprising a contiguous span of at least 12 nucleotides of SEQ ID No 1 or the complementary sequence thereof, wherein said contiguous span comprises either :

- 5 - at least 1 of the following nucleotide positions of SEQ ID No 1: 1-39624, 39705-40589, 40666-43629, 43710-44203, 44311-45125, 45210-45440, 45622-45717, 45791-68580, 68675-70246, 70396-72421, 72601-73295, 73434-74648, 74898-83055, 83175-85192, 85279-85609, 85740-85906, 86070-88304, 88396-90585, 90705-91767, 91824-94380, 94490-96296, 96364-97184, 97270-101167, 101274-109465, 109581-110228, 110363-111819, 111882-113636, 113783-10 113945, 114186-117002, 117075-119676, and 119677-121162; or,
- a G at position 7159 of SEQ ID No 1.

2. An isolated, purified or recombinant polynucleotide comprising a contiguous span of at least 12 nucleotides of SEQ ID No 2 or the complementary sequence thereof, wherein said
15 contiguous span comprises:

- at least 1 of the following nucleotide positions of SEQ ID No 2: 1-1600, 1751-2138, 2332-2539, 2659-3829 and 8885-10884; or,
- a C either at position 2551 or 4500 of SEQ ID No 2.

20 3. An isolated, purified or recombinant polynucleotide comprising a contiguous span of at least 12 nucleotides of SEQ ID No 3 or the complementary sequence thereof, wherein said contiguous span comprises either :

- at least 1 of the following nucleotide positions of SEQ ID No 3: 391-1619 and 6988-10682; or,
- 25 - a C at position 5487, or a C at position 6265 of SEQ ID No 3.

4. An isolated, purified, or recombinant polynucleotide consisting essentially of a contiguous span of 8 to 50 nucleotides of SEQ ID Nos 1-3 and 5-8 or the complement thereof, wherein said span includes a *HKLP*-related biallelic marker in said sequence.

30

5. A polynucleotide according to claim 4, wherein said *HKLP*-related biallelic marker is selected from the group consisting of A1 to A32, and the complements thereof.

6. A polynucleotide according to claim 4, wherein said *HKLP*-related biallelic marker is
35 selected from the group consisting of A1 to A22 and A25 to A32, and the complements thereof.

7. A polynucleotide according to claim 4, wherein said *HKLP*-related biallelic marker is selected from the group consisting of A23 and A24, and the complements thereof.

8. A polynucleotide according to any one of claims 4 to 7, wherein said contiguous span is 18 to 35 nucleotides in length and said biallelic marker is within 4 nucleotides of the center of said polynucleotide.

9. A polynucleotide according to claim 8, wherein said polynucleotide consists of said contiguous span and said contiguous span is 25 nucleotides in length and said biallelic marker is at the center of said polynucleotide.

10. A polynucleotide according to claim 9, wherein said polynucleotide consists essentially of a sequence selected from the following sequences: P1 to P30, and the complementary sequences thereto.

11. A polynucleotide according to any one of claims 4 to 7, wherein the 3' end of said contiguous span is located at the 3' end of said polynucleotide and said biallelic marker is present at the 3' end of said polynucleotide.

12. A polynucleotide according to any one of claims 1 to 3, wherein the 3' end of said contiguous span is present at the 3' end of said polynucleotide.

13. A polynucleotide according to claim 12, wherein the 3' end of said polynucleotide is located within 20 nucleotides upstream of a *HKLP*-related biallelic marker in said sequence.

14. A polynucleotide according to claim 13, wherein the 3' end of said polynucleotide is located 1 nucleotide upstream of said *HKLP*-related biallelic marker in said sequence.

15. A polynucleotide according to claim 14, wherein said polynucleotide consists essentially of a sequence selected from the following sequences: D1 to D30, and E1 to E30.

16. An isolated, purified, or recombinant polynucleotide consisting essentially of a sequence selected from the following sequences: B1 to B19 and C1 to C25.

17. An isolated, purified, or recombinant polynucleotide which encodes a polypeptide comprising a contiguous span of at least 6 amino acids of SEQ ID No 4, wherein said contiguous span includes at least 1 of the amino acid positions 1-478 of the SEQ ID No 4.

18. A polynucleotide according to any one of claims 1 to 17 attached to a solid support.
19. An array of polynucleotides comprising at least one polynucleotide according to claim
5 18.
20. An array according to claim 19, wherein said array is addressable.
21. A polynucleotide according to any one of claims 1 to 17 further comprising a label.
10
22. A recombinant vector comprising a polynucleotide according to any one of claims 1 to
17.
23. A host cell comprising a recombinant vector according to claim 22.
15
24. A non-human host animal or mammal comprising a recombinant vector according to
claim 22.
25. A mammalian host cell comprising an *HKLP* gene disrupted by homologous
20 recombination with a knock out vector, comprising a polynucleotide according to any one of claims
1 to 17.
26. A non-human host mammal comprising a *HKLP* gene disrupted by homologous
recombination with a knock out vector, comprising a polynucleotide according to any one of claims
25 1 to 17.
27. A method of genotyping comprising determining the identity of a nucleotide at a *HKLP*-
related biallelic marker or the complement thereof in a biological sample.
- 30 28. A method according to claim 27, wherein said biological sample is derived from a single
subject.
29. A method according to claim 28, wherein the identity of the nucleotides at said biallelic
marker is determined for both copies of said biallelic marker present in said individual's genome.
35
30. A method according to claim 27, wherein said biological sample is derived from
multiple subjects.

31. A method according to claim 27, further comprising amplifying a portion of said sequence comprising the biallelic marker prior to said determining step.

5 32. A method according to claim 31, wherein said amplifying is performed by PCR.

33. A method according to claim 27, wherein said determining is performed by a hybridization assay.

10 34. A method according to claim 27, wherein said determining is performed by a sequencing assay.

35. A method according to claim 27, wherein said determining is performed by a microsequencing assay.

15 36. A method according to claim 27, wherein said determining is performed by an enzyme-based mismatch detection assay.

37. A method of estimating the frequency of an allele of a *HKLP*-related biallelic marker in
20 a population comprising:

a) genotyping individuals from said population for said biallelic marker according to the method of claim 27; and

b) determining the proportional representation of said biallelic marker in said population.

25 38. A method of detecting an association between a genotype and a trait, comprising the steps of:

a) determining the frequency of at least one *HKLP*-related biallelic marker in trait positive population according to the method of claim 37;

b) determining the frequency of said *HKLP*-related biallelic marker in a control population
30 according to the method of claim 37; and

c) determining whether a statistically significant association exists between said genotype and said trait.

39. A method of estimating the frequency of a haplotype for a set of biallelic markers in a
35 population, comprising:

a) genotyping at least one *HKLP*-related biallelic marker according to claim 29 for each individual in said population;

b) genotyping a second biallelic marker by determining the identity of the nucleotides at said second biallelic marker for both copies of said second biallelic marker present in the genome of each individual in said population; and

c) applying a haplotype determination method to the identities of the nucleotides determined
5 in steps a) and b) to obtain an estimate of said frequency.

40. A method according to claim 39, wherein said haplotype determination method is selected from the group consisting of asymmetric PCR amplification, double PCR amplification of specific alleles, the Clark method, or an expectation maximization algorithm.

10

41. A method of detecting an association between a haplotype and a trait, comprising the steps of:

a) estimating the frequency of at least one haplotype in a trait positive population according to the method of claim 39;

15 b) estimating the frequency of said haplotype in a control population according to the method of claim 39; and

c) determining whether a statistically significant association exists between said haplotype and said trait.

20 42. A method according to claim 38, wherein said genotyping steps a) and b) are performed on a single pooled biological sample derived from each of said populations.

43. A method according to claim 38, wherein said genotyping steps a) and b) performed separately on biological samples derived from each individual in said populations.

25

44. A method according to either claim 38 or 41, wherein said control population is a trait negative population.

45. A method according to either claim 38 or 41, wherein said case control population is a
30 random population.

46. An isolated, purified, or recombinant polypeptide comprising a contiguous span of at least 6 amino acids of SEQ ID No 4, wherein said contiguous span includes at least 1 of the amino acid positions 1-478 of the SEQ ID No 4.

35

47. An isolated or purified antibody composition are capable of selectively binding to an epitope-containing fragment of a polypeptide according to claim 46, wherein said epitope comprises at least 1 of the amino acid positions 1-478 of the SEQ ID No 4.

5 48. A method according to any one of claims 27 to 45 wherein said *HKLP*-related biallelic marker is selected from the group consisting of A1 to A32, and the complements thereof.

49. A method according to any one of claims 27 to 45 wherein said *HKLP*-related biallelic marker is selected from the group consisting of A1 to A22 and A25 to A32, and the complements
10 thereof.

50. A method according to any one of claims 27 to 45 wherein said *HKLP*-related biallelic marker is selected from the group consisting of A23 and A24, and the complements thereof.

15 51. A diagnostic kit comprising a polynucleotide according to any one of claims 1 to 21.

52. Use of a polynucleotide comprising a contiguous span of at least 12 nucleotides of a sequence selected from the group consisting of the SEQ ID Nos 1-3 and 5-8 or the complementary sequence thereto for determining the identity of the nucleotide at a *HKLP*-related biallelic marker
20

53. Use according to claim 52 in a microsequencing assay, wherein the 3' end of said contiguous span is located at the 3' end of said polynucleotide and wherein the 3' end of said polynucleotide is located 1 nucleotide upstream of said *HKLP* related biallelic marker in said sequence.

25 54. Use according to claim 52 in a hybridization assay, wherein said span includes said *HKLP* -related biallelic marker.

55. Use according to claim 52 in a specific amplification assay, wherein the 3' end of said
30 contiguous span is located at the 3' end of said polynucleotide and said biallelic marker is present at the 3' end of said polynucleotide.

56. Use according to claim 52 in a sequencing assay, wherein the 3' end of said contiguous span is located at the 3' end of said polynucleotide.

35 57. Use according to any one of claims 52-56, wherein said *HKLP* -related biallelic is a biallelic marker selected in the group consisting of the biallelic markers A1 to A32.

58. A computer readable medium having stored thereon a sequence selected from the group consisting of a nucleic acid code comprising one of the following:

- a) a contiguous span of at least 12 nucleotides of SEQ ID No 1, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 1: 1-39624, 39705-40589, 40666-43629, 43710-44203, 44311-45125, 45210-45440, 45622-45717, 45791-68580, 68675-70246, 70396-72421, 72601-73295, 73434-74648, 74898-83055, 83175-85192, 85279-85609, 85740-85906, 86070-88304, 88396-90585, 90705-91767, 91824-94380, 94490-96296, 96364-97184, 97270-101167, 101274-109465, 109581-110228, 110363-111819, 111882-113636, 113783-113945, 114186-117002, 117075-119676, and 119677-121162;
- b) a contiguous span of at least 12 nucleotides of SEQ ID No 2 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 2: 1-1600, 1751-2138, 2332-2539, 2659-3829 and 8885-10884;
- c) a contiguous span of at least 12 nucleotides of SEQ ID No 1 or the complements thereof, wherein said contiguous span comprises a G at position 7159 of SEQ ID No 1;
- d) a contiguous span of at least 12 nucleotides of SEQ ID No 4 or the complements thereof, wherein said contiguous span comprises a C either at position 2551 or 4500 of SEQ ID No 4;
- e) a contiguous span of at least 12 nucleotides of SEQ ID No 3 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 3: 391-1619 and 6988-10682;
- f) a contiguous span of at least 12 nucleotides of SEQ ID No 3 or the complements thereof, wherein said contiguous span comprises a nucleotide selected in the group consisting of a C at position 5487, and a C at position 6265 of SEQ ID No 3; and
- g) a nucleotide sequence complementary to any one of the contiguous spans of a), b), c), d), e), or f).

59. A computer readable medium having stored thereon a sequence consisting of a polypeptide code comprising a contiguous span of at least 6 amino acids of SEQ ID No 4, wherein said contiguous span includes at least 1 of the amino acid positions 1-478 of the SEQ ID No 4.

60. A computer system comprising a processor and a data storage device wherein said data storage device a computer readable medium according to with claim 58 or 59.

61. A computer system according to claim 60, further comprising a sequence comparer and a data storage device having reference sequences stored thereon.

62. A computer system of Claim 61 wherein said sequence comparer comprises a computer program which indicates polymorphisms.

63. A computer system of Claim 60 further comprising an identifier which identifies
5 features in said sequence.

64. A method for comparing a first sequence to a reference sequence, comprising the steps
of:

reading said first sequence and said reference sequence through use of a computer program
10 which compares sequences; and

determining differences between said first sequence and said reference sequence with said
computer program,

wherein said first sequence is selected from the group consisting of a nucleic acid code
comprising one of the following:

15 a) a contiguous span of at least 12 nucleotides of SEQ ID No 1, wherein said
contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of
SEQ ID No 1: 1-39624, 39705-40589, 40666-43629, 43710-44203, 44311-45125, 45210-
45440, 45622-45717, 45791-68580, 68675-70246, 70396-72421, 72601-73295, 73434-
74648, 74898-83055, 83175-85192, 85279-85609, 85740-85906, 86070-88304, 88396-
20 90585, 90705-91767, 91824-94380, 94490-96296, 96364-97184, 97270-101167, 101274-
109465, 109581-110228, 110363-111819, 111882-113636, 113783-113945, 114186-
117002, 117075-119676, and 119677-121162;

b) a contiguous span of at least 12 nucleotides of SEQ ID No 2 or the complements
thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following
25 nucleotide positions of SEQ ID No 2: 1-1600, 1751-2138, 2332-2539, 2659-3829 and 8885-
10884;

c) a contiguous span of at least 12 nucleotides of SEQ ID No 1 or the complements
thereof, wherein said contiguous span comprises a G at position 7159 of SEQ ID No 1;

d) a contiguous span of at least 12 nucleotides of SEQ ID No 4 or the complements
30 thereof, wherein said contiguous span comprises a C either at position 2551 or 4500 of SEQ
ID No 4;

e) a contiguous span of at least 12 nucleotides of SEQ ID No 3 or the complements
thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following
nucleotide positions of SEQ ID No 3: 391-1619 and 6988-10682;

35 f) a contiguous span of at least 12 nucleotides of SEQ ID No 3 or the complements
thereof, wherein said contiguous span comprises a nucleotide selected in the group
consisting of a C at position 5487, and a C at position 6265 of SEQ ID No 3; and

g) a nucleotide sequence complementary to any one of the contiguous spans of a), b), c), d), e), or f); and

a polypeptide code comprising a contiguous span of at least 6 amino acids of SEQ ID No 4, wherein said contiguous span includes at least 1 of the amino acid positions 1-478 of the SEQ ID No 4.

65. A method according to Claim 64, wherein said step of determining differences between the first sequence and the reference sequence comprises identifying at least one polymorphism.

66. A method for identifying a feature in a sequence, comprising the steps of:
reading said sequence through the use of a computer program which identifies features in sequences; and

identifying features in said sequence with said computer program;

wherein said sequence is selected from the group consisting of a nucleic acid code

comprising one of the following:

a) a contiguous span of at least 12 nucleotides of SEQ ID No 1, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 1: 1-39624, 39705-40589, 40666-43629, 43710-44203, 44311-45125, 45210-45440, 45622-45717, 45791-68580, 68675-70246, 70396-72421, 72601-73295, 73434-74648, 74898-83055, 83175-85192, 85279-85609, 85740-85906, 86070-88304, 88396-90585, 90705-91767, 91824-94380, 94490-96296, 96364-97184, 97270-101167, 101274-109465, 109581-110228, 110363-111819, 111882-113636, 113783-113945, 114186-117002, 117075-119676, and 119677-121162;

b) a contiguous span of at least 12 nucleotides of SEQ ID No 2 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 2: 1-1600, 1751-2138, 2332-2539, 2659-3829 and 8885-10884;

c) a contiguous span of at least 12 nucleotides of SEQ ID No 1 or the complements thereof, wherein said contiguous span comprises a G at position 7159 of SEQ ID No 1;

d) a contiguous span of at least 12 nucleotides of SEQ ID No 4 or the complements thereof, wherein said contiguous span comprises a C either at position 2551 or 4500 of SEQ ID No 4;

e) a contiguous span of at least 12 nucleotides of SEQ ID No 3 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 3: 391-1619 and 6988-10682;

f) a contiguous span of at least 12 nucleotides of SEQ ID No 3 or the complements thereof, wherein said contiguous span comprises a nucleotide selected in the group consisting of a C at position 5487, and a C at position 6265 of SEQ ID No 3; and

5 g) a nucleotide sequence complementary to any one of the contiguous spans of a), b), c), d), e), or f); and

a polypeptide code comprising a contiguous span of at least 6 amino acids of SEQ ID No 4, wherein said contiguous span includes at least 1 of the amino acid positions 1-478 of the SEQ ID No 4.

10 67. A method for the screening of a candidate substance interacting with a HKLP polypeptide comprising the following steps :

- a) providing a polypeptide consisting of a HKLP protein or a fragment comprising a contiguous span of at least 6 amino acids amino acids of SEQ ID No 4, wherein said contiguous span includes at least 1 of the amino acid positions 1-478 of the SEQ ID No 4 or a variant thereof;
- 15 b) obtaining a candidate substance;
- c) bringing into contact said polypeptide with said candidate substance;
- d) detecting the complexes formed between said polypeptide and said candidate substance.

1/4




FIG.1

2/4




FIG.2

3/4




FIG.3

4/4




FIG.4

<110> Genset

<120> DNA encoding a novel kinesin-like protein

<130> 64.W01

<150> US 60/130,217

<151> 1999-04-20

<160> 10

<170> Patent.pm

<210> 1

<211> 121162

<212> DNA

<213> Homo sapiens

<220>

<221> exon

<222> 4844..4920

<223> exon

<220>

<221> exon

<222> 7058..7237

<223> exon

<220>

<221> exon

<222> 10469..10534

<223> exon

<220>

<221> exon

<222> 15940..16118

<223> exon

<220>

<221> exon

<222> 16712..16823

<223> exon

<220>

<221> exon

<222> 20054..20131

<223> exon

<220>

<221> exon

<222> 20793..20858

<223> exon

<220>

<221> exon

<222> 21558..21575

<223> exon

<220>

<221> exon

<222> 23971..24046

<223> exon

<220>
<221> exon
<222> 24864..24942
<223> exon

<220>
<221> exon
<222> 26529..26671
<223> exon

<220>
<221> exon
<222> 27640..27681
<223> exon

<220>
<221> exon
<222> 30863..31074
<223> exon

<220>
<221> exon
<222> 39625..39704
<223> exon

<220>
<221> exon
<222> 40590..40665
<223> exon

<220>
<221> exon
<222> 43630..43709
<223> exon

<220>
<221> exon
<222> 44204..44310
<223> exon

<220>
<221> exon
<222> 45126..45209
<223> exon

<220>
<221> exon
<222> 45441..45621
<223> exon

<220>
<221> exon
<222> 45718..45790
<223> exon

<220>
<221> exon
<222> 68581..68674
<223> exon

<220>
<221> exon

<222> 70247..70395
<223> exon

<220>
<221> exon
<222> 72422..72600
<223> exon

<220>
<221> exon
<222> 73296..73433
<223> exon

<220>
<221> exon
<222> 74649..74897
<223> exon

<220>
<221> exon
<222> 83056..83174
<223> exon

<220>
<221> exon
<222> 85193..85278
<223> exon

<220>
<221> exon
<222> 85610..85739
<223> exon

<220>
<221> exon
<222> 85907..86069
<223> exon

<220>
<221> exon
<222> 88305..88395
<223> exon

<220>
<221> exon
<222> 90586..90704
<223> exon

<220>
<221> exon
<222> 91768..91823
<223> exon

<220>
<221> exon
<222> 94381..94489
<223> exon

<220>
<221> exon
<222> 96297..96363
<223> exon

<220>
<221> exon
<222> 97185..97269
<223> exon

<220>
<221> exon
<222> 101168..101273
<223> exon

<220>
<221> exon
<222> 109466..109580
<223> exon

<220>
<221> exon
<222> 110229..110362
<223> exon

<220>
<221> exon
<222> 111820..111881
<223> exon

<220>
<221> exon
<222> 113637..113782
<223> exon

<220>
<221> exon
<222> 113946..114185
<223> exon

<220>
<221> exon
<222> 117003..117074
<223> exon

<220>
<221> exon
<222> 119677..119798
<223> exon

<220>
<221> allele
<222> 7159
<223> 12-809-119 : polymorphic base G or C

<220>
<221> allele
<222> 16369
<223> 12-805-115 : polymorphic base A or G

<220>
<221> allele
<222> 26310
<223> 12-790-396 : polymorphic base A or G

<220>
<221> allele
<222> 31112

5

<223> 12-791-211 : polymorphic bas A or G

<220>
<221> allele
<222> 33808
<223> 12-803-125 : polymorphic base A or T

<220>
<221> allele
<222> 34255
<223> 99-33040-321 : polymorphic base C or T

<220>
<221> allele
<222> 35293
<223> 12-810-77 : polymorphic base A or G

<220>
<221> allele
<222> 45892
<223> 12-787-103 : polymorphic base A or G

<220>
<221> allele
<222> 54497
<223> 12-793-383 : polymorphic base G or T

<220>
<221> allele
<222> 56582
<223> 12-792-233 : polymorphic base A or G

<220>
<221> allele
<222> 60336
<223> 99-41009-244 : polymorphic base A or G

<220>
<221> allele
<222> 60469
<223> 99-41009-111 : polymorphic base C or T

<220>
<221> allele
<222> 79063
<223> 12-593-287 : insertion of TAAAT

<220>
<221> allele
<222> 79176
<223> 12-593-174 : polymorphic base C or T

<220>
<221> allele
<222> 80016
<223> 12-589-152 : polymorphic base G or T

<220>
<221> allele
<222> 87463
<223> 12-785-200 : polymorphic base C or T

<220>

<221> allele
<222> 87656
<223> 12-785-393 : polymorphic base A or G

<220>
<221> allele
<222> 101602
<223> 12-588-103 : polymorphic base A or G

<220>
<221> allele
<222> 104391
<223> 12-603-191 : polymorphic base C or T

<220>
<221> allele
<222> 117430
<223> 12-586-414 : polymorphic base A or G

<220>
<221> allele
<222> 117459
<223> 12-586-443 : deletion of C

<220>
<221> primer_bind
<222> 7041..7060
<223> 12-809.pu

<220>
<221> primer_bind
<222> 7545..7565
<223> 12-809.rp complement

<220>
<221> primer_bind
<222> 16004..16024
<223> 12-805.rp

<220>
<221> primer_bind
<222> 16464..16483
<223> 12-805.pu complement

<220>
<221> primer_bind
<222> 26232..26252
<223> 12-790.rp

<220>
<221> primer_bind
<222> 26685..26705
<223> 12-790.pu complement

<220>
<221> primer_bind
<222> 30902..30922
<223> 12-791.pu

<220>
<221> primer_bind
<222> 31282..31301
<223> 12-791.rp complement

<220>
<221> primer_bind
<222> 33476..33496
<223> 12-803.rp

<220>
<221> primer_bind
<222> 33912..33932
<223> 12-803.pu complement

<220>
<221> primer_bind
<222> 33934..33953
<223> 99-33040.pu

<220>
<221> primer_bind
<222> 34364..34383
<223> 99-33040.rp complement

<220>
<221> primer_bind
<222> 34918..34938
<223> 12-810.rp

<220>
<221> primer_bind
<222> 35350..35369
<223> 12-810.pu complement

<220>
<221> primer_bind
<222> 45465..45485
<223> 12-787.rp

<220>
<221> primer_bind
<222> 45974..45993
<223> 12-787.pu complement

<220>
<221> primer_bind
<222> 54361..54381
<223> 12-793.rp

<220>
<221> primer_bind
<222> 54861..54879
<223> 12-793.pu complement

<220>
<221> primer_bind
<222> 56365..56385
<223> 12-792.rp

<220>
<221> primer_bind
<222> 56793..56813
<223> 12-792.pu complement

<220>
<221> primer_bind

<222> 60111..60130
<223> 99-41009.rp

<220>
<221> primer_bind
<222> 60562..60580
<223> 99-41009.pu complement

<220>
<221> primer_bind
<222> 78815..78835
<223> 12-593.rp

<220>
<221> primer_bind
<222> 79332..79349
<223> 12-593.pu complement

<220>
<221> primer_bind
<222> 79706..79726
<223> 12-589.rp

<220>
<221> primer_bind
<222> 80149..80167
<223> 12-589.pu complement

<220>
<221> primer_bind
<222> 87264..87284
<223> 12-785.pu

<220>
<221> primer_bind
<222> 87753..87773
<223> 12-785.rp complement

<220>
<221> primer_bind
<222> 101161..101178
<223> 12-588.rp

<220>
<221> primer_bind
<222> 101686..101704
<223> 12-588.pu complement

<220>
<221> primer_bind
<222> 104131..104151
<223> 12-603.rp

<220>
<221> primer_bind
<222> 104558..104578
<223> 12-603.pu complement

<220>
<221> primer_bind
<222> 117017..117036
<223> 12-586.pu

<220>
<221> primer_bind
<222> 117481..117501
<223> 12-586.rp complement

<220>
<221> primer_bind
<222> 7140..7158
<223> 12-809-119.mis

<220>
<221> primer_bind
<222> 7160..7178
<223> 12-809-119.mis complement

<220>
<221> primer_bind
<222> 16350..16368
<223> 12-805-115.mis

<220>
<221> primer_bind
<222> 16370..16388
<223> 12-805-115.mis complement

<220>
<221> primer_bind
<222> 26291..26309
<223> 12-790-396.mis

<220>
<221> primer_bind
<222> 26311..26329
<223> 12-790-396.mis complement

<220>
<221> primer_bind
<222> 31093..31111
<223> 12-791-211.mis

<220>
<221> primer_bind
<222> 31113..31131
<223> 12-791-211.mis complement

<220>
<221> primer_bind
<222> 33789..33807
<223> 12-803-125.mis

<220>
<221> primer_bind
<222> 33809..33827
<223> 12-803-125.mis complement

<220>
<221> primer_bind
<222> 34236..34254
<223> 99-33040-321.mis

<220>
<221> primer_bind
<222> 34256..34274

<223> 99-33040-321.mis complement

<220>

<221> primer_bind

<222> 35274..35292

<223> 12-810-77.mis

<220>

<221> primer_bind

<222> 35294..35312

<223> 12-810-77.mis complement

<220>

<221> primer_bind

<222> 45873..45891

<223> 12-787-103.mis

<220>

<221> primer_bind

<222> 45893..45911

<223> 12-787-103.mis complement

<220>

<221> primer_bind

<222> 54478..54496

<223> 12-793-383.mis

<220>

<221> primer_bind

<222> 54498..54516

<223> 12-793-383.mis complement

<220>

<221> primer_bind

<222> 56563..56581

<223> 12-792-233.mis

<220>

<221> primer_bind

<222> 56583..56601

<223> 12-792-233.mis complement

<220>

<221> primer_bind

<222> 60317..60335

<223> 99-41009-244.mis

<220>

<221> primer_bind

<222> 60337..60355

<223> 99-41009-244.mis complement

<220>

<221> primer_bind

<222> 60450..60468

<223> 99-41009-111.mis

<220>

<221> primer_bind

<222> 60470..60488

<223> 99-41009-111.mis complement

<220>

<221> primer_bind
<222> 79157..79175
<223> 12-593-174.mis

<220>
<221> primer_bind
<222> 79177..79195
<223> 12-593-174.mis complement

<220>
<221> primer_bind
<222> 79997..80015
<223> 12-589-152.mis

<220>
<221> primer_bind
<222> 80017..80035
<223> 12-589-152.mis complement

<220>
<221> primer_bind
<222> 87444..87462
<223> 12-785-200.mis

<220>
<221> primer_bind
<222> 87464..87482
<223> 12-785-200.mis complement

<220>
<221> primer_bind
<222> 87637..87655
<223> 12-785-393.mis

<220>
<221> primer_bind
<222> 87657..87675
<223> 12-785-393.mis complement

<220>
<221> primer_bind
<222> 101583..101601
<223> 12-588-103.mis

<220>
<221> primer_bind
<222> 101603..101621
<223> 12-588-103.mis complement

<220>
<221> primer_bind
<222> 104372..104390
<223> 12-603-191.mis

<220>
<221> primer_bind
<222> 104392..104410
<223> 12-603-191.mis complement

<220>
<221> primer_bind
<222> 117411..117429
<223> 12-586-414.mis

<220>
<221> primer_bind
<222> 117431..117449
<223> 12-586-414.mis complement

<220>
<221> misc_binding
<222> 7147..7171
<223> 12-809-119.probe

<220>
<221> misc_binding
<222> 16357..16381
<223> 12-805-115.probe

<220>
<221> misc_binding
<222> 26298..26322
<223> 12-790-396.probe

<220>
<221> misc_binding
<222> 31100..31124
<223> 12-791-211.probe

<220>
<221> misc_binding
<222> 33796..33820
<223> 12-803-125.probe

<220>
<221> misc_binding
<222> 34243..34267
<223> 99-33040-321.probe

<220>
<221> misc_binding
<222> 35281..35305
<223> 12-810-77.probe

<220>
<221> misc_binding
<222> 45880..45904
<223> 12-787-103.probe

<220>
<221> misc_binding
<222> 54485..54509
<223> 12-793-383.probe

<220>
<221> misc_binding
<222> 56570..56594
<223> 12-792-233.probe

<220>
<221> misc_binding
<222> 60324..60348
<223> 99-41009-244.probe

<220>
<221> misc_binding

<222> 60457..60481
 <223> 99-41009-111.probe

<220>
 <221> misc_binding
 <222> 79164..79188
 <223> 12-593-174.probe

<220>
 <221> misc_binding
 <222> 80004..80028
 <223> 12-589-152.probe

<220>
 <221> misc_binding
 <222> 87451..87475
 <223> 12-785-200.probe

<220>
 <221> misc_binding
 <222> 87644..87668
 <223> 12-785-393.probe

<220>
 <221> misc_binding
 <222> 101590..101614
 <223> 12-588-103.probe

<220>
 <221> misc_binding
 <222> 104379..104403
 <223> 12-603-191.probe

<220>
 <221> misc_binding
 <222> 117418..117442
 <223> 12-586-414.probe

<220>
 <221> misc_feature
 <222> 20352,102280
 <223> n=a, g, c or t

<400> 1
 ttggtagaat tggtagctaa aaggctgagt gaaatatggc ttaaaagctt tattggctgg 60
 gcgcggtggc tcaccctgt aatcccagca ctttgggtgg cttaggtggg tggataacct 120
 gaggtcagga gtttgagaac agctgaccaa catggtgaaa ccctgtctct actagacata 180
 caaaattagc caggtgtggt gatgcatgcc tgtaatccta gatacgtggc aggctgaggc 240
 aggagaatta cttgaaccga ggaggcggag gttgcagtga gctgagatcc taccattgca 300
 ctccagcctg ggcaacaaga gcaaaactcc gtctcaaaaa aaaaaaagct ttatcattta 360
 ttttttgccc ctgtcttatg gtgcagaggc ttaaaagtgt tttgacagca aattttctag 420
 aggctaggag tggtttattat aaccatgttt ttgagcgggt aggactacct cagagggcat 480
 gccttggtgc atcattgttc ttattgctga gctaccgaaa cctagaatct gactcacaca 540
 atatgacact tatttcctgt ttcttggtag agtttgtgtg gtcattcatc ttttagatct 600
 tttagaaact acagtcctcc tagttccac ttttatattt atttatttat ttatttttga 660
 gatggagtct tgctctgttg ccaggtgtg agtgacgtgg tgcaatctcg gctcactgca 720
 acctctgcct ccaggttca agcaattctc ccgctcagc ctccaagta gctgggatta 780
 caggtgcgtg ccaccatgcc tcgctaactt ttgtattttt tgtagagacg gggtttcacc 840
 acgttgccca ggctggtcgc aaactcctga cctcaagtga tcaaccccc tcagcctccc 900
 aaagtgcgtg gattacaggc gtgagtgacc gtgccagcc atcagttccc acttttaaat 960
 gagaggttct ttgtttttgt ggggtttttg ttgttgtgtg tggtgtgtgt gttgtgtgtg 1020
 ttgttacttg agacagggtc ttagtctgtc atccaggatg gagtgacgtg gcacaatctt 1080
 ggctaactgc agcctggacc tcctgggct caggtgatct tcctacctaa gccccctgag 1140

tagctgggag	tgcaggcaca	caccaccatg	cctagctaac	ttttgtattt	tttgtggaga	1200
tgagatttca	ccatgttgcc	caggctggtc	tcaaatccct	gtactccagc	gatcctcctg	1260
cctgagcttc	ccaaagtgt	gggattataa	gtgtgagtca	ccgagccagg	tcataaagac	1320
agtttttttt	tgttttgaga	tggagttttg	ctcttttgta	cccaggctgg	agtacagtgg	1380
tgtgatcttg	gctcactgca	acctccacct	ctcgggttca	agcaattctc	ctgtccttact	1440
tagcctccc	agtagctggg	attacaggcg	tgccccacca	cgctgcaact	atTTTTtata	1500
tttttaacag	agatgggggt	tcaccatgtt	ggacaggctg	gtctgcaact	cctgacctca	1560
ggtgatctgc	ccacctcagc	ctcccaaagt	actgggatta	cagatgtgag	ccactgtgcc	1620
tggccaataa	agagaatttt	aaacaatgcc	tacagtgtga	ttttagtggg	gtggccacat	1680
acatgtagg	gtgcacatgt	atttccatga	aaatgccttc	ctgccttttag	ccttgacaat	1740
gctgcatata	gggaataagc	atttaaataa	tggtgtctgg	gctaagagt	tgaactctgg	1800
agccaaaccc	tccaaaattg	catcctgaat	ctgccactag	ctagctctgg	agccatgaat	1860
aacttattta	acctctctgt	gatggtttat	ctatctgtaa	aatgggggta	atgaaacatc	1920
tggctcattg	agtgtctggg	agcatcaaat	aggtagtag	gtctgggtgt	gtggctcaca	1980
cctgtaatcc	cagcactttg	ggaagctgag	gcatgaggat	ggcttgagcc	caggagtgtg	2040
agaccagcct	gggcaacata	gtgagactcc	catctctaca	aaaaagaaaa	ataatagccg	2100
gtgtagtggc	atgaacctgt	ggtcccatct	actcagaggc	tgagggtggg	gatcactgga	2160
gcccaggagg	ttgagtctgt	agtgaattgt	ggttgtacca	ctgtactcca	gtctgggcct	2220
cagagttaga	tcctgtctca	ggaaaaaaa	agggttcgtg	acagaaagta	cttatggctc	2280
agagtaagt	ttctgtgaat	gttggtctgt	ttttaaatct	tgcatTTaat	tcctctttgg	2340
ttagaagtat	acttctttgc	ttggattacc	atTTgcatgt	ggagagatgt	tccttctaga	2400
accgtctctc	cttgtcacat	gattatgtac	agtgtgagct	ttattaatac	tctgtcagca	2460
tttgagggtg	gaacataaat	cggtggctgt	accagtgga	atccagcaag	agagagaact	2520
aagcaccgta	gcaactgttg	ctattttgtg	actgagcaac	agctgtgcga	gacagattta	2580
ttttttaatc	ctgatctcac	tgatagcctg	ggatgtggcc	ttgccacatc	tcaaaggagt	2640
agattgggtg	agatatccag	tctttgagga	gatttcatga	ccacatgcc	taaagattct	2700
gtttcatttc	aaatatgagg	agctaaagct	ttgtgtgtat	gagtgaagga	gaaagattaa	2760
gctttaaaag	attcatagtc	agtcctgaa	atatagtgtg	actggattaa	tgacatgata	2820
aatcactttt	aatgtgggtg	gtgagacttc	atgcatttca	gtcaaaaaag	taattttgct	2880
gtaatgtggg	tcatttccaa	atTTtgctg	ctactttacc	taggatggat	aaacattatc	2940
acagagtatt	tactgagcat	ctactttata	ccagggtctt	ttctgaatac	tggtaacatg	3000
cagttaatta	cactccaaaa	tgacttagtc	tgtgattcca	ctatatacta	cattacataa	3060
atacacacac	actaatcatt	tacatacttg	tcaaccttag	atttcagaaa	taacactgta	3120
tttgtaatga	atgcttcatt	accaaagtct	tttctgaagt	acatagtaag	atctttaaaa	3180
agagctttca	tggccaggcg	cggtggctca	cgcttgtaat	cccagcaact	tgggaggccg	3240
agggcgggcg	atcacaaggt	caggagatca	agaccatcct	ggctaacacg	gtgaaacccc	3300
gtctctacta	aaaatacaaa	aaaattagct	gggcgtgggtg	gcgggcgcct	gtagtcttag	3360
ctactcagga	ggctgaggca	ggagaatggc	gtgaacctgg	gagggtggagc	ttgcagtga	3420
ccgagatcgt	gccactcccc	ttcagctagg	gcgacagagg	gagactctgt	ctcaaaaaaa	3480
aaaaaaaaaa	aaaaaaaggc	gctttcaaat	agtgggttga	ttcattcaat	aactttctaa	3540
ttttcatttc	tctttttttt	cttgagatga	agtcttactc	tgtcaccocag	gctggagtgc	3600
agtgcattaa	tatcagctca	gtgcaacctc	tgccctcccg	gttcaagtga	ttctcctgcc	3660
tcaacctccc	cagtagctgg	gattataggc	acacaccact	atTTctggct	aatttttgtg	3720
tttttggtag	agatgggggt	tcaccatgtt	gcggaggcta	gtctcgaatt	ctcgacctca	3780
agtatacac	ctgccttggc	ctcccaaagt	gctgggatta	caagtgggag	ccaccgcgcc	3840
cagctccatt	tctgatgtgt	tccaatatTT	cagggtctta	cggtgcctt	gaaatagtaa	3900
tgtgggccag	gtgcagtggc	tcacgcctgt	aatcccagca	gttagggagg	cggagatgcg	3960
cagatcacct	gaggtcagga	gttcgagacc	agcctggcca	acatgggtga	accctgtctc	4020
tactaaaaat	aaaaaaatta	gccagtgtga	gtgggtggatg	ccagtaatcc	cagctactca	4080
ggaggctgag	gcgggagaat	ttcttgaacc	cggttgagg	tggttgcaat	gagccgagat	4140
cacaccactg	cactccagct	tggggaatag	agcaagactc	tgtcttcaaa	aagaaaagaa	4200
atagaaatgt	cgttcatttc	caaatttttg	ctggtaactt	acctaaggag	gataaacatt	4260
gtcacagaat	atttactgag	catctacttt	ataccagggt	ctgttctgaa	tactggtaac	4320
acgcagttaa	tgtgttgggt	ctgggttttc	atgttccatg	tataaattac	ctcttaaaac	4380
tatgattgtt	ttcattcatt	cattcatttt	tgagatcggt	tctcaaaata	ggctgggggtg	4440
cagtggctcg	ataacagcta	actacatcct	caaacctcca	ggctcaagcc	atcctccgac	4500
ctcagcttcc	taagtagctg	ggactgcagg	catgtgccac	catgccacgc	tagttgttgt	4560
tgtttttttt	ttttttggta	tttgttggat	agacagggtt	tcacctgtt	gcccagggtg	4620
gtcttgtaact	cctgacctct	agtgatccac	ccagctcgga	ctctcaaaat	gctgaaattta	4680
caggcatgag	ccactgagcc	cagcctatta	atTTttgatt	gccctattca	gatttagtgg	4740
tatgaattta	ctgattggga	tgTTTTgaac	acacagaata	acatgtttta	ctaaagaact	4800
tgtaattgag	tgacttataa	aatgaaacat	ttttatcttc	taggtattat	taacccaaag	4860
aatccaaagg	aagctccaaa	gtccttcagc	ttcgactatt	cctactgggtc	tcataacctca	4920

gtgagtaccc	tcatgccaca	gcactgccag	ctcctgcctc	ctttcctctt	tccttgccga	4980
tttgtctttt	ctcttagctg	agcagatcca	agttttgctt	cttaactggt	gtgtagcatg	5040
agttcattct	ttgatacagt	gatatagtat	tttcattcat	aaagtatttg	ggttgatgga	5100
aagttttatg	acacatatat	ataaatatat	acacatatat	acatacagaa	catatttagt	5160
ttacaccaaa	tttatactga	atgtacacta	aattttatact	gatttagaca	tttcttgagg	5220
atttatttat	cagtatctta	ggatcctcat	taccaacatt	attgcccatt	gatcttagga	5280
aatgccaaat	aataataatt	attattatta	ttattatttt	cttctgagac	ggagtctcac	5340
tctgtcgccc	aggctggagt	acagtgggtc	catcttggtc	cactgcaacc	tccgctcgag	5400
gttcaagcga	ttctcccggc	tcacctcccg	agtagctggg	attacaggcg	cgtggcacta	5460
tgcccagcta	atTTTTgtat	tttttagtaag	acaggatttg	accatgtggc	cagctggtct	5520
cgagcttctg	accttagtga	tccacctgtg	tttgcctccc	aaagtgtctg	gattacagggt	5580
gtgagccact	gtgcctggct	ggaaatgcc	aattaaaatt	gcttagggta	ctgttaattt	5640
taagttgaat	tttgaggga	tgggcagcaa	gattggaatg	accacagaga	aattgttttt	5700
tttttgtttg	tttgtttgtt	tttggctagg	ctgatcttga	actcctgacc	tcctgatcca	5760
cctgcctcga	cctctcaaag	tgttgggatt	aaaggagtga	gccaccgtgc	ctggccaatt	5820
tttttttttt	aattgtagta	gagcaatgtg	ggttctgaac	atTTTgaaag	atgataccgt	5880
gcctctacta	aaaattccaa	aaaaaaaaaa	aattagggtg	gcttgggtgg	gggtgcctgt	5940
aatcccagct	acttgggagg	ctgagggtca	tgacctcgcc	gtaacccgat	ggcgtcccac	6000
actgcactcc	agcctgggtg	atggaacatg	actgtgtctc	aaaaaaaaaa	aaaggaaaga	6060
ggctgcagat	ccacaagaaa	ctaaaagatt	accagtatta	cgaaaatgca	gagtagggcc	6120
cttctgggta	cccttactca	gtgtggaacc	ccaaatgttg	ctgcttctta	gggattgttc	6180
atagagttta	cttctgcatc	attcatctct	ttgttacatg	tttttggttt	tgttttctct	6240
gttttgtatt	tttttttaaa	gactgagtct	cgctctgttg	cctaggctgg	agtgcagtgg	6300
catgatctca	gctcactgca	acctacacct	cctgggttca	agcagttctc	ctgcttcagc	6360
ctcctgagta	tctgagatta	caggcagggtg	ccaccacacc	cgctgatttt	ttttaatttt	6420
tagtagagat	ggggtttcgc	tgtgttggcc	aggtgtgtct	cgaattccta	tcctcaagtg	6480
atctgcccg	cttggcctcc	cagagtgatg	ggattacagg	catgagccac	tgtgccctgc	6540
cctggggaat	cgatgtttta	aactattgac	aaagccta	tgtcaaggga	agggactgag	6600
aaggggattg	gaaatgctgt	ttcctgaaag	ctgtgtttat	tgagaaatag	tcccagtagc	6660
tggagtga	tcaaagaagt	atTTTTatat	gtggacttga	cctttggtct	tttattctca	6720
ttttccactt	aagaaaatct	tggctgtgtg	aatgaaagag	tgatactttt	aaggttatag	6780
aaagtgaat	gtaatcatgc	cagataattt	tatatagata	tttttatgta	tggctgacct	6840
ggatgactct	aacagtgc	gtgtttgtga	gtgtgtgtgt	gtgcatgtgc	gtgtatttaa	6900
tgagaaaagt	aaacttgtgt	ataggaggct	taaaaaatgt	gtagggaatt	ttaggtgact	6960
gttctgattc	cagacacttt	tattatgaa	gcaactaagt	aagtatagga	agaaatatta	7020
ataaaagggt	atTTTattct	ccttttactc	tttacagccc	gaagatccct	gttttgcctc	7080
tcaaaaccgt	gtgtacaatg	acattggcaa	ggaaatgctc	ttacacgcct	ttgagggata	7140
taatgtctgt	atTTTTgcst	atgggcagac	tgggtgctga	aaatcttata	caatgatggg	7200
taaacaagaa	gaaagccagg	ctggcatcat	tccacagggtg	aaaaacaaaa	caaaacaaaa	7260
atcttctctt	cattattagt	gttagtctta	aattgtctta	acagttattt	ttatttggcg	7320
aacatttatg	cggggattgt	tttatgtcag	gcacaaagat	gaacaaccca	ttattttccc	7380
tcagaggga	tcacaattga	atgggaagga	ttgacatgta	cacatgtctg	tcattaaagg	7440
tgggaaagtc	agtgttttgt	aatgattttg	ccatatatcc	aatgccatat	tattttgtca	7500
tttgaaaagt	gttaccagct	tgttaaagct	ctgttctaag	tcctggagat	ggggggatat	7560
tgttgatctg	atTTTTtttc	aaattccatg	catagattac	ctctgaagaa	tgtgattatt	7620
tttggttttt	ttgatagcct	tattcggaga	tactttcatt	ttattttcct	aatttaatac	7680
tatgacttta	aacttcaaac	actctgagat	tctccctctt	tttttttttt	tttagaacag	7740
ttagtattat	tgggtttgat	cctttattgt	ttgggaatga	agaagcgttc	tgacatagat	7800
atttatttat	ttttattttc	attattcgtt	tcacttagga	tttaattggtg	aaagacataa	7860
gtatttaaaa	agaacttcat	tatttactta	ttattttgag	acagggtctc	actctgttgc	7920
ccatgtctga	gtgcagtggg	gtgatcttgg	ctcactgagg	ccttgtcctc	ccaggctcaa	7980
gtgatcctcc	tgcttagacc	tccagagtag	ctgggactaa	agccatgtgc	caccacacct	8040
agctcatttt	taaattaaaa	aaaaaaaaaa	tttttttttt	tttttgagg	ggagtctctc	8100
tctgttggcc	aggctggagt	gcaatggcgc	catcttggct	cactgcaacc	tccgcctcct	8160
gggttcaagt	gatttctctg	cctcagcctc	ccaagttagct	gggacaatag	gcacccacca	8220
ctatggcctg	ctaatttttg	tatttttagt	agaggcgagg	tttcaccatg	ttggctcctg	8280
tctggatctc	ctgacctcag	gcaatccgcc	cacctccgcc	tcccaaagtg	cagggaattgc	8340
aggcatgagt	caccatgccc	ggcctttaaa	atTTTttttt	tagagacagg	gtctcaccat	8400
gttgtccagg	ctgggtctga	atggacctgg	gctccatcct	gggcttggga	tgatctgcct	8460
gcctcgactt	ccagaagtac	tgggattaca	ggcatgagcc	actgtgcggg	gcctaaaaac	8520
ttaagaaaaa	aaaaagatta	agtgagaaga	ttgtatttaa	aattttcctt	ttatagtcac	8580
gcatacctta	agggtggggg	tacattttga	gaaatgcac	attagtcgtt	agggtccttt	8640
gtcattgtac	agatatcata	gagtgtactt	acacaaaccc	tggaagggtg	aaactgctcc	8700

acacctagac	tgcgttgtat	agcctgttgc	tccaaggcta	taaacttgtg	tagcatgcta	8760
ctgtactgaa	tactacaggc	aattataata	ggatggtaag	tatttgtata	tctaaacata	8820
gaaaagtaca	ttaaaaatat	gatataaaag	atttgaaatg	gcacacctgc	atagtgtctc	8880
tacactggag	cttgcagggc	tggaaagtgc	tctgggtgag	tcagtgagca	agtgggtgagt	8940
aaatgtgaag	gcctaggaca	ttgctttgca	gtattgtaga	ctttataaat	gctgcacact	9000
taggctacac	taaatttagt	tttaaaaaa	gtttcccttc	aataattaac	cttagttaaa	9060
ttactataac	ttttttactt	tataaacttt	aaaatttttt	aaactttctg	atccttttat	9120
aataaccttt	agttaaaac	acaaacagcc	ccagcatggt	ggctcatgcc	tgtaatccca	9180
tcactttggg	aggccaaggc	gggtggatca	ctggaggcca	ggagtccgag	accagcctga	9240
ccaacatggt	gaaaccccg	gtctactaaa	aatacaaaaa	aatgagccag	gcatggtggc	9300
gcatgcctgt	aatcccagct	acttgggagg	ctgaggcagg	agaattgctt	gaacccaggga	9360
gacggaggtt	gcagtggagt	gagatcgcg	cattgcactc	tagcatgagc	aacaagagt	9420
aactcggctc	caaaaaaaa	aaaaagaaac	cacacacaca	ctttgtacag	ctgtacaaaa	9480
atattttctt	tcttcatatc	tttattctat	aagctttttt	gtatttttta	attttttttt	9540
tctttcttga	gacagagtct	tgctttgtca	cccaggctgg	agtacagtgg	cgccttatca	9600
gctcactgca	acctccgctt	cctgggttca	attaattctt	gtgccttagc	ctcctgagta	9660
gctgggacta	caggcacgta	ccaccacgtc	tggctaattt	ttgtattttt	agtagagacg	9720
gggtttcaca	atgttggcca	ggctgggtct	gaactcctgg	cctcaagtga	tctgcctgcc	9780
ttggcctccc	aaagtgctga	gatttacagg	cgtgagccac	tgacacctgc	caaatttttt	9840
tttttttttt	ttagacagag	tttcaactct	cttgccagg	ctggaatgca	gtggcacgat	9900
cttggctcac	tgcaacctcc	gcctcctggg	ttcaagcgat	tctcctgcct	cagcctccag	9960
aatagctggg	actatttttg	tatttttagt	agagatgggg	tttcaccatg	ttggccaaga	10020
tgttctcgat	ctcctgacct	tgtgatctgc	ctgcctcgtc	ctcccaaagt	gctgggatta	10080
caggcatgag	ccaccttgcc	cagcccaaat	ttttaatttt	tgaaaaaaaa	tttaaacttt	10140
tttgttaaaa	acttaggcac	acattttatta	gcctaggcct	acacagggtc	aggatcatca	10200
agatatcaact	aggtagacag	aattttccag	ctttcttata	atcttatggg	gatcactatc	10260
atatatgcag	tccatcggtg	accgaatcat	ggttatgcag	cgcatgactg	actttattca	10320
aagcattaat	cgtatcttga	tgtttatgac	ataatatatt	ttgagatgga	gcaagaaagg	10380
acgtctggct	aattcattga	gcacgcgtgg	atgacttagt	actcctctca	tttgtgtctc	10440
tcatgcctct	ctcattctac	ttccctagtt	atgtgaagaa	ctttttgaga	aaatcaatga	10500
caactgtaat	gaagaaatgt	cttactctgt	agagggtgag	acagccgtga	gttgacaccg	10560
taagcccttg	ttttccattc	tctcaagcat	cacttaaatg	gctccaaatt	atgactgtgg	10620
tacacatcac	ttcaccattc	cttcattttt	gtcctttcag	gctatttctg	ggttttgggt	10680
ataatgaata	gtgtcctgct	ttaatctttt	ttttgggggg	tgcgcggtgg	ggacagggtc	10740
tcactctggt	gcccaggccc	tggagtgcag	tggcgtgatc	acagctcact	gcagccttga	10800
ccttctgggc	caaagcgatc	ttaccccttc	agcttctcaa	gtagctggga	caacagggtg	10860
gtgccaccac	cacccttaga	taattgattt	ttttgtagag	atggagtttc	cctatgttgc	10920
ccaggttggg	gttgaactct	tgggctcaag	caatcttgcc	tccctccacct	ctgaaagtgc	10980
tggggttgca	gtgtgaatca	ctgagcccag	ccacaattaa	atctttgttc	ataagtctgt	11040
gcatttattt	gcagttatta	tcttttgtat	gttctttggt	ttttaaataa	tacatttgaa	11100
ttgccaaatg	ccttttttga	aagtaatgct	gggtaacagt	tttcagtcac	actaggagg	11160
tgtaagcacc	cattgtgccc	ttgccaatat	ttttctattt	tttaatcttt	gctaatttga	11220
taatgtaagt	attctcttac	ttaaattttg	aatttctttg	attactagta	tgggtgaacg	11280
tcttgtaaat	tatgcatatg	cctgtttaca	ttttgggttc	acaatgtctt	tctgattcat	11340
ttgtgggaac	tcttctcctt	taatatgttt	aataatggtt	tttgacaggc	attgggttta	11400
aatatatata	atcaaatctc	cccccttccc	acttttttaa	ttgcttataa	aaacttggaa	11460
agtaattatc	catcctgtgt	ttaccctggc	tctttcatga	agtctataga	attactttat	11520
acacatttgc	ttatttgata	aacattctct	ctgttttgta	tgtgcttaaa	gaagctgttt	11580
tggggctcat	acctgtaatc	ccagcacttc	gggaggcgaa	tggatcactt	gaggccaggga	11640
gtttgagacc	agcgtggcca	acctggcaaa	accctgtctc	tactaaaaac	acaaaaaatt	11700
agccggctgt	ggtggtgcac	acctgtaatc	ccagctactc	gggaggctga	ggcacgagaa	11760
tcgcttgaac	ccaggaggca	gaggttacag	tgagccgaga	tcatgccagt	gcacccagc	11820
ctggagtctg	tctcaaaaaa	aaaataataa	taataaataa	ataaaaagaa	gaagctgctg	11880
tttctttctc	ttcctccctc	tctctctcac	cgccaccccc	ccaccacacc	cttttttggg	11940
attcactttc	atagtaactg	aaattattaa	agtgcttctg	taccatgagt	ctgttttgaa	12000
aaaataggct	gccttttaag	cgatcttgag	attttttttc	ttttcttttt	ggtatttggg	12060
ctgattctct	gtagccttac	agcatttcat	tcctagttaa	ttcatttaat	tgctttggag	12120
ttttttcttc	acctagaagg	tggtgcataa	atatgtaaat	tctaaggaat	tctgtcttgt	12180
gactgtcata	cacaaaaaag	agaatgtgct	gccctaagca	tattttgcgt	taaaatgaat	12240
ttctgaata	gcttgtccca	tagttattct	tttttgaatc	aggtgataga	atttttgagc	12300
ttacaggtaa	taaaagaaga	tgcccttttc	ttttgtatta	gagtttccag	gcttaacttg	12360
gagtgaggca	tagattccta	gttgacactg	tgtttctctg	ccacatttgc	aggacctctc	12420
tacctgtctc	ttctacttcc	ttccctgttt	actctgctcc	acccatattg	ccttttttcc	12480

tgtttcaaga	acataccaga	aacatttcctg	cttcaaggcc	tctacattta	cttctcctct	12540
tgctgaaat	gttggtcact	tggtatgcat	tttgactaag	tttctcacgt	tcttcaagtc	12600
tgataaaatc	acacactctc	agtggagctt	accttgacca	ccttatttta	tattgtatct	12660
tgccctttac	cccacatcct	gagctctctt	ccccattcta	ttttcttttt	tccatcatac	12720
tcatcatgtt	ctaacaatgct	ttctcagtta	cctatttttt	atgtttgttg	tttattatca	12780
gtatcccttg	ctagaagcat	aagctcactg	gggcagggtt	ctttgtctgc	tttatttagt	12840
ggtgtatacc	aattgcctag	aacagtgcct	gtaagagaac	ggctcctcagt	gagttggatc	12900
tgccagggtg	catctggagt	ggttggtgca	gaagtaaaag	aatgatgat	ggctttggat	12960
ggattcacat	atcagagcat	aaggaatgga	gcacatgaac	ttttacctag	attatttggg	13020
ttctctttgt	ctcacatttg	tgtgaaaagg	tacgagtgtt	ccaaaagatg	taatatattc	13080
cggggattga	gtattgtatg	tacaaattta	agtacacagc	aatgattgaa	gaatcatctc	13140
actttctctg	atgacttgct	gtcagaaaaga	atattttgtg	ttctgttact	ttactctcat	13200
aatagattaa	ttactttatc	tttcagagtt	gtcttttttt	ttggagacag	agtctcactc	13260
tgtcacccag	gctggagtgc	agcggcgctg	tctcggtcca	cttcaacctc	cgctcccgag	13320
gtcaagcaa	ttctcctgcc	tcagcctccc	gagaggctgg	gattacaggc	tcctgctacc	13380
acacctggct	aattttctgt	attttttagta	gacacagggt	ttcaccatgt	tgccagggt	13440
ggtctogaac	tcctgacctc	atgatccacc	cgcttggtgc	tcccaaaggg	ctgggattat	13500
aggcgtgagc	cactgcgccc	ggcctttttt	tctttttctt	cttttttttt	aagacaagggt	13560
cttgctctgt	taccagact	ggagtgcagt	ggtgcaaaaca	gggttcgctg	cagccttgac	13620
ctcctgggct	cagtgatcgt	cctgcctcag	cctcctgagt	aatttgacc	acaggcatgt	13680
gccacagtgt	gtggctccca	aagtgtctaa	attacaggca	tgagctactg	catctggcca	13740
agttcttaat	tttttaaaat	ggattttatt	tattttattta	tttattttatt	tattttattta	13800
tttattttatt	tttagagaca	ggatctcact	gtcaccagg	ctggagtgc	gtggcgtgat	13860
ctcagctcac	tgcatcccca	acctcctggg	cttgagtgat	cttccacact	cagcctcccg	13920
agtagctggg	aacacaggca	cacactacca	tgcttggtca	attcctttgta	tttttggtag	13980
aagtggggtt	ttaccatggt	accaggttg	gtctcgaatt	cctgagctca	agcaatctgc	14040
ccaccttgac	ctcccaaaga	gtggggattc	caggcatgag	ccaccatgcc	cagcctttaa	14100
ctgattttta	aacacaaatt	aaggccagggt	gcggtggctc	acacctgtaa	tccttgact	14160
ttgggaggcc	aagatgggag	gattgcttga	gccaggagt	tcaagaccag	cctgggcaac	14220
gtggtgagac	cctgtctctg	caaaaaaaaa	ttttttttta	ttagctaggc	atggtggcac	14280
gtgcctgtag	tcccagttac	tccaggagct	gaggcaggag	gattgcttaa	gccagggaag	14340
tcgaggctgc	agttagccat	gtttgagcca	ctgcactcca	gcctgggtga	cagagtaaga	14400
tacagtctca	aaaaatatat	aaaaataaaa	tacagtataa	aaatacagtt	agagactgaa	14460
tgaagaaaaa	aggcaataga	aatggaatta	agattccatt	ccagatactc	tgaaaacatg	14520
tccatagaca	tttaaagttt	aaaaatgcag	gagctgtaat	gtaagcttga	atagctccat	14580
gatacagaag	gatctgtttc	ctgggtagca	tcgtaatgtg	gcctattgtg	atgaatttac	14640
cctagttcag	agccttggtt	gcttagaagt	ctgattttgt	agattcatca	agaaaaggac	14700
tatccttaca	agttaagact	tgagtaaaag	tcatttgact	gatgttatga	aaccaacttg	14760
ttggtttgtt	actttaccat	gagaaataac	tattactaag	gctagaacac	tcaagccatt	14820
cgcaagtacta	tttaatatat	ttctgtggac	agacctcgct	gttggaagcc	tgttggttgt	14880
ttggtgtgca	tggaattaga	aggagaggag	ttgcttagat	acatgtttta	ttgccccagg	14940
ttaagaaatg	atagcttaat	gcattaatca	ccatgatcga	atggctcctg	caatgagaca	15000
agaattttatg	aagtgtattg	tctgaggctt	ctttctccct	cacaaatctt	ggataggtcc	15060
tgccaccacc	agctttcatg	ctggtcaacc	ctcatccctt	ctggaggata	catcgtccat	15120
agttagtaac	agtgtggtga	gccttaaaaga	attagcatca	ccgctctgtt	ttgtcttctc	15180
tgctctgtaa	tattgatatt	gttccttggt	tttaactcta	atagaaatct	tatttccgag	15240
ttcttgaggt	aacctcaatg	caaatcccag	ttactgacaa	cagtatacat	tttaaaaaac	15300
attgataact	gataaattgc	tactaaaata	tttttagct	gtcactacat	tccagagata	15360
atctgtttac	ctctccctta	cagagaaaag	ctgaaggctg	gttctaacct	attcagtgt	15420
attacctgtg	cgtattgtta	atagaataaa	agggcctaga	gttacggcat	ttcactttgt	15480
aaaggttgct	gatctctaatt	cttagagctt	tttaataaaag	ttaaaaaaat	tttttttctg	15540
tatgtgattt	cttttggttt	ctttttcttt	ctttctttct	ttcttttttt	ttattaagac	15600
agagcctcac	tctgtcaccg	aggctgaagt	gcaggggtgt	gatatcggt	cactctaacc	15660
tccacctccc	aggtacaggc	gatttctcct	gccttagcct	cctgagtagc	tgagactaca	15720
ggcatgtgcc	accacgcctg	gctccttttt	gtatttttgg	tagagacggg	atttcaccat	15780
gttggccaga	ctggtctcga	actcgtagtt	cgagtgtatc	caagtgtacc	gccctccttg	15840
gcctcccaaa	gtgctgggat	tacaggcgtg	agccaccgct	cccggcttct	gtatgtgatt	15900
tctttttcac	tctaattcac	tttactaatt	tgttcatagg	tgagctacat	ggaaatttac	15960
tgtgaaagag	tacgagattt	gctgaatcca	aaaaacaagg	gtaatttgcg	tgtgcgtgaa	16020
caccacttcc	ttggacccta	tgtggaggat	ctgtccaagt	tggcagttac	ttcctacaca	16080
gacattgctg	acctcatgga	tgctgggaac	aaagccagggt	atggtaggaa	atagagtaat	16140
gactgaggtc	tttggcacct	ttttaggtcc	ttttttccca	gttaagggtt	tgaggccaca	16200
tttatagcta	tgaaagtgtc	tttaattgtg	gagtcctctg	atcctttgac	tgtgctgtaa	16260

cagtgagggc	ttcagagtgt	taagtatctt	ctcatctaaa	cagtacattg	ctgaaacatg	16320
tttagagctct	cattgagttt	tactcttagg	ttttgatcaa	gtcatggtra	aatgttagtt	16380
atcatgtata	tatgcggctg	tggtagtaat	tgattctgga	ataaatgagc	accaataaag	16440
aataaaaaatt	aaatgcttca	ttggtttcta	aagggatagc	tgaggtatga	aaaatgattc	16500
tgatgggggtg	cctttggaat	tgagagtaa	cttgatgggc	tcacaaataa	ggcagttattg	16560
agagcaattg	tgtaagtac	aatcttgcaa	tctgtgaata	agtttaagac	tattgtctgt	16620
gattgttatg	cttataatgt	ggagcctgct	gtccatttca	actctcatct	aagaattccc	16680
ttgtcacgtg	gtcaccttta	tgtttattta	ggacagtggc	agctacaaac	atgaatgaaa	16740
caagtagccg	ttcccacgct	gtgtttacga	ttgttttcac	ccagaagaaa	cacgataatg	16800
agaccaacct	ttccactgag	aaggtaggag	agtttcagtc	tctaggcttg	agttgtgaag	16860
gatggagatt	ttgagaagtc	ccttttggtg	ccctctgctt	tttgtggaag	gttgggtggt	16920
gggggggtgct	cctttactta	atggagggtg	ttttatacct	ctgcttatca	tttttcttcc	16980
accctacacc	agttttctgt	ataggttctt	aaatgaagtt	gtaagataga	gaatagggtt	17040
aagaaaaaaa	caatccagga	ccttgggtct	caaaaactac	tcaatacaag	acagttaagc	17100
agtcttgacg	gagaaagttc	tttcttttct	tcttctgac	tgaacctttt	tattacctta	17160
ttagctgcct	ttggtatgtg	tattcttttt	tttttttttt	ttgagctatg	ataatgttat	17220
agtgggatta	aatgtttaag	caaatgatat	gtttcattcc	tggaataata	ctccacaaca	17280
gttctgggtg	tcttcaaata	cctttctttg	cctattgatt	tttcattttc	tatagtgggt	17340
aaatgtcaca	gacttttagcg	ttggattgct	tgctttgaat	tctgattctg	ccactcttca	17400
gctatattac	tttgggcaaa	ctacttaacc	tctttgttcc	tcagtttccc	catccataac	17460
agcctgatag	gattgttaaa	aagattaagc	gttaatatag	gttaagtga	tacaataatg	17520
tctggcacag	agtagccact	caataatggt	aattttattg	ttattgttag	taataaaaaa	17580
tttatgttga	atactactct	atctgcatcc	ctaagaagga	tgacacctga	tttttaagct	17640
tcaagttagg	tttattgtac	tggaattat	aaaagatcct	ttgccttata	aaatgagtac	17700
tttagacacc	actgtcacta	attattaaaa	tcacctttta	tttgttgact	tatttctagc	17760
tgaaaaatgat	ttttaaaaa	atactcctgc	tgggccgggt	atgggtggctc	acacctgtaa	17820
tcccagcact	gtgggaggcc	aaggcagaca	gatcaagagg	tcaggagttt	gggaccagcc	17880
tgccagcat	ggtgaaaccc	agtctctact	aaaaatataa	aaaaattggc	tgtgcgtggt	17940
ggcatgccct	gtggtccac	tgctcagagg	ctaggcagga	gaatcgcttg	aaccagagg	18000
tgagggttca	gtgagccaag	atcgtgccat	tgactccag	catgggtgac	agagcaactc	18060
tgtctcaaaa	aaaaaaaaaa	acacactcct	actgggcattg	gtggctcaca	cctgtaatcc	18120
aagcaacttg	ggaggctgag	gcgggtggat	cacctgaggt	caggagttcg	agaccagcct	18180
ggtcaactgat	gtgaaaccca	gtctctacta	aaaatacaaa	aaattagccg	gtcatgtgtg	18240
gtgccggtaa	tcctagctac	tccagaggct	gaggcaggag	aatcgcttga	acctgggagg	18300
cagaggttgc	agttggctga	ggttgcgcca	ttgcacttca	gcctgggtaa	caagagcgaa	18360
acttcgtctc	aaaaaatata	tatatatttc	ttatatagat	gatcaaataa	cttgttttagc	18420
cagaggttat	aatattagtg	ctatttatta	gttagtgttt	atctgaatgc	cagcaaagaa	18480
cattgtatgt	gaaggtgttt	tttgccttc	tcccatgatg	tcgaatctac	aaaattaatg	18540
ttaatcccat	aatcatagct	ttccttgtaa	ttcatttttg	atacttttgt	ctaaactttc	18600
ctgaagcttt	atttttttac	agtgttattg	aggtattatt	gaagtataca	gtaaattggca	18660
catattttaa	gcgtaaaatt	tggtcagttt	tgacgtattt	gtataacctgt	gaaactagcg	18720
ccacaatcca	gataatgata	atttttattc	tcaaaagttg	cctcatatct	cctttagtgc	18780
tttctctatc	ccttcttctg	tggaattcc	cgggcaattc	cggacctgcc	cctttgtcac	18840
tatatattaa	tttacatttt	ctagaatttt	atataaatgg	aatcatagag	tatgtactgt	18900
tttttgtctg	gagtctttca	cttagcatag	tgattttgag	attcatctat	gttgttgtgt	18960
ttatcaatag	ttccttttca	ttgttgatga	gtattgcgtt	atacagatac	accacagttt	19020
attaattcaa	ctgttgatgg	acttttaaac	gtttgccagt	ttttgattat	tacaaacgaa	19080
gctgatgtga	acatttgtgt	acaagtcttt	gtgtggacat	ttgtttttgt	tctgttggta	19140
aatacttagg	agtaggatga	ttgagtctta	ttgtagctcc	ttaagaaatg	tcaaggctgg	19200
gcacggtggc	tcacgcctgt	actcccagca	ccatgggagg	ctgaggcggg	tggtatcaca	19260
ggtcaggagt	ttgagaccag	cctggtcaac	atggtgaaac	cccgtctcta	ctaaaaatac	19320
aaaaaaaaatt	agctgggcgt	ggtggtgggc	gcctgtaagc	tcagctactc	gggaggctga	19380
ggtgggagaa	tcgtttgaac	ccaggagggtg	gaggttgacg	tgagccaaga	ttgcaccatt	19440
gcattccagc	ctgggtgaca	gagcgagact	ctgtctcaaa	aaaaaaaaaa	aaaggaaaga	19500
aaataaatgt	caaattgttt	tccaaagtga	tgttaccatt	ttacattccc	accagaaata	19560
tgtgattgtt	ccagttgctt	aaagcttcac	ttttagtaag	ccatttaatg	agttgatgag	19620
ttttatata	ttttataaca	tactctagta	aattatgtca	tgtttatcct	aaagttactt	19680
ttttttgaac	cgtgataatt	gctatgaatt	ttgtgcatgc	tctttttttt	tttaattaaa	19740
aaaaattttt	ttttagagac	acaagggtgt	tatatgttgc	caggctggag	tgacgagac	19800
agctttgaac	tcctgggctc	aaaggagctt	cctgagtagc	tggtgactaca	ggcatgtgcc	19860
attgcacctg	ccgtgtgtac	tgtttaaaact	aactgataga	tggtgtcttg	atctttttgt	19920
gcatatttta	aaagctctaa	tacttttagt	ttttccata	ccttaattctt	ttaaaaagca	19980
ttctgcctct	aaatatttcc	tgcttctatc	ttgcttattt	ttgaattgcc	cccatgttat	20040

tggtctttat	caggtcagta	aaatcagctt	gggtgatcta	gcaggaagtg	aacgagctga	20100
ttcaactggg	gccaaaggga	ctcgattaaa	gggtatttatt	ttagcaaata	aatggcctga	20160
tcaataaatg	gccactacct	gtttttgtaa	ataaaatttt	attgaaacac	agctacatac	20220
atttgtttat	gtattgtcta	tggctgcttt	tgcactacaa	tggcagtggt	gtgacagaga	20280
ccgtatggcc	cagaaaaacct	aaaataattt	gtgtctgtcc	ttctacggaa	aaattttgct	20340
gactcctgat	cngtaacttc	aatggaagga	taagttatct	aactaaatag	ttagggagca	20400
catcatagtc	aggggtgagag	ttgaattgca	ttttaaggaa	caggtagaat	ttggctagat	20460
ggaattaaga	gatggcagtg	ggaataggat	atacaagatt	cagatgtgag	aacagaaatt	20520
tcattgtcgat	aagttgaggg	aagaaggctg	cacacatctg	taagcttttg	gagcacaat	20580
atctcgaaga	tacttcagga	ctgtgagtta	ggcatgtggg	ctcagcatgt	tattatcaag	20640
tggaaaagag	gacagatgga	ggactaaagg	ttgttttttag	aacttgtaat	aaagaaatat	20700
gctagcatat	ggcaaatcat	attttatgtt	ctgtatgata	tagcagtagt	agaaattctt	20760
catcattctt	tcataatttg	tattttattt	aggaaggagc	aaatattaat	aagtctctta	20820
caactttggg	caaagtcatt	tcagccttgg	ccgaggtggt	aagttttata	cttcattata	20880
aggggagttg	tgtctgttca	gcaaataatac	ttgatgatat	tttattattg	tttataggct	20940
gtctttttgt	gcttttctgt	aatctgcctt	tctcctttta	gagcttttca	gataagatac	21000
taggtataac	acagtttgac	tttgactga	aagtaaaaga	tactaaggag	aacaaaagta	21060
aatatagatc	tgtaaaaact	agagcaaaaca	ctgaagggtt	gactcatttt	tttgagcctt	21120
attttatggt	tctcacaaaa	accaagtata	tttattgtct	cttcttttcc	tgtataatct	21180
ttctctagta	agatgcagta	tcttttgag	tattttgagc	agtttggtt	tgggtgaatgc	21240
tgttgatttg	agggccagta	gtttttttt	tttttttccc	ccaatagttt	tgttctttcc	21300
tactccttta	tggtagaaca	gtaaaaactaa	aactattttt	tttaaaattag	gggctttggt	21360
gctacattat	ttaaattagt	cattgtgtgt	tctagtttag	tttaaaacct	ctttttaatt	21420
gctaaaaaaa	tgaatgctt	tcttgcttgc	cttgagaaaa	tcataatcta	tgacatgaaa	21480
tttaaacaaa	atctaatttt	ctacttgag	gcttatcctg	tgttcttatt	ttctccttta	21540
aatgctttca	cctgtaggat	aactgcacta	gcaagggtaca	gtgggggattg	gtagagataa	21600
actagaattg	acttttatgt	tttaaatcct	cactaggatg	tatggaggca	taagtaggaa	21660
tggaaacctc	aaaaatcttt	tcattcatttg	tctcttggt	ctggaattta	gagtggctgt	21720
aaatttaggg	tgaccacaaa	tcattgaggtt	tcacacccaa	attaatttct	agaacatgat	21780
atgctttgga	gaaggaatta	agaataaagg	aacaaggccg	agcatggttg	ctcatgcctg	21840
taatcccagc	actttgggag	gccaaggcgg	gcgatcatt	tgaggtcaga	agtttgagac	21900
cagtctggcc	aacatggtga	aactctgtct	ctactaaaaa	tacaaaaatt	agctgggtgt	21960
ggtagcgcgt	gcccatagtg	tcccaactat	tcacgggttt	taaactgaag	ttttaggcct	22020
caaggattac	atagtggagt	gaaagtttgt	tccactaatc	tgaatgactg	aaaggcttaa	22080
ttctcccttg	aggtctgtgc	aatctaaaaa	tgaacctgtc	attggaaatg	actttctgat	22140
tgggtgattg	tgtgtctgtg	ggtagagtgg	aatgagatga	tgatgatgtg	gacataattta	22200
ataagagatg	gaattaaaat	gtttcttgcc	tactcccttt	cctcctcctc	ctcaaaaaaa	22260
aaaaaaaaaa	aaaaaaaaaa	aaccacaaca	acaccacaag	actagccact	aagaggccag	22320
ccatgtagta	ccctttttct	gaatagaaaa	agtgtctgtg	caatagtagg	atcatcctta	22380
gtttaacctt	ggattatacc	aggactgaat	tataaaactg	ctgatttgct	ggagtgggtg	22440
ttagcagtag	gaccctggct	ttctccttag	tagctctttt	tttttttttt	ttttttggag	22500
accaggtctc	tcactgctgc	ccaggctggg	gtacagtggc	acgatctcgg	ctcagtgcaa	22560
cctccacctt	ctgggttcaa	gcgattgtcc	tgcctcagcc	tcccaagtag	ctgagattac	22620
aggtgcccgc	caccaccccc	ggctaatttt	ttgtattttt	agtagagatg	gggtttcatc	22680
acgttggccca	ggctggtctc	gaactcctga	cctcaagtga	tccaccacc	tcggcctccc	22740
aaagtactgg	gattacaggc	atgagccact	gtgcccggcc	tccccttaat	atcttcttaa	22800
agtgtcacia	cgcagtagtc	agccagcata	tatttgagta	gctgttacct	gtaaagcatt	22860
tgcatttaatt	ttacttttta	gagaatataa	ttttttttca	tttaagcact	gccttttcat	22920
tgggtgctttt	tgctttgtcc	tgtggtataa	agaaggatca	tgaagaaagc	agaataagta	22980
gacttaatat	aacattcatc	tttctaaatt	accaaattcat	atctctttct	tagacctttt	23040
tacttttagt	ctttgcttta	ataattggat	tatttttggc	ataatttcct	ttcaaagaga	23100
ccaaataaat	attatttttac	aagagttact	agatttgcaa	caaccacaacc	aatgacatgg	23160
tggttggcca	agtaaatcca	ctaactagat	gacttgctcc	tacacatcca	acctggaatg	23220
cacaagatta	gtattttccc	ctgtgcaaaa	ataaaaattg	aggatggggg	atttgatgat	23280
gttagggaat	tattgtttat	ttaatgcaca	agattagtag	tttcccctgt	gcaaaaaataa	23340
aaattgagga	tgggtatttt	gatgatgtta	gggaattatt	gtttattttg	gtaagtgtga	23400
taatggctct	gtgggtgttt	gttttggttt	taaaaaacat	gaccttttcc	ttcacagtaa	23460
gaaaaaggat	aaaaaatgat	atggacagga	tattaggtaa	tagtattgtg	ttgatgttta	23520
attttttgaa	tttggtcatt	gttttctgtg	taagagaatg	ttcttatatg	ctgaagtgtt	23580
taggggtaaa	tggagatgag	gtccgcaact	gacttcaaaa	tgggtcagga	aaataagttg	23640
tagaaaagtg	tttgggggat	aattgaaaga	tgattggcag	agtattttgg	taatggttga	23700
agctgcacgt	tgggtatggt	agctcattat	actgctctct	caactgttgt	gtatattaca	23760
cagttttcaa	aataaaatgt	aaaatgaaaa	aattacctta	ttgcgtgagt	ttttgggttt	23820

gtgaagaatt	gaaactatga	atgataaact	tccacagtgt	cgggggttga	ctgtaattta	23880
atttacttgg	gaatttttaga	ctgattttgcc	tttcttggga	atttttttcc	ctaacgaaaa	23940
atgctaagac	cattttctttt	cctttaatag	agtataaaaga	agaagaaaac	agatttttatt	24000
ccctacaggg	attctgtact	tacttggctc	cttcgagaaa	atttaggtat	gttgaccact	24060
agtgaaaaagt	agttatcttt	ttaactttga	ttatcttttg	tggtaattg	tcctgtgtct	24120
gttttgagaga	caaataatct	ctagatattc	atgttactac	ttgagtgttc	cttgccctgtc	24180
ttcattttta	ctcctaattt	cagttgggtg	gattcccttg	gttttaagcc	agggataact	24240
cccatatctc	attttcttca	gggcatgaac	tgtgcagtag	ttgtcctccc	gactcccat	24300
acttgcttca	gattattgct	ctttcttcga	agttattttca	aaactccatg	cttctctgag	24360
gtgcacaaaa	ttagacattg	ccaaccctta	atcctcctcg	ttgctgatgt	tttcagacct	24420
tcaaactcact	gttcttcatt	cacttctcct	actttttttt	tttttttgag	atggagtctg	24480
tgtctgtgcc	caggctggag	tgcggtgggtg	caatcttggc	tacttgcage	ctccgcctcc	24540
tgggttcaag	cagttctcct	gcctcagcct	cccagtagc	tgggattaca	gacacacacc	24600
accttgccca	ggtaattttt	ttgtactttt	agtagagaca	gggttttgcc	atgttggcca	24660
ggctggtctc	gaactcctga	cctctggtga	tccgcccacc	tgggtgaaag	agtgagactc	24720
cctctcaaaa	aaaaaataaa	agaaagaaat	ttgacattac	attaaatctt	aggatctgag	24780
attacaatat	cagatttgac	acattccata	aaatttttct	tctaaaatga	gtgtgatttg	24840
atactcatga	ttaatctttt	taggtggcaa	ttctcggact	gcaatgggtg	ctgctctgag	24900
ccccgcggat	atcaactacg	atgagacttt	gagcactctg	aggtaacttc	ttttgatctc	24960
agtaacaaca	tagaccacat	tgccatcaga	agccattttg	tgataccatg	gatgttttta	25020
tgctatctgg	gtagttattt	atgtaataaa	tgttcccttc	ctctcataac	agggaaaaat	25080
tagggaaaaa	gtgatttgta	ttataataga	atttacctct	ttatgggttt	cagtttccact	25140
gggaaagata	tgaacttttt	atttttttat	ccctcatgga	tacctagata	aggcaggagc	25200
tttttcatag	gaaagatttt	taaattgctc	aagtgagatt	tttgggtaac	ttttaggtaa	25260
tagtgacatt	ttcttattca	tttctataaa	aagaagtgtc	ccccttcttg	tagtgtttgt	25320
cgttatagaa	atgctgaagt	cggccaggcg	cagtggcccc	cacctgtaat	cccagcactt	25380
tgggtggccg	aggtgggagg	atcacttgag	gtcaggagtt	tgagaccagc	ctggccaaca	25440
tgatgaaacc	ccatctctac	taaaaataca	aaaatgagtc	gggcgtgggtg	ccatgcacct	25500
gtaatcctag	ctactcagga	ggctgaggca	ggagaattgc	ttgaacctgg	gaggcggagg	25560
ttgcagttag	ccgagatggc	gccactgcac	tccagcctgg	agacagagcg	agactccatc	25620
tcaaaaaaaa	aaaaaaataa	attctgaagt	cataataagt	aataaaatta	taggaatatg	25680
tagcaatatt	aaaaaattac	caacatagcc	aggtgcagtg	gtgtgcacct	gtagtccatg	25740
ctacttgga	agctgaggca	ggaggatcac	ttgagcacag	gagtttgagg	ccagcccagg	25800
caacgttagtg	agactctgtc	tcttaaaaaa	aaaaaaagaa	aaaaacaatt	accaacagaa	25860
tgttcataaa	aagttttttt	tgtttgttag	tttttgtttt	tgagtcttgc	tatgtggccc	25920
aagctggagt	gcagtggcaa	gattgtagct	cgctgcagcc	tcagacacct	gggctcgtgt	25980
gattctccta	cttcagcctc	tcaagtagct	aaaactatag	gtgcatgcc	ctaggcctgg	26040
ctaattttta	aaaatttctt	gtagagatgg	ggggtcttgt	taggttactc	aggctgggtt	26100
caaattcctg	gcctcaagtg	atcctcccca	cttggcctcc	taaagtgttg	ggattacagg	26160
tgtgagctgc	ctcccagggc	ctgtattttt	ttatgtatga	gtaggtttca	ttactttagt	26220
ctggaaactc	ctacccttct	tccatgaaag	acacttgatg	taatggaata	aacactatac	26280
agtctcataa	gttgggttta	agtcctctcr	ctgttatttc	cattgtgatc	ttgggcaaat	26340
cacttaaat	catctgagcc	tttttcttca	tctttaatag	gtactataaa	gctttataca	26400
aatgtattat	tatcaaggct	cataaaatgt	aaacactcag	gatatttgat	ttagagataa	26460
tagtatgtta	cttatgagaa	gtagaacata	tgagaaatga	caagaacaaa	ttttcttttt	26520
ggatctagat	atgcagatcg	tgcaaaaaca	attaaatgca	atgctgttat	caatgaggac	26580
cccaatgcca	aactggttcg	tgaattaaag	gaggaggtga	cacggctgaa	ggaccttctt	26640
cgtgctcagg	gcctgggaga	tattattgat	agtaagtga	ttaaggatcg	ttacaaaatc	26700
taatcctttc	ttcttcaggg	ttcttattca	gcgttcttat	atttaaaata	aacttcaagt	26760
taaggagcat	gatgaagcta	aatggtagtg	aaaatgtttt	gttttggttt	gttttctttt	26820
tccttaaaaa	acccaatgga	aatgatctct	agattccttt	atgtaatgtg	taccagacta	26880
tccatttgaa	ctcaggaaat	tagtttttaa	tcgcaaatat	gtcactagta	tttttggttag	26940
aattctgttt	tagatcagca	atgaagttta	tttggaaact	ttaaactgtt	atcactgagc	27000
aattaattac	ttcatagaaa	tgttatttga	aattttgagt	tttgtttcat	gggttgctct	27060
ggattagtgt	cgtggcagta	atataagaaa	cgcaccattc	aagcctacct	tcagataaga	27120
ttcacctttg	aatggcccc	tcctcctgta	aaagtctgga	gtgccccgtg	ctgcctatga	27180
acatatgaag	gggaaaagtt	caattcctat	gaagtccaag	aactacactt	ccagagagta	27240
ttgttttaca	atgttgtatt	aaaattttatc	ttaaaaaatc	tccgtgatgt	gcctatccag	27300
tggcttattc	agaagtctc	ttaaagttagt	aaaatggatt	catataggaa	tagaatatgg	27360
tagaatatgg	gttagtttag	taaagcattt	tttttagtg	taaagttaaa	cttatattcc	27420
ttgaagttct	aataagattt	tcaaggcctc	ttctagttaa	attcttctta	ttgattagag	27480
tcgagtttta	gatgaatacc	gtttgtaaaa	atgataccgt	attctaagta	tttaactttc	27540
catctcccca	aatgcaaaaa	actgctctgt	tccctctctg	tgtcactgct	gaacctgaac	27600

tttgaccctt	ctcgtatatt	ccctcctgct	taccttcagt	tgatccattg	atcgatgatt	27660
actctggaag	tggaagcaaa	tgtgtgtatt	tcacatattg	gttattttcca	gtactctgac	27720
ttgctaactc	gcctctgctg	gatcagcctt	aaaataagct	ttgcatgctg	gcaatcctta	27780
cagggaaatc	ccagacagag	catagtatag	aatcagaaat	gatcctatatt	ttgcccttct	27840
tcatttagca	aactaatctt	tgctgaattt	cagtttgtgg	cttgagtga	gttgaaaggc	27900
ttaagaatct	cttgctgctt	ctactcttca	gtttttccca	ttgtgattca	attaaatcca	27960
aaaatcatag	atgatagcaa	ctgacactta	ctgatttttg	ataaatatgcc	aggccctatt	28020
ctaggcacct	tgaatatatt	gtccctttta	attctgttaa	tggctctgaa	aatatatttat	28080
tatgttcctt	ttaccaataa	tcatttttag	tcagcaaaac	tatctatgat	ttaagagctc	28140
tgtagatggt	gggctaagac	tcatgagtgc	ttgtgctaag	ttgcttttct	ccataacttg	28200
ggcctcggaa	tgattagata	aacgtttttt	cctttttttt	tttttttttt	tttttttaga	28260
cagtctcact	cttgtcacc	aagctggagt	gtaatgacat	gatcgtggcc	cactgtaacc	28320
ttgaactctt	gggctcaagt	gatctttcca	gctcagcctc	ctgagtatct	gagactacag	28380
ttgttcgctg	ccacacctgg	ctaatttttt	aattttttat	ttgtagaggt	gaggggtctc	28440
gctgtgtttc	ccaggctggt	ctggaacccc	tgacctcaag	cagtctcctc	gcctcggcct	28500
tccaaagtgc	taggattata	gatgtcagcc	accacaccca	accattcctc	ttttcataag	28560
tcacaaatta	aaataactta	ggtgtctaaa	atgactcaaa	cattttttaa	gagtcacatg	28620
aacctcatta	ttctctccac	atatatctta	agtgtgtgag	gaaattatatt	cttacccttt	28680
taaggtaatt	catcaagaaa	ttgttatctt	agaagaaaaa	tgtcttgatt	ttggggctaa	28740
tttttagatt	tgtactgagt	tatttctttt	tttttctttt	ttcttttctt	ttttgagatg	28800
gagtcttgct	ttgtcgctag	gctggagttc	agtggcgga	tctcggctca	ctgcaacttc	28860
cgctgcctgg	gttcaagcga	tttctcctgc	ctcagcctcc	cgagtagctg	ggactatagg	28920
tgcgccacc	cacgcccagc	ttatttttat	atttttaata	gagatggggt	ttcatcatgt	28980
tagccaggat	ggtctcaatc	tcttgacctc	gtgacctcc	tgcttgggcc	tcccaaagtg	29040
ctggcattac	aggtgtgagc	catcaagccc	agccacctga	gttgcttatt	tctacagtaa	29100
ggcatgtaat	gaggcttggc	agacagtga	aaaatgcata	ggagaaacca	aacatagccc	29160
tggtctggtg	tatgaggtta	gggcttaaa	gcactataat	gtcctgatac	agaagtgact	29220
cttggaactg	aacagctctg	tctgagattt	ccaaagctca	acgatactgg	tgtggaaaag	29280
gtgccttctt	ctttttgctt	ttgtgctcaa	ggtgttgctg	ctgccagctt	gttgcttacc	29340
tcttttctta	gcagctgcgg	catttctctga	aagaccatgc	atgactaagt	tttaattctg	29400
tattcctgag	ggaaagcagc	cctgactcct	tctcagtttc	cattttgatc	tatttctctt	29460
tattcctttt	agggcttgca	tgtgaccatt	ggacctaaac	tctgtccatc	ctggcttttt	29520
ctccaatgga	ctaatttttt	gctctaatec	ctatccccct	tttaaaagat	gcctctttgc	29580
ttgttgccct	atgttctgta	tttgcagatg	gtggttttca	aactgtcttg	aagcaaccac	29640
tggttttaag	ggagttaaaa	aactggcttt	aattaaatag	tagagatgag	taagcccagg	29700
gagaactaat	agtatttttt	tccagctatg	tgtattgact	catctccttt	gaggaaaact	29760
gacttagttt	tgatatgaat	tttagtttct	caggttttaa	aataagctaa	ttttaaagtt	29820
ggattgcacc	aatacttaag	atttaccttt	ggtatctttt	tgcaaaggat	gactaatgaa	29880
agaaagtaat	gaaagaaagt	acaaacagca	aattaattaa	ttttgttggt	tgtggcttaa	29940
tttttttttc	ttttaccaat	actgagtgtg	aaagcaaaat	taacagttct	attaaagact	30000
tagtaaatgg	aagtatgttc	ctaaaacttg	cattttatag	taggaagcag	gcttagcagt	30060
gacctatggt	tgtactgtta	ggctaattaa	gggctttaa	ttagcaata	atcagcctta	30120
ttttttgctt	gctctgacag	agaatggaa	cctgcactaa	aactcattaa	tctgctaatt	30180
gttcttccct	gttgaggcaa	tcaattagtc	ctgagttttt	aaccaggact	tatacaaaat	30240
tctctcatt	tctataatta	cccctttttt	atttaagaca	tattaaaaat	ccttttagca	30300
tgttgaaatg	ttgccataca	aagcatacta	ctcagtgcac	aaattttggt	tttaactgtaa	30360
cattttcttg	gaaatttttg	ctgaattaca	cgtaaacactg	agccagacca	atatggatta	30420
ttattgataa	ataaatgac	attgttttcc	cttactgtta	caaactactg	gaaaagtgtg	30480
aatctgaaga	cattgtagac	taggaacact	agaaagacta	aatggaaagt	agtttggtt	30540
tttttagggag	agtgcctaag	cagtatcctt	tgaggctctg	gtttatcccc	ctgcgagata	30600
cagcgtaaac	ctacattgga	atcttttagt	ttgtaaatgc	agaaatgatt	ctgaacctgt	30660
aggataccac	tattaaagac	agacatgttt	tagaactgcc	atctagaaca	gttgtaacct	30720
gtgtttgtat	tatagcgtgt	cttggctctg	gcgctttaat	gctgtccttt	cttacaacga	30780
tattccttcc	tgtctttttt	tcttctctctg	cttctttccc	ttttccctac	ttttcctgcc	30840
ttctcttctt	tctatctccc	agatctgaaa	gattttcaga	acaataagca	tagatacttg	30900
ctagcctctg	agaatcaacg	ccctggccat	ttttccacag	catccatggg	gtccctcact	30960
tcacccccat	cttctctgctc	actcagtagt	caggtgggct	tgacgtctgt	gaccagtatt	31020
caagagagga	tcatgtctac	acctggagga	gaggaagcta	ttgaacgttt	aaaggtaagt	31080
aatagttcag	actgaataca	aggtattcta	trtagctcca	caaggaagaa	ctaggagtaa	31140
aaatcactaa	gatttcgact	cagcatatgg	agaactcttt	gactcttaga	agtgtcctga	31200
aaattgaatt	ttgtgctttg	taagttaatt	tcttttcatt	agaatgcctg	agtctgacat	31260
ggccaggcag	agttgggaga	agtgcattgg	cagttcatag	gctggcaagg	cagagtaaca	31320
gattaataga	tgtgtacgtt	aattctggga	tagtacatca	agttacagtg	taattgtttt	31380

gttagaaatt	tatttgga	cgtggtttca	ttgaaagatg	gaattctgtg	tagagccatg	31440
cttctgttaa	taaagatgaa	gtttgggccc	ggtgcagtg	ctcacgcctg	taatcccagc	31500
actttgggag	gccgagggcg	ggtggatcac	gaggtcagga	gatcaagacc	atcctggcta	31560
acacggtag	acctgtctc	tactaaaaat	acaaaaaatt	agccgggctg	ggtggcgggc	31620
acctgtatgc	ccagctactc	ctgaggctga	ggcaggagaa	tggcatgaac	ccaggaggcg	31680
gagcttgca	tgagccgaga	tcgtgccact	gcattccagc	ctgggggaca	cagtgaact	31740
ccgtctcaaa	aaaaaaaaaa	aaaaaaaaaa	aagataaagt	tagaacattt	gagtagctgt	31800
tccaggttta	ttcagatagt	tattaaatta	tatcttctag	aagttcacag	aaagccttga	31860
tgccctatga	ttactgtaca	ggaaggttat	ctcctcctat	gctggtaccc	atggggtcac	31920
agggccatag	tgctgcatgt	atgcagatta	ctggcctatg	gctcccagct	ttgtgttgg	31980
aggtatttcc	cagccccttg	ctggctctgt	caggttcgat	cttatctctt	tgtatgaata	32040
tgaaagttaa	gagcagcaga	gccttctgac	tcataagctg	ttctcttcct	cctcaattaa	32100
ttcagtgca	agacagttgt	cacctgcagt	gctgaaagct	gttcaggaa	cctgccaccc	32160
atcagatgac	tgtgacagca	tcattctgct	tgaagcaaat	attttttctt	tccctacttg	32220
tcacctccag	gaagatagtt	tgaccatac	cttttagcca	ttgtgttcaa	gagtgctagg	32280
gtaggggaca	cagggaggac	aaaaaacact	ttttaagagt	ctgagaagtt	gcagtcagtc	32340
tgccagccag	cagtcttctt	cacagtcact	tggtattccc	cttgactgcc	attcatgcaa	32400
ggacgataga	gaacagtgac	acattgacac	atttaataag	taaaaggctt	gccgaaaaac	32460
tctttaacac	taaagtgaat	taagcttgat	aattgaatat	gtcagatgct	ttcattttaa	32520
ttttcatgtg	aggctgaaaa	agattttata	tgaagaaaga	ggttctgctt	tttctttctt	32580
aatcagccta	ccttttccct	tgacatttta	cttacttcaa	aagtacagt	tgggccgggc	32640
atggtggctc	acacctgtaa	accagtgact	ttgggaggct	gagggaggag	gatcatttta	32700
gctcaggagt	ttaaggctgc	agtgaagctg	gattgtactg	ctgcactcca	gcctgagtga	32760
cagagcaaga	ctctatctca	aaaaaacatg	ctttagaac	aaaaataagg	ccgggcacgg	32820
tagctcacac	ctgtaattcc	agcactttgg	gaggccaagg	tgggcggatc	atctgaggtc	32880
aggagttcaa	gaccgcttg	gccaacagtg	aaaccccatc	tctactaaaa	atacaaaaaa	32940
attaagccgg	gcttagtggc	tcagtcctgt	aatcccagca	ctttgggagg	ctgaggcagg	33000
cagatcactt	aaggctcagga	gttcaagacc	agcctggcca	acatggtgaa	accccatctc	33060
aactaaaaat	aagaaaatta	gccaagcatg	gtggcacacg	cctgtaatcc	cagctgcttg	33120
gaggggtgag	cacaagaatc	acttgaactt	gggaggcgga	ggttgccacg	cctggagatt	33180
gcgccattgc	attccagttc	gggcggcaga	gtgagactct	gtctcaaaaa	ataaataaat	33240
aaaaatagaa	aaattagccg	gccatggtga	cgtgcactca	tagtcccagc	cactcgggaa	33300
gctgagggca	gagaatccct	tgaacccagg	agggcgaggt	tacagtgagc	taagattgctg	33360
ccaatgccct	ccagcctggg	caacagagtg	agactccatc	tcaaaaaaaa	aacaaaaaca	33420
aaaataatgt	gcttgtttct	tccttcagaa	catgtaagt	gaaagtgagt	gaagttctgc	33480
aacacctagc	aaaggagaaa	ggagtgaaga	atctttccat	taaatgttgt	aggctgcagg	33540
gtaactttta	ctgttggtaa	aattgacttt	gaaattttac	taaagacctt	agtgaagagg	33600
agtggtaaga	ctccattcac	acattttctc	acaataggcg	tcagtgaacca	tggtgctttg	33660
cagggctctt	ttcagagcca	gagtgagaaa	tagggagctc	cttgggaaatg	agaagtaaaa	33720
ctcccactta	ctgagtcctt	aatataaaat	gttctttttc	tttggggatc	ttagttctta	33780
actatttgct	tattaaagcc	ttagctcwtc	cacttgagat	taccctttcc	tttgaccttc	33840
ctcttgatgt	aaaggagaaa	tttctcccta	actgccaaat	gggaccaatt	gaaaataatg	33900
tcacatgtta	ggagtcaaga	cattaaccag	actcttcaag	attcaggtaa	cccagtgctc	33960
tccaaataat	ttctagggaa	taacactgtc	attattcagg	gaactctcct	tccgacttcc	34020
ctatgtccca	ctaaagcaag	acaaaactgt	attacttcaa	tcagaaaaaca	aatttctctg	34080
ttttaattct	tccttacgcc	aaaataaaga	actcttctgt	tttccctagt	tttattctag	34140
aaagaaaaca	gcttccttag	tgttttgtgt	tgtcttgagt	ttttaatgca	gtgtggtagt	34200
tcactgaagc	ataagcacct	tagttctgct	cttattttgt	gaatggtgag	aatttctttc	34260
gggcagacct	gcattttaat	gctgcacgca	gataacatct	ttcatccgta	gcgccagggc	34320
aatggttggt	cagttagaat	gcttgagaat	atctaggaac	tggtttccta	aggcatggtg	34380
ctgcaggaga	aaaatattat	aatggaaaac	agcaatagag	gaggttagtt	tattttccta	34440
acattttatt	tccctttggt	tctttttgac	tttaaaaaag	gtcagttgat	tggtactgt	34500
ttttacacat	gatgttgctg	tggtttgtgt	cactgaatag	cctttttcca	tatgttaggt	34560
actcatgtca	caccattgta	aagatgtttt	tttttttga	gacagagtct	tgctctgttg	34620
cccaggctgg	agtgcagtga	cgcgatctcg	gctcactgca	acctccgcct	cccaggttca	34680
agcaactctc	ctgcctcagc	cttctgagaa	cgtgggatta	caggcatgta	ccaccacatc	34740
tggttaattt	ttgtattttt	agtagagacg	gggtttcgcc	atgttgcca	ggctggtctc	34800
aaacttctga	ctacaggtga	tctgcctgcc	tcggcctccc	aaagtgtctg	gattacaagc	34860
atgagccacc	atgccaggtc	ttcctgtcaa	ataaggtttt	aaacatgcag	ctgtagccac	34920
agctctcgtg	tttattagtc	tggcatcaca	gtcagtactg	tgagtgccgt	tgcttaggct	34980
tagcctataa	atattccaaa	atatccatta	cagaagcagg	caagggcaaa	tgctgtgcta	35040
agcagcccaa	gggccacatt	tggaaccac	ttaggaagca	tgctatgttc	agtacagtaa	35100
aaccttggtg	attcaatggc	ttcaacaaat	gttctgtgta	gtaagtgttt	ttcatggaaa	35160

tattttctgac	ttgtcctggt	gattttctgtg	tttccttaac	tgaatcatca	tcaagaagag	35220
attaaattgt	ttagctatat	gaacataatt	tattaaccag	tgctatacaa	ataactagag	35280
atgcagacag	atrtgacgtg	agaccaaaga	gtgaatgaca	cctgagcctg	gcattccctcg	35340
gagcttaaaag	aagagttcct	ggtgtccctc	tgtttttttg	aatttttttt	ttttcaaaga	35400
aaggaggag	gattaaagat	ttatctcctc	aaccctcat	ttctgaacca	tgaatccaca	35460
tttcaagttc	atcttacatt	aggcccctag	agtggggtat	gggaagccct	ggacaccagc	35520
atgcctgcct	gccagcagaa	tgagttgatg	ctcctgagac	tgaatgctgt	tttgacttg	35580
gcttccttat	ttatactaag	tttgctactc	aggcttgaac	agaatattcct	tcactttttc	35640
ttaacttagg	gtgaactgtt	tcatagaaca	tttacttgaa	agcctattga	tcctgtttta	35700
tttgaagaaa	gaacagttag	tcagcaaata	gctctagacc	tgtctcagtg	agtagtggtg	35760
gctcacttag	gtctctcgtg	ctgttggaac	acagtggcac	catcttggtc	cagtgcaccc	35820
ttcgccttcc	aggttcaagc	gaatctcctg	cttcagcctc	ccaagtaact	gggactacag	35880
gtgctgcca	ctgtgcccg	ctaattttcg	tatttttagt	agagatggg	tttcaccatg	35940
tgagccagct	gatcttgaac	tcttgtctct	ggtccagttt	tcttatctcg	tgaatggttc	36000
cagaataaac	gattttgcag	gcttttttca	gccctgaagt	tgtgctaata	tgactcacag	36060
ttattatatt	ctctaagttt	cccccaaagt	ttttaaaaat	agctttctag	acaacctaaa	36120
aattgcagct	gaaaactgta	ataaaaataa	aagttgtacc	ttactatta	aatctctttt	36180
ctgcaaagag	atcatgttct	gattttaccct	tcagttggta	agtaaaatta	gataagtcac	36240
attagagttg	gctgcaaaat	ggtaaagagga	ttcctagaca	gtaattttgc	tttcttagag	36300
tgagttctac	agttgcccc	aaataggatg	gctttaatat	gactagataa	tcctctgctc	36360
tgaataaacc	tttttctgaa	ggtaatcaac	taggaagaaa	tatttttaaa	atatctcata	36420
tgtcagtttg	tattccataa	tcaacagtgg	tttattgtgg	ttagtactat	aataagtact	36480
agtcaagatg	ccttggttag	gtcgggtgca	tcctgttgtc	ctgtctgttg	aaggttttca	36540
ctttgagaag	attgtataga	tttctaccca	gctatcttta	atagacctta	aaattatgta	36600
ggttatatac	tcagcttgct	ttgtatttgc	ctatttaaaa	tcatatgcag	ggtggggctg	36660
ggaggaaaag	aaaaaaaagg	aaaaaagcag	tcatatgcaa	aagcattttc	tttttttaaa	36720
gaggggctca	ctgtgttgct	caggctggag	tgtagtgggtg	cagtcatagc	tcacttatgc	36780
agaagcggtt	aacaccactc	ttagtgggtg	ttgaataact	tttgtgggtg	tttatagata	36840
aactattgaa	aacttggttc	ttggccttct	cactccttag	ggaattgtgt	ggatttgtct	36900
tgctcgctgg	ccctctagta	tgacactttt	gttctccatt	tgtggggcgg	catgtgcact	36960
gcatggtaat	catttctgct	gaggactgtg	tagagatgcc	attcaggtag	ctgttttagc	37020
ctttgagcct	ctgtgtgtgt	tctgaagcct	aatgagagcc	atttcagggtg	gtttttgtca	37080
cttgccaccc	ccaactgcag	acttgttcta	gcttcttcca	ccttgtaagt	ataactcttt	37140
tctgaaaaag	tacctcctac	atggctgcct	tactgatgcc	caggttcctt	ctttaatctg	37200
ttttgctctt	tgctacctt	tgctgggtgt	tggtgttctt	catctttctc	tttttttaaa	37260
attttatatt	aatcatacc	ctcctatttg	ttccaccaaa	ctcatatact	actttgcttt	37320
tgcagctgct	gctttgaatc	tttactcttg	cttttttcac	gtgttccctt	gccttgccct	37380
tggtatgttt	tttgttttga	ggtgacatcc	agttcatgta	atattctgga	agttccaagag	37440
atccagaaaag	tcaaacacta	gacaacttaa	ggtttattac	tcacaagtcc	tttctgtgta	37500
gatattttac	gttttgggtg	cacaggcttc	gttatgtaacc	tgttatcata	tgaatcatac	37560
attgaagtgt	ttgatgaact	ataacacttt	tccttgttaa	tcactactag	catcattact	37620
atatttcatt	atcatcatct	gactttctta	cttcatctcc	atgaccagca	ttttcacctg	37680
ttttggcatg	tgctcgctcc	ccttttttgt	actccttttc	cttccattct	ttgcctaatt	37740
gctactgttc	cttcatttct	ctgtctgggc	ctcactcagc	cttgattcac	ccatgttaca	37800
gagacctgtg	aatactccag	ttctagctct	tagtacactg	ttctcttata	tcctactgta	37860
ttcgtttctc	gtcctcagac	atgctgctct	agcaacacat	tattcctaga	aaactccagg	37920
tactttccca	ctctcagcat	accacatgg	cagcttctgc	tcttctttga	tctctgtcca	37980
catatcactg	agtctgtgag	gcctgtcctg	acaacctcac	tttagattgc	tcctcacct	38040
acccttcccc	atgtgttctt	tcctcttgtt	gctttatttt	tgtctggagc	acttgccact	38100
atctgacata	ctgtgtactt	attgattaaa	atggattata	tcatggcggg	gcacgggtggc	38160
tcacacctgt	aatcccagca	ctttgggagg	ctgaagcggg	ccgatcacct	gaggtcagga	38220
gttcaagacc	agcctgacca	atatggtgaa	acctcatatc	tactaaaaat	acaaaaactt	38280
gccggacata	gtggtgtgca	cctgcagtgc	cagctactca	ggaggctggg	acaggagaat	38340
cgcttgaacc	tgggaggtgg	aagttgcagt	gagccgagat	cgcaccactg	tacgtatgcc	38400
tggtatgacag	aatgagactc	tgtctcaaac	agacaaaaca	acaaacaaac	aaaataaata	38460
ataggattat	atcagtctcc	ttactacaat	gtaaatctta	tgaaggcaga	gattttgttc	38520
tctgacatat	ccataggtgc	cttgaagaag	gcttgatatg	taatgggggt	gtatataagt	38580
attgaatgaa	taaatacatca	cttacttctc	tctctctctc	tctggtatac	agtaacctcc	38640
ttgagagcag	gaaacatgtt	ttcgttcttt	ttaaattaaa	ttaaatttaa	tttaattaat	38700
ttattttttt	aattttactt	tacgttccag	gatgcagtgt	cagaatgtgt	aggcttggtta	38760
cataggaata	catgtgccat	ggtggtttgc	tgcccctatc	aacctgttat	ctgggtttta	38820
aacctgttat	gcattaggtt	cttatattta	tataagtgta	aacactcagc	aagtgattgg	38880
taaattgtca	cttcagtgtg	tgtcttcaga	tgcgatttaa	tagtaaacct	ttctatgttg	38940

actgctgggt	tgtgttcagg	aggtttagtgt	gttatggcga	aattgtagtg	aaacaaaagc	39000
agtaaatcacg	tttgggaact	ttattctttt	ttttctttt	ttgagatgga	gttttgcctc	39060
gttgctcagg	ctgcagtgca	gtggcgcgat	ctcggtctgc	tgcaagctct	gcctcccggg	39120
ttcatgccat	tctccttcc	cagcctcctg	agtagctggg	attacaggcg	cctgccacca	39180
tgcccacgtg	gatcacgagg	tcaagagctt	gagaccatcc	tggccaacat	ggtgaaaccc	39240
cgtctctact	aaaaatacaa	aaattaactg	ggcgtgggtg	cacgtgcctg	tagtcccage	39300
tgctcgggag	gctgaggcag	gagaatcact	tgatccccag	aggtgagggt	gcagtgatct	39360
gagatcgcg	cactgcagcc	tggtgacaaa	ggaagactcc	gtctcaaaaa	aaaaaaagac	39420
ttggaggatt	tatttgtttg	gggttgattt	agtttttaggt	ctaattcttta	tgattcttat	39480
cttatgtatt	aataaaagaa	aaatctcttt	acattttaatt	ctggcctgta	tcctcaaagg	39540
gcatgatattg	cttcttgctt	ttctagcatg	gtatgttaaa	ttataagact	ctttgcctct	39600
ttactgtgc	gcttccttcc	ttaggaatca	gagaagatca	ttgctgaggt	gaatgaaact	39660
tggaagaga	agcttcgtaa	aacagaggcc	atcagaatgg	agaggtcagg	aggttaaaat	39720
ctggaaatgt	ttctaagttt	tctaggggat	gtgattact	ttatcgtaag	aaaagataaa	39780
atataaatta	aaatcaaaga	gtctgccctt	ctaaaacaca	aaacgaaacc	aaaaaaatgg	39840
acagggattt	atggccagtt	gaatttatgc	tttgaaatga	catctacata	tttaaatatt	39900
atctagggag	tcacaggttt	attagttacc	atattacatg	ttgctaggga	ttcataaaaa	39960
tggaacatat	tggccaggcg	cgggtggctca	cgcctgtaat	gccagcactt	tgggaggctg	40020
aggtgggcag	atcacgaggt	cgggagatcg	agaccatcct	ggctagcatg	gtgaaacccc	40080
atctctacta	aaaatacaaa	acaaattagc	cgggcatggt	ggtgggcacc	tgtagtccca	40140
gctactcggg	aggctgaggc	aggagaatgg	cgtgaacctg	ggaggcagag	tttgacagtga	40200
gccgagatgg	tgccactgca	ctgcagcctg	ggtgacagca	agactctgtc	tcaaaaaaaa	40260
aaaaaaggaa	tagcctcttc	ttttctttt	tatattaata	tgcatatcat	ttatgaatta	40320
ttgactatcc	caaagtcaag	tgagacaaag	ttcttatctc	agtgaatata	actttaacaa	40380
agcaagattt	cattgatattt	gaatgtgatg	ttgcttttag	aaataacttc	tcattaatga	40440
tgagataagc	acattatattt	atgtttgcat	tattggctat	aaacatggcc	agttgttttt	40500
gcttttgagg	ctttattttc	caattcatat	tagactgatt	taattgacat	tttggatta	40560
gtgttctgat	atacctgttt	ttttcctaga	gaggctttgt	tggctgagat	gggagttgcc	40620
attcgggaag	atggaggaac	cctaggggtt	ttctcaccta	aaaaggtagg	aaacaatgct	40680
gtgaaaccta	atcagacaga	gacacttttt	gtttgtcttt	gttactgggg	cacttcatgt	40740
ctttcaggac	ctactctccc	ccttattatc	ttgatacttg	ccttaatttc	agatatttct	40800
gatttcatct	tagagcatgg	tcttatagta	caaaggcagc	ctatttctaa	aatgcctatc	40860
ttgtttactg	tttacaanaag	aaaaaaccta	aggcaatatg	taacctttgg	ccatgattca	40920
gtttgtttag	tcatctttat	ttgttgcttt	ctcttaaat	cccagcctaa	cctagcgtac	40980
tttcagtttt	ggctactctt	tcatgtagat	tttccacag	aagtcgtcta	ccctgcttta	41040
taactttttg	gaaccaattt	cagctgtaca	ggtttggga	gaataatact	gcttccattt	41100
gcttctgcaa	tttgatttat	agacagccat	tgtactaagt	aaagcaattc	ctgtccctat	41160
ccttctagac	tcctctgctt	ccttcttcat	tttttaaaaa	gtgtctcaa	tagttacett	41220
cctttttaca	taatcatagt	tatgaagtag	ttacgtagct	tttagcta	tagagtagta	41280
agggtttttt	tgtttttttc	ttaaactaag	ttgcaacgtc	ttgttgata	ctagaaaatg	41340
actatcaaat	ggttattttg	tgtgcttatg	tatcagaaaa	taggagaaca	tatatcgatg	41400
tacaaagcat	gtattttatg	ccactagtaa	attgtccttt	atttcaagtg	acctcacatt	41460
ggctattgga	gacttaaaaa	tctttattca	taggttctgt	tgatgatctc	tctaagttag	41520
gatacaaggg	gtggtgagtt	ttgaattgg	acttttatgt	gatatgaaat	attcctgtgg	41580
ttttcgctac	aattcttctg	gtcaggtttt	tctatcctgg	ccacataact	tttttttttt	41640
ttttcttccc	aaattccaat	gtgttatatt	caacagaaaa	gactgatgta	aagacatttt	41700
agatgcatat	tgtagaatag	aacagttaag	tattttatat	gtttatatta	tgaagggtta	41760
gtgggtttca	cagttttta	agtttttcc	tcagtaacac	atgtattcat	tcagatatgt	41820
tacttaggat	tatatagatt	ttagcagtta	cagttttgg	ggaattta	tgtgcctatg	41880
catatgtcta	ataaagggtat	tttataacat	cttacagtat	taaagactgt	gaggaaattt	41940
tttttttttt	ttttttttga	gacagagtct	cactctgtcg	ccaggctgga	ctgcagtggt	42000
gtgatcttg	ctcactgcaa	cctccgcctc	ccgggttcaa	gtgattctcc	tgccctagcc	42060
tcccaagtag	ctggactgca	ggcacatgcc	accacgccc	gctaagtttt	gtatttttag	42120
tagagacggg	ttcaccatgt	tggccaagg	tggctctgat	ctcttgactt	tgtgatccac	42180
ccacttcagc	ctctcgaagt	gctgggatta	caggcatgag	ccactgtgcc	tggctggaaa	42240
atttttatta	taacaactag	tgaanaatt	atgcatttaa	cataaatcat	aaataacttg	42300
aacaaatgtt	gagatgtaca	tgcagtgaa	ttccaaagt	agatctctgc	ttccaaagct	42360
ttttcttact	tttaattttac	ccatcttgct	ttagctttct	agacttcttc	atgaaaggaa	42420
gggatatacc	agtgaccaca	ttccaaactca	gtcttaggac	ctgtttctga	gattccattt	42480
ttcaaattat	gatgtgcttt	aaagaatgct	actcacaag	ggctttgact	ttctcaaacc	42540
agtcaagttc	ttcaactaca	aagggggatg	ctagttgaga	agatgtggag	gtttttcaga	42600
tttctcttta	accatctttt	atagttcctt	ttcctacccc	ctcatctgac	tcttctttca	42660
ttcttcacaa	tttgctggca	ctctcttgag	tagattttgg	cctcaagacc	atgtgactta	42720

tttttggatt	ccagtggggt	attttcacta	aagaagggtg	gttttttaaa	aagatttttag	42780
aaatattaga	gttagatgaa	gaggattggt	agaagatggt	tatcttatcg	aatatcagta	42840
attttacttt	tttctttaag	gtaaagacta	ataaattaaa	atcttttaag	gtaaagacaa	42900
attaataact	gggatcagaa	gaactgataa	aaattatagt	gggaagtggg	gggggggtat	42960
ggggacattt	taataaaaagt	aaatatttct	tgagattttt	accaggcat	ttacaaaaat	43020
tataattatt	ccgttcagag	aattggcctg	tcattttttc	cttgtgcttt	gaccttatca	43080
aaaattaggt	cttgtttttt	tatcccagaa	ctcatgtttt	aaaatggaat	gccaggcaca	43140
gtggctcact	cctatagtcc	caacactttg	ggaggccaag	gagggtgggt	cagttgaggt	43200
cagtagttcg	agaccagcct	ggccaacatg	gtgaaacctt	gcctccacta	aaaatagaaa	43260
aattagccag	gcgtgggtgg	gaacacctgt	aatcccagct	acttgagagg	ctgaggcagg	43320
agaatcactt	gaacctggaa	ggcagagggt	gcagtgaagc	aaaatcatgc	cgctgtactc	43380
tagacagagc	gagactctgt	ctcaaattaa	ttaattaaatt	aaaataaaat	agaatgcact	43440
tggttccgtt	tggtgaggcc	tgaagagagt	gttttctctt	tattatggaa	ccgcaactag	43500
ggataaaaaag	aaatccctct	ctaggctctg	tggagagttg	gagaagcctg	gcctcttgta	43560
ggccctctgt	gttaccacag	ctctctcatg	gtgccctgct	ccaaattgat	catctgtaat	43620
ctttttcaga	ccccacatct	tgtaaacctc	aatgaagacc	cactaatgtc	tgagtgccta	43680
ctttattaca	tcaaagatgg	aattacaagg	tatattttatt	tcctgttttg	gtcacttcgt	43740
gtgttttccc	cctcttagat	aattgaataa	ctaaagggaa	ggggttgaaa	aaattaacgt	43800
aatgatttgc	tgtatttttt	gtctgaaata	gttacaaact	atgctctctt	tccaaataat	43860
gtgtttttgc	cactggagcc	agttactatg	tagtttttct	ctgaagaccc	taaataaatt	43920
ttttttcctt	taacaaatat	acattcctta	gggattttat	ttgactcatg	tctttataat	43980
actgtatgag	ttacattggg	atatcagtgc	tcctgttttc	ttattctgat	gtagagaggt	44040
tgatcatatg	tgaggataga	attcatagtt	gaaagcttct	gatgcaacaa	ggcaataacct	44100
tttttgtagc	taccaaagga	tattcttttg	aataaagagg	ttcattgttt	gtagtgcctt	44160
ctgtgtctga	atttccctgg	gaaacacttt	ctcttggtgt	cagggttggc	caagcagatg	44220
ctgagcggcg	ccaggacata	gtgctgagcg	gggctcacat	taaagaagag	cattgtatct	44280
tccggagtga	gagaagcaac	agcggggaag	gtgagcattc	ctggctggag	cttcagcaac	44340
aacattttca	ttttatatta	tgagaaatcc	ttaagacttt	gtattctctg	tctatcagta	44400
gtactttctt	atacaatcta	attctgaaaa	atggagagac	ctgggctgct	tatgaatgca	44460
gagatggaca	aggctgcttt	acatgaaaat	agcttggaca	aaagaagccc	tttttactg	44520
ccaagaactg	agaaggacat	aggcaattag	gcttggctctg	gaatgttaat	tattttaatag	44580
aaaagtaaga	aaatagcaga	tatcctgggt	aataggagat	ttgaaggaca	ttaagtcaac	44640
ccagcagaat	ttatttttat	ctaaaaggga	agaaaaagtc	agtcttgatt	tttgccctggg	44700
ttattaacaa	aacaacaatt	taatggtttt	tttctgttat	ataagtcact	cattccctta	44760
tcaaaatatt	agcttctcag	cttttagttt	ctgtttatta	cctatatctc	atccttacaa	44820
tttctgatgg	ttctgagttt	tattgactga	accgtcagag	atccctgaaa	ctaataattc	44880
ctatcatctt	cttaggttta	tcaaatagag	ttaaatgttc	ttgtgttagc	catggccaca	44940
gatagcctct	tctattgggg	ctagttctgg	taccccaaaa	tgaactactg	tatagacaac	45000
ttcagccact	tgattgattg	cagggattta	ttctacttac	tgcaaatcct	gataagcaac	45060
tgctttccat	tatttgattc	caatagtttg	taatgataac	attagtttgt	gtttgttcct	45120
cttagttatc	gtgaccttag	agccctgtga	gcgctcagaa	acctacgtaa	atggcaagag	45180
ggtgtcccag	cctgttcagc	tgcgctcagg	tgagactggg	agaggtttgc	catcttcagc	45240
aatgtgcaca	tggttctctg	gacaactcta	atttttggct	gtttaaaagg	tgaagtaata	45300
gtcagcatta	ggatttttgt	tcttgtaaaa	acaacagctc	tgaaagctgt	cttttcacat	45360
tagggagggg	tgaggttggt	aaatagatac	tatatattaa	aaaaattatt	tcttaaccct	45420
atttttctgt	tttgtgctag	gaaaccgtat	catcatgggt	aaaaaccatg	ttttccgctt	45480
taaccacccg	gaacaagcac	gagctgagcg	agagaagact	ccttctgctg	agaccccttc	45540
tgagcctgtg	gactggacat	ttgccagag	ggagcttctg	gaaaaacaag	gaattgatat	45600
gaaacaagag	atggagaaaa	ggtaatgcac	agttacgcag	cccatatgac	tgtttcttct	45660
tttaaacatg	taataactaat	agcattcttg	aatttttttt	ttttttttta	cttctaggct	45720
acaggaaatg	gagatcttat	acaaaaagga	gaagggaaga	gcagatcttc	ttttggagca	45780
gcagagactg	gtaggagtcc	tgaatctgct	aaactgttgg	gaaaagggca	gcttgttccc	45840
atactttccc	tgttccacag	agcagtactc	acccaaattg	cttctgtctc	artgatacca	45900
agcactattc	tttaattttc	ttaatggaga	atgaacttaa	atctcccggg	agccttagcc	45960
tgaaaaaata	gtccacagag	gtactctttt	gggcttttta	tgtcttaaag	ccaaatctta	46020
acttctgtta	caaccaaata	ctttttaagg	aaatagagct	tttctggtag	cctttgcctc	46080
ttgatagtgg	ttttggaatt	tgcttcagtg	gtggttcttt	aaatgataat	tactctgaat	46140
attgaatttg	gtgagagttt	gccttggttt	tgtttctgat	cacttgatag	tactaattct	46200
ctgctcttgg	gctgactttg	ggattgttct	tacgctgggc	agactttttt	tttttaagtt	46260
aaactgtgtc	taaaagtgtt	gctgcacagt	tgcatgtgtt	actccttttc	ttatcccttg	46320
catggagtct	gaattctcaa	tcagggttctc	agtggcatgt	gtggtagcgg	tgggagcaaa	46380
ggctgcatac	ccagcccggg	caggacagaa	ggcttgcttc	ctttgaggga	aggaggattt	46440
gagtgagcag	ctggaaaagtc	tgtttaaggt	cccagctatt	gatacaatac	taatggcttc	46500

agcggttttat	aatggtaatt	gatttcccag	atttgataga	aactgaatct	catgaataaa	46560
taagttgatc	cattagttag	attttcattt	ttaaagccaa	ccaaacaacc	aatttttggtt	46620
aacaaagctg	tggttataag	aaccttctca	gcattcattt	gatatgtaag	aataaagtat	46680
aactaacctt	tggatactgt	ggatgcagtt	taggaagttc	tcatttttta	agatttcagtt	46740
gtatctttct	tttaaggtaa	gggggcatta	gggagaaagt	tcaagcactg	gcttctttcc	46800
ttagtataaa	atacagcatg	taaactacag	ccttctagtc	atgctctgtg	ctttggagaa	46860
gaaggggctt	agtgtctaag	gggccagaaa	cgtgaaggac	ttgagcacia	gcaacctttt	46920
ccagaatacc	tcagttctca	ggcatgggca	tggattaaac	cagttagttt	gttatcttgt	46980
aaattatagt	aaagccctaa	aaagaatgga	ttgtaagaat	aaatcaaggg	tttggggcaa	47040
gataatagca	gtgttaccaa	cagaggctct	ttaggaggcc	tttttagcat	taagttttta	47100
tgaaaacgaa	agaactagct	acatgtgatt	agtggcttga	aaggttttgt	gacacaaaaa	47160
tgtcttttat	cttaataatg	tagtacattc	taataacatg	cctctgatgg	tctgtgagag	47220
atcaaatgag	gacaggtgta	attaagctct	attattaaca	ataatgcctt	gtatttgtat	47280
agtgtcttat	aatttacaga	gagcgttaac	atccatcaat	ttatttgacc	ctcaaaacaa	47340
ccctctttgt	ttacttatgc	atatgtataa	gtaaaatatg	taatatattacc	tccagttgag	47400
gaaacccaaa	cttggctggg	cgcggtgggt	cacgcctgta	atcccagcac	tttgggaagc	47460
ctaggcaggt	ggatcctggg	acctcactta	gaataactaa	gcacggtgct	cgcttcggca	47520
gcacatatat	taaaattgga	acgatacaga	gaagattagc	atggcccctg	cgcaaggatg	47580
acacgcaaat	tcgtgaagcg	ttccatattt	aaaaaaaaaa	aaaaagaaac	aaaaaccaca	47640
aaaacaaaaa	aaaacaagac	aaacaaaaaa	aaagaataac	taagcacggc	agaataagga	47700
tgcctgggtg	tatccataac	caagagttgg	tgatcattct	gtgccctaaa	ataaagatgg	47760
ccactaaata	aagagaaagt	gaatgtctaa	gcagtattta	gtttctaggg	aatgtacata	47820
ccctgcagat	aatcagattc	tggttgttgt	ttgctctgtg	tatctgaaag	caagccagga	47880
tttcaagagc	agcttcttaa	agcaaggaag	ttgctttcct	ctcaaaggct	tgtctgttcc	47940
actttcattt	cttgacttaa	gggaggaact	gatttctaac	acttcagcct	gaagaatatc	48000
taccagtagt	aatactgaat	ggaaaatgtc	tttatatat	accctgtatt	ggtcaattaa	48060
cagatgtatt	tctaattcac	cctgtgaata	aacattgaga	gaatacaaa	gaattattaa	48120
tcagtaataa	tttatttaat	atttaccata	tgttaactct	ttttgcataa	gttcagcatt	48180
aaacaatgga	attcaggctc	atgagttttc	tattctgata	gttcccaaat	gcatcacatt	48240
gttaggggtg	tgagtatttt	cttcagttag	tgctctagga	taccctgggt	ctttctgtgc	48300
ttgcttttta	cacacagttt	tgggatcttc	tctcttctta	tttgcataag	ttccttgagc	48360
agtctcttta	ggaggcctac	tttgcatattg	gcatttaattg	aaagccgaag	gatcattttg	48420
tgacatgtga	attgaccaga	cctcagtagc	ttttccaact	ccttaataaa	tgaatctaaa	48480
tattttaaaa	atctggccag	gcgcggtggc	tcacgcctgt	aatcccagca	ctttgggagg	48540
ccagggctgg	cagatcactt	gaggtcagga	gttcgagacc	agcctggcca	acatagttaa	48600
accccatctc	tactgaaaat	acaaaaatta	gcaggacgtc	ttgggtggga	cctgtaatcc	48660
cagctgcttg	ggaggctgag	gcaggagaat	tgcttaaac	tagcaggcgg	aggttgagc	48720
gagccaagat	cgcgcaactg	cactcaagcc	tgggtgataa	agctagactc	agtgtcaaat	48780
aataataata	ataataataa	taataataat	atagataaat	ataaaaaatt	ctttttttga	48840
catggagata	ggatttatat	cacataaaat	ttgcatacag	tttttagtat	caccatcaat	48900
atgtgacagg	gtgatacatt	tgaactttta	agaggacata	ctctacagag	tcaaaatggc	48960
agtattttta	aaagaaagaa	aatgacagct	gcagtagttt	acttgagttg	agttcttttg	49020
gatttatata	gacttattta	tattttat	tagagtttaa	agataacttt	tgtctcttaa	49080
gcatgatatt	aaaacaattc	ttagaatact	tgtcatttcc	tgaattaggt	aatattcatt	49140
tcacctctaa	gcaatttgaa	gaggaaaaga	aaaacagcca	gaaaaaaagt	gatagttgtt	49200
ggttatgtgg	gaaagaatca	aagataat	gtttatgttc	tgtaaaattc	agtttataca	49260
tatttgcaaa	aaatcctctt	caggagtcag	acatagtttt	ttctctaata	tttatgaatg	49320
tcataattat	cacttaaaat	tagccattat	gattttaaaa	actaaaaatg	aaaagaaaaa	49380
ttaatgtaga	gtctgtgtaa	atttagcaaa	ctttgttg	taagttatca	aataggtggg	49440
agttataaat	gggcttagta	ctcatttgta	gggtcaataga	aataatacaa	atcagtgcca	49500
agatttttaa	attcagaggc	caaattatta	ttattattgt	ggaaagtac	ttttttgttc	49560
cttagaaagt	gactatacag	gcaaactctt	ggttttgtca	gggtgtgatga	tgtgtgcctg	49620
tagtcccggc	taaggcagga	agatcacttg	agcgcaagtg	tttgagacca	gtttgggtcc	49680
ctgtctcaaa	aaacaaagca	aacaaagaaa	caacaaaaaa	caacaaactt	tgggtctagga	49740
ggcaaggtac	tatttatcag	tttcaagcaa	acaaactctt	ctagagatat	ttttggtaac	49800
ccagtgtaat	aaaagattga	ctgttagtaa	ctattttcat	ttactagact	gccctgaact	49860
actcatctgt	gagttgagta	tttttttttt	taagtgaag	catgtttatt	aagaaaaata	49920
aggaataggc	tgggtgcggt	ggctcacccc	tgtaatccca	gcactttggg	aggccaaggt	49980
gggtggatca	cctgaagtca	ggagttcaag	accagccggg	ccaacatggg	gaaaccccat	50040
ctctaccaaa	aatgcaaaaa	ttaggttaggc	gtggtggcag	gtgcctgtag	tcccagctac	50100
tcggggaggct	gaggcaggag	aattacttga	acccaagagg	tggaggttgc	agtgagccga	50160
gatcggggac	tccagcctgg	gcgacagagc	aagattctgt	ctcaaaaaaa	aaagaaacaa	50220
agaaaagaa	taaagcaata	agaatggcta	cttcataggg	agagcagctg	agtattttct	50280

tttcccagca	acttttatct	cccgaataga	aattctttct	aatcccactt	actatttggg	50340
attctaaatc	caacattgct	ttcccagaga	tccatgttat	caagatttgc	ttgtattagt	50400
taagagcgat	tgcttattat	ctatacagcc	cttgaaatta	cattgttgtg	aagtcaggct	50460
gctgttttag	ggctaagtaa	ggttttctga	aaaagaggca	atggtgcaaa	gacagctgga	50520
aagacagagt	taaacatacc	ttttaaagtg	gatcaatttt	atgtaaccag	agtatgatct	50580
ttctgtttta	tgtgtttatt	ggaagagttg	tcacttaaag	tgatattcct	caaatagcta	50640
aaaatgtcct	gggttttcta	gatagaaaat	atctctaagt	tgtacttatt	aaaatgtgac	50700
ttgggggaatt	ttaaattata	ttcagtgga	gtatatttgc	ccagctttat	ttttgttcgg	50760
gtaattttatt	tctcaaaatt	ttggaggtga	ggctcttatt	gatttgtgtg	tgtgaaaaac	50820
aagacaaaac	acaccacta	tgaggatcct	ttaaaaaaa	agctttcttg	ttatccagat	50880
aacacacttt	aatcgtacga	cggaaatgga	tacatttaac	atgacaaaa	atagcaaatc	50940
ctctgctcat	ggaaagatga	tttccaaacc	ctctgtctt	acagaagaaa	acatttgggt	51000
gactaagaac	tgacatggtc	tgctcatagg	gatcttagc	atttttaaca	tcatccctgt	51060
ttgactgctt	ctttttctac	attctcatct	tctaactttt	ttctagggct	gctcttttta	51120
gtaaagctaa	acaaaacctg	gcgtcctaca	gccttactaa	cctttgttaa	tccgcctgct	51180
cgtcacatct	ttgtatggca	tctgtgctca	ctcatggggg	aaagagattc	tggagaagc	51240
aacagcaata	accaaggcaa	atgtggggaa	gtgtctgcat	gggttgttag	catgatgtgt	51300
ggggcctagg	ttttacttca	ttttcttcga	atagttaggt	cgtaaagtca	cgccatttga	51360
ataattttgt	atgaccatag	ggtgacatac	tcactccctgt	tgtcaattct	cctaagaaat	51420
ttggtggcaa	atttcacac	ttaagccaga	aagtggaaat	ttggattagg	gatgccaaat	51480
aagcaaactct	ttatattgat	agaaatgtac	ataggcgat	gtcttaatag	acttccacat	51540
tattgagggg	tttttttttg	ttttgttttg	tttttttagt	tttttggct	tatttttaaa	51600
tttggttaag	gcttttttgc	ttgctttttc	ttcgatttaa	ttttcctttt	ttacatttta	51660
attttggtta	ttttggggcc	attttttgat	tttgtttttc	caaaggacgc	ggattctgat	51720
agcggggacg	attctgacaa	gaggtcgtgt	gaagagagct	ggaaactgat	tacttctctg	51780
agagaaaagc	tacctcccag	caagttgcaa	accattgtta	aaaaatgtgg	cctcccaagc	51840
agtgggaaga	aacgtgaacc	aattaaaatg	tatcagatac	cccaaagaag	gcgcttgagt	51900
aaagattcca	agtgggtcac	aatctcagat	cttaaaaattc	aggctgtcaa	agagatttgc	51960
tatgaggttg	ctctcaatga	cttcaggcac	agtcggcagg	agattgaagc	cctggccatt	52020
gtcaagatga	aggagctttg	tgccatgtat	ggcaagaaag	accccaatga	gcgggactcc	52080
tggagggcag	tggccaggga	cgtctgggat	accgtcgggtg	ttggggatga	gaagatcgaa	52140
gacgtcatgg	ccactgggaa	aggcagcact	gatgtagatg	acctcaaggt	tcatatagac	52200
aagctggaag	atattttgca	agaagtcaaa	aagcaaaata	acatgaaaag	cgaggagata	52260
aaagtcttaa	gaaataaaat	gctcaaaatg	gaaaaagtct	tgccactgat	cggatctcag	52320
gaacagaaaa	gccaggaag	ccacaaagca	aaggagcctg	ttggtgctgg	tggttagtgc	52380
acctctgaga	ataatgtaag	taaaggagac	aatggagaac	ttgcaaaaga	agaacgtgtt	52440
tcccagctga	tgaatgggga	tccagctttt	agacgtggac	gtctgcgctg	gatgaggcaa	52500
gagcaaattc	ggtttaagaa	cttgcaacag	caggagataa	caaagcagct	tcgtcggcag	52560
aatgtacctc	ataggttcat	ccctcctgag	aaccggaagc	cccgttccc	ctttaagagc	52620
aaccctaacc	acagaaactc	ttggagtcct	gggacacata	tcatcataac	agaagatgag	52680
gttatagagc	ttaggattcc	aaaagacgat	gaagcaagga	aagggaataa	agaagagagc	52740
caagaaaaag	ggggtaaaag	agcttttaag	gatccccagt	ttccatgggg	ctctcaagga	52800
atgagaagtc	aagatcacat	ccaagttagc	aagcagcaca	ttaataatca	gcaacagcca	52860
cctcaactac	gttggaaga	caattctctc	aataatggcc	agccgaaaag	tacgcgctgc	52920
caggcatctg	cctccgcgga	gtcattaaac	tcccacagtg	gtcaccacac	tgctgatgta	52980
cagactttcc	aggcaaagcg	ccatattcat	caacaccgtc	agtcttactg	taattataac	53040
actggaggtc	agttagaggg	caatgcagcc	acttccatc	agaagcagac	tgacaaaacc	53100
agccactgta	gccagtttgt	gacacctccg	cggatgagga	gacagttctc	agcacccaat	53160
ctcaaagctg	gtcgagaaac	cacagtataa	atcagttact	ggacaaactt	gaaatcatgg	53220
tggagaagaa	agacagtgtt	agctcatgat	ttgatttggg	tctacctttg	gccttgagtt	53280
cttattattt	acattataaa	tattaactgg	ttttatattg	ttaagacaaa	acactggtaa	53340
aagtttcaac	acctcccttt	tgcttgata	ccataaatgg	gcagtttctg	aaattttgga	53400
taaagcatca	agaactcctt	tttctgaaac	gttcctcctt	tttagtgcc	taattaatat	53460
acttacttac	acagacttgt	cccatcttga	tgtaaagttg	ttatgttttt	ataatgccta	53520
taaattaatc	tgacatccag	aaagacctgc	ctcttagttt	atgcagacat	tcaaaaggag	53580
aatttgataa	gcagaattaa	atgtctgtta	ttcataaatg	ttagaaattg	gtttaaaaat	53640
aataacatgc	aactgggtgg	gcacagcttt	ggattttctc	tccacgtctg	ttcccttctt	53700
taaatatgat	tagaattttc	gtaagccaaa	actaaaggtc	ttcaaacata	tcttgcaata	53760
catagctttg	catcttccac	acaacttcca	cgtcttcttt	tattctgtat	tttcaaaaa	53820
actcatataa	tggttttttc	ttttgaatgt	atcagaatat	gtatgaaatt	tgcatatca	53880
gaatctataa	attggcactg	tgtctgggat	tctgtttgtg	gtttgcattt	tgaagttcaa	53940
tgtatcattt	gacacacatg	cacacaaagt	gaactattca	ccaaaatcat	gccaacagaa	54000
ataccaacaa	aaaacttgtg	tagcttgtaa	ggaggttatt	tggcttgaga	gggaagaggt	54060

gggacagaca	tgggggacaa	agctacagca	tatggctctg	tgggtgctgct	tttttccaat	54120
agttgatgaa	atagtgaat	aaaagttaac	tttggttggg	cttggaataa	aggagatgat	54180
acataataaa	ctattctttg	gttagcatta	gtaatgcttg	tagacactca	aacaaggaa	54240
gcgaagtcct	gatgatgatt	tattgtatca	atacctcatt	ttatttggtt	gtggctctcag	54300
tttgccctcac	tggaatcc	actaagaaag	atggatttta	atgggaagaa	aagaattatt	54360
ttctacatcg	ttcctcatct	cttgagtttg	tcaaaaagtt	ttctaagagg	cagtgtgaag	54420
agccagagat	gtccgaaatt	gcctgacttc	atgttttggg	agattcctaa	gtcacatgaa	54480
atcatctatc	aaacacktgt	cttccaaacg	gtgtgaaaaa	ctacagccaa	gatagcagtt	54540
agcattcaag	gaagccttaa	agattgtgat	tatgtttttt	tttcccttgc	gtgtgggcac	54600
tatgctttat	tagaacacat	tatttttaat	catagtattt	tttctttgcc	tctaagaaaa	54660
tactttctta	atgctcagaa	agttgcttcc	aattaatggt	tttttccctt	taaaaagaga	54720
agctttgaga	gatatttttg	ctttcatagc	tagaacagtt	gaagtcttca	actgaggttt	54780
tatagcagat	tagacatggg	taaatgatgt	ctgtaattgg	ttgagttact	gagatgacaa	54840
tctcctgtgc	catttggttt	gaatgtactt	gataggctgc	ttcaaatcag	tcacttcaat	54900
gcattttgtg	taaacccagt	tgtccttttt	tattcctctt	tagacataaa	tgtgctaact	54960
attcctttct	atacacaatt	tattttctaa	gattaaaaat	aataattgct	aggtttggtg	55020
gcacatgcgt	acagccccag	ctacttagga	ggctgaagca	ggaggatccc	tagagcccag	55080
gagttctggg	ctgtggtaca	ctatgccagt	tgggtgtaag	tttggcatca	atatggtgat	55140
gtccctgaag	caggaaacca	caaacttgca	tcaggacggg	tgaacccagc	ccaggttgga	55200
aatggagcga	gccaaaactc	ccaggctgat	cagtaatggg	atcttgccct	tgaatagcca	55260
gtgcactgca	gccaggacaa	cagagtgaga	acctgtcttt	aaaaaaaaaa	aaaaaaaaaa	55320
aaaaaggtaa	tatttattat	attattcagg	tttcaactct	gtagcaaaaa	tgggctctca	55380
tttccctaac	ttgagataac	atagggttag	tccatatatt	ttcattctta	cagtgttctt	55440
ttcatgggag	tgaatgagtt	actctccact	ggtgattagg	taatactgta	gaatgaagag	55500
ttgtataata	tattcattta	cagctgtgga	ttgtggtaag	gactatgtcc	acagtgatat	55560
tccaaagaat	tgggtttata	tttgtgtctc	atctgttaat	cccagggtgc	ctcatgttgc	55620
tgaatatatt	agatagctaa	aatatccctt	aatttcacag	atgaccagga	agaaattaac	55680
caaggtttta	ttgactgcc	tgtatgtccc	atgatgcatt	tctgagcaaa	tgcttatcct	55740
agagaataac	tctgtatgaa	taaaattgct	taattgagtc	tcttactaaa	taagtaacta	55800
gtgccatgct	tttgtgagct	cttggtatgg	cccataattac	tttgtttttt	gtttttgtta	55860
ttgttgtttt	gtgatagtct	tgtctgtcgc	cccaggctgc	agtgcagtgg	cacaatctca	55920
gtcactgca	acctctgcct	cctgggttca	agcaattctc	ctgtctcagc	ctcctgggta	55980
gctgggacta	caggtgcatg	ccaccatgcc	tggctaactt	ttgtattttt	agtagagaca	56040
gggtttcacc	acgttggtca	ggctggcttc	gaattcctaa	cctcaggtga	tccacctgcc	56100
tttgccctccc	aaagtgtgca	gattacaggc	gtgagccacc	gcgcctggcc	tgtttgtttt	56160
tttaacatga	tttttctcta	agcttaata	ccacaaggcc	aaagagaaat	ggtcataatt	56220
taaaccatta	ttatattggt	gaggtatccc	tagctattat	tatagcaaa	tgggaaaaaa	56280
gtgtttattc	tattgaagtt	atgtaatgat	ccgacattaa	tgggaatata	gaggagtcc	56340
aattaattgg	tataatttca	caaagcggaa	tgggtattcct	tggagagtta	aagacattct	56400
cttttagtaag	tgtaaacca	agggcatttt	ctttattcct	gcttctaatt	ccttctagcc	56460
cagtgaatta	tttctctttt	cactggtaat	gtgatgaatg	ggaattgttt	attacattga	56520
agtgaattga	agtgaccttt	tgtgctttag	gtgcaggttg	acactgaaaa	aaaaacaaaa	56580
crctgaattt	ttcacaccta	tgtctgcatt	aaaggctgtt	ttactaccgg	aagttacata	56640
gacttctctg	agtcagctga	tgtgccccag	tgcccttactg	gtcctttgta	gatttgcctt	56700
aatgatttgt	acaaatgact	gggaggcggg	gatgctgcct	gtgtcctggt	gaaccttaat	56760
gaaggggccc	tcttaggcac	agtgcaaaac	aagcatttgt	cctgtactgt	tagagccaaa	56820
attgtgatga	gcaatactga	taattgtcca	gtttatgtca	tctttcccag	attttaaaa	56880
ctgttctaga	tattcttagc	ttgaaccact	tttgattgtg	aaatgtatta	ggtgttgtcc	56940
cattatttact	gtaaaatgaa	gttttgaatc	tcttgtttaa	taaactgtgg	atttccccct	57000
tcaattttct	aaacaacaac	aaaaaaatgc	ttgaagattg	tctttgagtg	taagatctgc	57060
cttttcagaa	agggagtgtt	agtttgtaat	gttaaaaaat	aaagacctca	ttcaataaaa	57120
gttgaagtca	tcttttaaga	gtgtgatttc	tctctatgtg	ggaagaggga	aaaggaaagc	57180
aaggtaatgc	taactaaacc	tgggttttgac	ttttatttat	ttgctttttc	atctgaagat	57240
ggttgtaaac	taaatcttct	ttttgatatt	ttctatgtga	acttgattag	ttttaaagct	57300
tttgcttttag	ctaccctttt	ctgtttaaca	aaattctatc	ttaagtgaga	tctttcagcc	57360
ctattttatg	ccacatttca	cttacataaa	atcttttctc	acaaaagatg	aataaggata	57420
ttactttctg	atttgagttt	agtttaacca	tgttttccagt	tttttctcta	catactcttg	57480
gtttgaagtt	cttttaaatg	aaatgattta	ttttccttta	tatgggcaga	aactaaattt	57540
ctaaaaagtt	ttgagtgact	ctgcaggttt	tctttctcct	ttaacctctg	cttttcttca	57600
cccagagtat	tgatggtaca	ctagagagga	aacaggacaa	agcagaatat	ttttctggaa	57660
tggaccacta	gtctcattga	aaatttgga	ctgacattat	tttttaattg	ttaaaaacaa	57720
cataatcacg	catcatggga	tagaatacaa	aatccacatg	agtttttaat	atgaaatctg	57780
aaactttaga	gaagtgttat	gcctgaatcg	gtctgcttcc	agcctcatat	ccttttctt	57840

caggtacttt	gtaatccttt	gctgtcggag	aaactgttgg	aaaagttgca	atgtccctac	57900
cgtttatagg	gtgacacaga	agattacagt	gtgatatgtg	tagtgacact	gtgctgatga	57960
gactgtggcc	acacagtgct	gctgggtggg	agaagcgaag	ttaagtgttg	tcaacaactg	58020
gggtgtcatc	ttgggtgtgg	atattttcct	tccagctatt	gcttgctaata	tataatctgc	58080
cattaggagc	caagccttta	atgtctaaaa	cagcctaacc	ggaaaactca	gaatgggtgat	58140
attgatttta	ctgtgaatat	caccttctca	gaggtgcctt	aattgtacca	ccatcatctc	58200
tcacgcgagag	taccctagct	tgtctctttg	cttctattct	tgacccttat	aatccaatgt	58260
ccacatagca	accaagatca	tcttttaaaa	acataaatca	tatcttgtca	gtgtccctcag	58320
gggctttcca	ttgcatgaaa	gaatcttagt	cttactgtag	tctccaaggc	tttacetgtt	58380
ttggctgctt	acctaccact	cttaacactg	tcactctcac	accacagaca	ctctgccctt	58440
ttttttgttc	cttgatatatg	ccaagtttct	tcccacctct	gggctcttgc	acttctcaag	58500
cctggagccc	atttccatca	tgtttacctc	cccttccaga	cttgagctcc	tcagctcttt	58560
ctgccagccc	caccacaagt	tgactcactg	ttttaataat	gatggcctgt	aattatcttt	58620
tctttgtatt	tgtttgggtc	ttccacgaga	atgtaaactc	acctagaata	aatgccgtat	58680
ctgtcttatt	cacttcgcca	gtcatataaa	tcataaacct	ggaagagtgg	ttctcaaat	58740
ttaatgtcag	ttagaatttt	gggtgtttat	gagaacacag	actattgggc	ccacctcag	58800
agtttctgat	tcactaggtc	tggggcgaga	cctgataaat	ttgcatacct	cataaattcc	58860
caggtgattc	tgatgctgct	ggtccagtaa	acacctttga	taaccagtga	tctggaatgg	58920
gggctctggt	aagcattttc	tgtaaaacac	tagatagtaa	atattttaga	cttagtgagc	58980
catgtggtct	tttttgtgaa	aactgttcaa	ctctattgtt	gtagtacgaa	agcagccata	59040
gacaatatgg	aaatgaacaa	gggtggccgt	gttccagtcg	aactgtgttt	acagaaacag	59100
gcagtgggct	ggatctagcc	cgaaggccat	agtttgccga	ccccgattag	aagatagcta	59160
ctttgataaa	tatttatcta	aaagatcagt	atgtgactgt	gtatgtaatc	tagaagacag	59220
atacatgcca	gttgctggga	tgagtggtaa	gtgacaaatg	tgtggtccct	ggcctcataa	59280
agttctcaga	gttatggagg	gagacagctg	taataaccac	accagtagtg	agtcagtaca	59340
tttgtgacaa	gtgccatgat	ggaaaaacac	aaggtgctct	gaaaggggtt	aacgggtgcc	59400
tttggaatg	aaaagtggcc	tctcattgct	cttagctgac	aatgtttttg	tttactgag	59460
gaaaaaaagc	aatcagaaga	aactttgttt	tcttcccttg	tctagttttg	cctgcatcat	59520
ccatccatcc	ctgtactggg	ctttccctcc	tgttagtgga	taaactgtcc	ttgatccctt	59580
cctagcccag	cccttgctcc	ctgcattcca	tgccctctca	agaacttggt	gaacatttac	59640
ccccttctct	tttacetcat	cagtttctct	cctggattat	tccctgccagc	atgcaaacat	59700
gtcaccagtt	gcaaaaagttt	tctgtactta	ttacgtctca	cttgccctacc	tccattcttt	59760
tttttttttt	ttttgagata	gagtctcact	ctgtcaccca	ggctggagtg	cagtggcacg	59820
atctgagctt	actgcaacct	ctaccttccg	ggttcaagta	attctcctgc	ctcctgagta	59880
gctgggacta	caggtgaacg	ccaccacacc	gagaattttt	gtatttttag	tagagacgga	59940
cggggtctca	ccatggtggc	taggctggtc	ttgaactcct	gacctcaggt	gatccacctg	60000
tctcagcctc	ccaaagtgtc	gggattacgg	gtgtgagcca	ccgtccagg	ccccattttt	60060
tcttttcatt	atttctctatc	ctcattcttt	gctggaactc	atctagtcag	actcttgttc	60120
tcccctcacc	accaaagctg	ctcttatcag	agtcacaaaa	gctcttcttg	atgatagatt	60180
cagtgcattt	tccagtcctc	atcttactga	atgtctctgg	catttgacac	agttgtctgc	60240
tcttcatggg	acactttctt	ctcttggtct	cttttccctg	aacctttcca	gctgtcccat	60300
catgtctgtg	ggacacagac	cttggtctat	tctatrtggt	tctctgttcc	cccttccatt	60360
tttgactgta	tgatgaagat	aatgataact	catataattga	gcatgtatta	catgtattaa	60420
tacacataaa	cctttttacat	gttttgactc	attgagtcct	cacaacacyc	ctcttctgtg	60480
aggtaggtat	tgtttattcc	cttttcacat	actttcccag	tgtaacacgg	ctactcagcg	60540
ggagagccag	agccagtcct	agggagtcct	tctgtctgca	aacctctgcc	ctgcctcgcc	60600
ttctgctgac	tctcaggtgt	cctctctctc	cagcctgggc	ccctaccctg	agctctagat	60660
gctgatgata	cacaactact	gcctgacatc	caaatacactg	gcaggcacct	tacattttaa	60720
tcatttgaac	tggagcagct	agctcaggcc	acaaatccaa	gtcaccttta	gtttctcttt	60780
ttcttcacat	cccatgatca	gatcatcagt	agatcctggt	tgtctctcct	gtaaaacata	60840
tctgactcc	atctcttttc	accacctttg	ctgtcaataa	cctgggtccaa	gccaccacag	60900
tgtgtggacag	tgtgttgaca	gcctcatgac	cattcttctt	atttccatag	ttctcctga	60960
gtcccacaga	gaataatgtt	tttaaaacat	atcccagctc	taaaccatcc	agcagtttcc	61020
attagaataa	tatccccaga	ccttaccatg	gtctataagg	ccatctgtac	tccatttgct	61080
acctgattac	atctgatgca	tctcctaccc	gtctccctac	tctgccattc	catccacaca	61140
agttttcttg	cttttctctga	aatatgtggt	gcccccttcc	atttcaggac	cttggcactt	61200
gctgttacta	tgaatttgag	tagaataacc	cctgcccagg	acctctgcct	gcttaacctc	61260
ctcagtcctag	aagtctgtct	cccattcttc	tccctggtag	tctctaaccc	tatcttttate	61320
tttcttcata	ctagtactcg	ccactaaaat	ccatctatat	tttgtttact	ctttatcatc	61380
tctgtctcct	ttgtcaatgg	aaccttgtc	ttattgacca	ctagcatttg	gcacatagta	61440
ggagcacaaa	aggtatttgc	tgagtggatg	gatggataat	ggtttaaaaa	tcatgctctg	61500
gattgcctgg	gctcaagtct	catttttagc	atttgcttaa	cctctttctt	tctctttaac	61560
atgaggctaa	ctatatagta	ccatctttat	ttttttgttt	tgttttgttt	tttgtttttt	61620

tattgagaca	gagtcttgc	ctgtcaccta	ggctggagtg	tgtgacacga	tcttggctca	61680
ctgcaacctc	tgctgccctg	gttcaagcga	ttcttgggc	tcagcctccc	aagtagctga	61740
gactacagcg	acacgccacc	acgccctact	tatttttttg	tatttttagt	agagatgagt	61800
tttgccatgg	tgaccagact	ggtttcaaac	tcccagacct	aggtgatccg	tccaccttgg	61860
cctcccaaaag	ttctgggatt	acaggcatga	gccaccacgc	ctagcctagt	accatcttta	61920
taaggttatt	gagaagattg	taataattca	tgtttaacat	taagaacact	tcctgataca	61980
aaataagctt	aaaaaccggg	gtagataggt	aggagagaac	attctagtag	agagaacagc	62040
cagtgcgtgtg	cctttataaa	tatggggcat	atthtgaact	agttaggatt	tttttttttt	62100
ttttttttga	gatggagtct	cgctctgtcg	cccaggctgg	agtgcagtg	cgcaatctcg	62160
gctcactgca	agctccgcct	cctgggttca	caccattctc	ctgcctcagg	ctcccagata	62220
gctgggacta	caggtgcccc	ccaccacacc	cggctaattt	ttttagtatt	tagtagagac	62280
gggttttcac	cgtgttagcc	aggatggctc	tgatctcctg	acctcgtgat	ctgcccacct	62340
tggtctcca	aagtgttgag	attacaggcg	tgagccaccg	tgcttggcca	ttaggatttc	62400
ttataaacat	aaatgtctac	tgaaaaaatt	attattatta	ttattattac	tgagacagag	62460
tttactctt	gttgcccagg	ccggagtga	gtggcgagat	cttggctcac	tgcaacctct	62520
gcctcctggg	ttcaagcagt	tctcctgtct	agccttccga	gtagctggga	ttacggcatg	62580
tgccaccatg	cctagcta	ttttgtatca	ttagtagaga	tggtgtttca	ccagtcagcc	62640
aggctggctc	cgaactccta	actcaccgcg	ctcggcctcc	caaagtgtct	ggattgcagg	62700
catgtccac	cgcgtccagc	caaaatata	tttttttaa	tgtgccttcc	gtaattagtt	62760
tttgtcacaa	aatgaacaga	acgatagtat	gtgaatctca	ttaattagcc	atttgtgtg	62820
ggaaatggtg	atgatataca	gtgtcctttt	taaaagctcg	taagtagcct	ggtttgggga	62880
gggggccaat	tcaatagatg	atgttgatat	tttgactatc	aattgtggtc	aaaatgcagt	62940
taaagttttt	tggtttgttg	ttttgttgt	tggttttgag	atgggtgtct	gctgtgtcct	63000
ccaggctgga	gtggagtggc	acagtcttgg	ctcactgcaa	cctccacttc	ctgggtctca	63060
gtgatectcc	caccttagcg	tcccagtaga	ctgggactac	atatgccctc	cactaagtcc	63120
ggctaatttt	tggtttttta	gcagaaacag	ggtctcacta	tggtgcccgg	actgggtctca	63180
aactcttag	ctcaagtgat	ccaccgcct	ctgcctccca	gagtgctggg	attacaggca	63240
tggtgccacca	cacctggccc	agttaaactt	tttagattca	tagggctatg	ggaaatcttt	63300
tcatgttagt	tttggttact	aaagcacagc	cacagctgtt	aggggaaggtc	atcctcatat	63360
cttaactgtc	agagtctgag	aattttgtat	tcattggtag	tttgaccagc	ctgtcaactt	63420
attcttgaag	aaatcctcaa	gataaagatt	aacatttttt	ttctacctga	tgatttttag	63480
gtaaagcagc	agctttctct	ttttgtcata	ataattcatg	taatgtatag	tttttccccg	63540
actccccagc	tttttgagac	atttgaaaca	gagaagctga	aaaaaaaaact	ataatgaact	63600
tttgtatacc	catacttata	tatttttctc	gaaattgtg	aaaattatag	atatcatggc	63660
agtttacc	aaataacttc	atgcttatct	cttaaaatca	agaatattct	tttttttttt	63720
tttttttttt	ttttaagacg	gagtctcact	ctgttgccca	ggctagagca	atggtgcaat	63780
ctcagctcac	tgcaaaactac	acctcccg	ttgaagtgat	tctccgcct	cagcctccca	63840
agtagctggg	actacaggca	cctgctatca	tgcccagcta	atthtgtgtt	ttttagagag	63900
tggtgtttca	ccatgttggc	caggctggtc	ttgaactcct	gacctcaggt	gatctgcctg	63960
cctcggcctc	ccaaagtgtc	gggattacag	gcattagcca	ctgcatccag	ccaggaatat	64020
tctcctaata	ccactatccc	atctaggaaa	tttaatactg	attcaatatt	gattcaatat	64080
tatatcatgt	acagtccatt	ttcaaaattc	gttaattaaa	tcaaaaatta	ggaaagagag	64140
tggttttaac	caagatccag	tcaaggttta	tatatthcat	ttggttatga	tttttttaaat	64200
ctttcattat	actgggtatt	ttgaattgtc	taggccacat	tctgatttgt	ctgattattt	64260
cctcagttat	agattcaagt	taaatctttt	tgtaagaat	gctataaagg	ggccgggtgc	64320
agtggctcac	gcctctaate	ccagcgcttt	gggaggccga	gacgggtgga	tcacctgagg	64380
tcaggagttc	aagaccagcc	tgccaacat	ggtaaaaccc	cctctctatt	aaaaatacaa	64440
aatttaactg	ggcatggtgg	tacacacctg	tagtcccagc	tacttgggag	gctgaggcag	64500
gagaattgct	tgaacctggg	aggcagaggt	tatagttagc	tgagattgtg	ccactgcact	64560
ccagcctggg	tgacagagca	aaactccatc	tcaaaaaaaa	aaaaaaaaaa	aagctatgaa	64620
ggggccaggc	acgttggctc	actctttagt	tcccagctac	ttgggaggct	gaggcaggag	64680
gatcacttga	gcctgagagg	tcaaggttat	agttagctat	gattgtacca	ttgcacccca	64740
gcctgggtga	cagagggaga	ccctttctcc	aaaaaaaaagg	aaaaagattg	ctataatggt	64800
gatgttgtgt	acttctgtt	gcatacacat	agtagacata	aaatgtcagt	ttgtcctatt	64860
atcaataatg	ttaaagtttg	gtaaatttgg	ttaatgtgat	gtttcccagc	tctctgcatt	64920
gtaaatatat	catttccctt	cttaaataga	tatctgggca	tggaacttgg	atcgtatgca	64980
tatcctgttt	cccatgcata	ttttactcaa	ttatttagc	ttttgttgt	ttgtgtctat	65040
tggtgttgtt	tttaagatgg	ggtctcttgc	tctgtagccc	aggctggagt	acagtgccac	65100
gatctgggct	cactgcaccc	tccgcctccc	gggtcaagt	gatcctccta	cctcagcccc	65160
tctagtagct	gggactacag	gtgtgtgcca	tcacacctgg	ataatttttg	tattcttttg	65220
tagacatggg	gtttggccat	gttgcccagg	ccagctctgga	attcctgggc	tcaagcagtt	65280
tgctgcctc	agcctcccaa	agtgtctggga	ttataggcat	tgagtcactg	tgcttggcct	65340
atthtagtat	ttattatcct	ttcttgattc	agttattaca	tctggggttg	caatatggtg	65400

atcccccaat	tgccatttct	tctgcattta	ttagccacct	tttttttgtt	tttttgtttt	65460
tcagatggga	tcttgctgtg	ttgcccaggc	tggagtggct	attcacaggc	ctgatcatag	65520
cgcactacag	ccttcaactc	ctgggctcca	gggatcccag	gggctgggca	tggtggctca	65580
cacctgtaat	cccagcactt	tgggagataa	tttaactttc	tcagaattag	aaggctagta	65640
agaaaaagtg	ccaggacccc	aaacccaagc	ctcttttagcc	tcgagtagct	gaaatgaaca	65700
ggtgcatgcc	ccatgcctgg	taattagcta	gcttttaaaa	aatttttatt	taaagaagag	65760
ttcccccttt	ttctctcctt	ctttcagtat	caccatggaa	tcatagaattc	tttctatatt	65820
caatgtactg	tgctattttt	ctggtgattt	attggccagt	gggatcctgt	gtccttttga	65880
aatggcccca	gtggtgtttg	agttcttcca	tcctttctga	cacaaaaatat	ttcagactca	65940
ccttgacttt	tccttgcttc	agacctggaa	gcagctattt	ttttcaaggga	accctgggttc	66000
ctttcagtg	cagtggtact	tagaaaccaa	gatctaggtg	ctaagtatgc	ttaaacaata	66060
accttttata	aaaaaacatt	gcctgagggc	aggcgactg	gctcacatct	ataatcccag	66120
cactttggga	ggccgaggtg	ggcgatcac	gaggtaaagg	gttcgagacc	agcctggcca	66180
acatagtga	accctgtctc	tactaaaaat	acaaaaaatt	agccaggcac	ggtggcgtgc	66240
acctgtagt	ccagctactt	gggaggctaa	ggcaggagaa	tcgcttgaac	ctgggaggca	66300
gaggtgtgg	tgagccgaga	tcgcatcact	gcactccaac	ctgggtgaca	gagcgagact	66360
ccacctcaaa	aaaaaaaaaa	attgcctgac	attttgat	tgattacca	tgtaagagg	66420
ctttcacgtg	tttttaaaat	ggttcactct	ttggtccttc	ctctagtact	gtgtgggtag	66480
gcaggtcaaa	tgctaaaatc	tttattttaa	gatgaagaag	gtgaagttaa	gagataattt	66540
aactttctca	gaattagaag	gctagtaaga	acaagtgcc	ggaccccaaa	cccaagcctc	66600
ttaccagttt	aatttttttt	tctttttttt	cataacatgt	agtctcctaa	aaaagtgtgt	66660
ttagatatct	agtaaagcta	tgtagttgg	tcgtggtcct	ggtcactgcc	tgcccactta	66720
aagagaacac	ttcctttgtg	agaataactg	acaaacgacc	agaaccatag	aggatatttg	66780
catccttctg	agcattgctt	gggagctcct	tttgtaatgg	agtttgctaa	atggctttca	66840
gcagtgtctt	atgatgcagt	gaatgtttga	tgtgggactt	ggagagctct	gtttggaac	66900
agtggatagt	catacttata	ccctttgtct	catgacagcc	tgtaattggg	tcactgctag	66960
ggaacccaaa	gaagcagggc	tggtgtcgg	ggctcacgcc	tgtaatccca	gcactttggg	67020
aggccaaggt	gggcagatcg	tgaggtcagg	agttcgagat	cagcctggcc	aacatggtga	67080
aaacccatct	ctactaaaat	ataaaaatta	gtcgggcagt	gtggcgggca	cctgtagtcc	67140
cagctactga	ggaagctgag	gcaggagaat	cgcttgaacc	cgggaggtgg	atgttgcatg	67200
gagccaagat	tgccgcatg	cactccagcc	tggaacaata	tagagtggga	ctccatctca	67260
aaaaagaagg	aggaaaccct	ggcagtaaat	cttaaaatgc	tatgcttatg	ctgtaattta	67320
gtattgcgct	gtttgaaggt	ctctttctta	accctactta	ttttattttc	actaaaattc	67380
aagtttttca	aattaaaca	ctaaggaaac	tttttagaca	tttgtcccct	attctctgcc	67440
caccatgaga	gtttaatacc	tcagatatat	tacatatctg	tatactgatt	atatttgtac	67500
ttcatatatt	aaaagactaa	gatttttttc	atcctctgag	aaccattttt	tgcccccttt	67560
ggggtgat	tacccttatt	gaaaaacgca	tggagtagat	ggaaaaatga	aggatacatc	67620
tgatttgtat	tacaataatc	cttttttttt	ttttttttga	gacagagtct	cgctctgtct	67680
ccaggctgga	gtgcagtggc	gcgatctctg	ttcacttcaa	cctctgcctc	ccgggttcaa	67740
gtgattctcc	tgccctcatc	tcaccaagtag	ctgggattat	aggcacgtgc	caccatggcc	67800
agctaatttt	tgtattttta	gtagagacag	ggtttctcca	tggttggtcag	gctggtctcg	67860
aactcccgac	ctcaggtgat	cacccacctc	ggcctcccaa	agtgtctggga	ttacaggtgt	67920
gagccactgc	accgggtcag	taatccatct	tatagttgca	tattgaaaat	aaagtgtctt	67980
acttgaataa	ctaaatgcag	tctataaatt	atttcagctc	taatagcact	atgagctttt	68040
ggttacctga	aggctaaact	catcctactg	atcaaagata	gtaatttttc	acaaagagga	68100
agcatcaaa	tctacaatct	taattttttt	tccttccaaa	gagttagata	attgagattt	68160
ttgatgatac	tggtgaattt	aatctttttc	aatattgttt	ttctctggaa	tttattgata	68220
tataagatat	tacttgaata	tattctaacg	cttaaggtat	acacgtgtta	tttcacctaa	68280
aaattccccc	tttctcttac	cactacccca	gatgaagaac	tcttgagtct	ttgagggagt	68340
agaaacatac	tgcatgtagt	tttttctttt	taacattttac	ccattacaga	ccatcttgtg	68400
tgctttgtgg	gtttttttcc	taaacatttc	cctgtatgtc	tttttattat	tgtttctctg	68460
ctttgtctct	tttcttctgc	ctttgctttc	tctacttccc	ccaatatttc	ttccatttct	68520
ctttccctgc	cctcttattt	cttccctctc	ctacatgtta	tctcctttct	tttcattcag	68580
gactatgaga	gtaaaattgca	ggccttgca	aagcaggttg	aaacccgatc	tctggctgca	68640
gaaacaactg	aagaggagga	agaagaggaa	gaaggtgaaa	tctagagacc	gaaagtgtcc	68700
tgtgtatata	tttttgaggt	tatattatca	aattagtcac	ttatgcataa	tcaaagctgg	68760
attcctctta	gttggtcccta	tcattttattg	acattttact	gagccagttt	acagatgatt	68820
gaatagatgt	gcagtgttga	gaatccaggc	tgtattttat	gaggtgggga	tgggagtgat	68880
ggtgttctta	gaccttcgca	cggtttctga	gcacgttcat	ttaaatccct	taggaagtgtg	68940
acctgcactg	tctctcatat	tgctgtagat	gtgactttta	gatattccatt	acaaatatag	69000
agctgaatct	ttgttaagac	agtctgttca	atttgaagcc	atttctggca	aatttctaat	69060
catattgctt	agtcatgttt	gcctatctta	ggttttccag	catttcttag	ttgagttagg	69120
gctttcatta	ttacttattg	ccttataatg	cagtatgaac	acttaccaca	attcgttttt	69180

ctaccaacc	tcttggaat	ctcaaggctt	caaactaact	ttgaaggaa	tcttaattta	69240
tgtattacat	ttcttattta	tttattttact	tattttatttt	tgagacagag	ttttgtctct	69300
gtcgcccacg	ctggagtga	atgggtgcagt	cttagctcac	tgcaacctcc	gcctcctgcc	69360
tcctaagtag	ttgggattac	agggatgcgc	caccatgccc	agctaatttt	gtgttttttag	69420
tggagatggg	gtttcaccat	gttggtcaga	ctggtctcaa	actcctgacc	tcagggtgatc	69480
tgcttgccat	ggcctctcaa	agtaatggga	ttacaggcat	gagccactgc	gcccagccaa	69540
tttcttattt	ttaaataagt	gagatggtac	cttttttctc	tctgtaacag	aaaaaaaaaa	69600
agaaatatat	gaagtcatac	aatgtaacag	aggtcaggag	tttgtctttt	tttttttttg	69660
agacagagtc	tcactttgtc	accagggtt	tagtgcagtg	gtgcagtctt	ggctcactgc	69720
aacctctgct	tcctgggttc	aagcaattct	catgcctcag	cctcctgagt	agctgggact	69780
acaggtgtgt	gccaccacgc	ctggctaatt	tttgtaattt	tagtagagat	gtggtttctc	69840
catgttgcc	aggctggcct	caaaccctg	acctcagggt	atccgcctac	ctctgccttc	69900
caaagtgcctg	ggattataga	cgtgagccac	tgtgcctagc	caggagtatt	tttttttatt	69960
gtggccacat	gttatacctt	tttttaaaaa	aaagaaaaat	ttatttcttt	gttgtaactat	70020
tttaataaaa	tacaaaagta	ggatttttaa	agattttctg	aaatgtatat	attgctttca	70080
taatatcaac	agaatttggg	aagctttact	aaaaagaaac	aaaagcagct	gaatggagag	70140
aaactggaaa	acacctcacc	ttggtaaaa	agataagcta	gaagagaggt	gtaattttta	70200
tatagtgtga	agattgccat	taaatatgca	tcattcattc	tttcagttcc	ttggacacag	70260
catgaatttg	agttggccca	atgggccttc	cggaaatgga	agtctcatca	gtttacttca	70320
ttacgggact	tactctgggg	caatgccgtg	tacctaaagg	aggccaatgc	catcagtgtg	70380
gaactgaaaa	agaaggtatg	gagcaggagg	acacaggaga	gctggaggca	aagccgagcc	70440
tgtgtgggt	gcatctgggt	tctcaccttg	aattaacctt	tcctttgggg	gaactcagct	70500
gctttgtgct	ataaaacagc	tttatatatg	tctttattta	atataatgtg	ctggaacata	70560
gtaggtgcta	attaaatttt	tgttaaataa	aatttttaaa	acatgttttc	tagcttattt	70620
tacaaataga	ctatatcct	acaaaaactg	gagaaacata	gtaatagaaa	aatacttctc	70680
ttgggtttct	gaataatttt	aatttttctc	cacttttctg	ttattaagtt	tgacactcta	70740
agtttggtat	gccctttg	cattgatctg	cttctctct	agggtctgag	aggcactgct	70800
gtgtagtga	caagcacatg	caccttgac	ccagacagtc	caggttccag	tcccagctcc	70860
acgagtggg	ttgatcccc	tagccctgtg	cctccattcc	cccatctaca	taatgggggt	70920
ggcagcagta	cctggcacca	agggattttg	tgaggaaatga	aaatgggcat	atgcataaag	70980
cacttatagt	gtctggcatg	gtaaaaggca	ataaatgtta	ttcatttttg	taagtattac	71040
caaaacttca	caccaaacct	tagattcagt	cttctgtatg	ccccctgtgt	tatcagtttc	71100
ttcactgcta	agattacagt	ggaagaccat	attgtgtagt	aggtaagagc	atggaccacc	71160
taactttggg	gaatttaatt	atcttctcct	agaactaaat	tctagtttta	gggaatacga	71220
tatgttcttc	attcataatc	ttggcagcct	ttctcatttt	ataattattt	attagccact	71280
cctctcctca	agtagactaa	ataaaggcag	taactgtctt	tttttcttac	tactgtgcat	71340
agcatttaat	taatattaaa	taattgttta	ataattgttg	gtgaaactat	aacgtggtag	71400
ggagaaataa	atgatgcaat	gtcgtgaagt	acttaccata	tttattaaac	acttttataa	71460
aaccatattt	attaaacact	tttttttatg	agataggatc	tggctctgtc	gcccaggctg	71520
gagtacagtg	gcgcaattat	ggctcactac	tgctccacc	ttctgggctg	ggctcagttt	71580
gtagagacag	ggtctcactg	tgttgcttag	gctgggtctc	aactcctggg	ctcaagtgat	71640
ccttccgcct	cggccttcta	gtgtgctggg	attacagggtg	tgaacttccg	tgcccagcca	71700
aataaatgct	tttaagagca	gtagtgtcaa	tcctctgaat	ttaaaatggg	aaaccttgct	71760
gtgttttcat	cctctctcat	tagtgaggga	ctagatatgc	attttgtggt	cgaccttttg	71820
ttctctggta	taaattctat	aaaagtcttc	attcattcaa	tcaaatgtta	tttataaaga	71880
ttttatgccg	ggtgtggtg	gttgtgctg	tgatctgagc	cgcgagggtg	ggtggatcat	71940
ttgagctcag	gagttcaaga	ccagcctggg	caatgtgggt	aaaccccatc	tctacaaaaa	72000
atttttaaaa	atcagctagg	ctgggtgatg	atgcctgtag	tcccagggtac	tcaggagggt	72060
gagatggtag	gatcacctga	gcctaggagg	tggaggctac	agtgtctgtg	gattgcatca	72120
ctgcacccag	cctgggtgac	agagtgtaac	cttgccccca	cgtgctctcc	cctcaagaag	72180
attttcgtat	gtgcaaaaac	actgccaggc	taggaaaaca	aagtccttaa	atagcttaat	72240
ttgaagttag	ggttattccc	tatgactgct	aacagggtgg	catatttttc	agtagaaagt	72300
tagtcacaag	atatttggtg	agcacattcg	acctctttcc	cattgggtat	gattgtatcg	72360
tctcatctct	gaagcttgct	tgtgaaaac	catttgtcaa	tggtttattc	tttctattca	72420
ggtgcagttt	cagtttggtc	tgtgactga	cacactgtac	tcccccttgc	ctcctgaatt	72480
acttccact	gagatggaaa	aaactcatga	ggacaggcct	ttccctcgca	cagtggtagc	72540
agtagaagtc	caggatttga	agaatggagc	aacacactat	tggcttttgg	agaaactcaa	72600
gtatgaaaac	attcataaag	gctgggtgtt	ttatttagga	aataacaatg	acctgctcaa	72660
gtgagctccc	tccagctctc	ctctctctga	ggacactgtt	gcttacttcc	ctgtactttc	72720
taaatgctgt	tgttgcccaa	tgggtattacc	atccatctgt	tttcttttaa	gtgaataaat	72780
gcactctca	cacactctga	catttgtaga	tatttaagtt	tacctccat	agacctctt	72840
ataaatttta	aatattccag	tttcatcata	tttctgttac	agatttggaa	cttattttatt	72900
ttttctatcc	tggacctcag	gataatcaat	aataaacatg	gaagttataa	tctggcttct	72960

tttacaatgta	gtataatgag	gtatttcccc	cgtctcagtg	tatttgagat	gtcacataca	73020
gttgacctac	aatattcccc	cttatgactt	ttcacccggt	agaggatagg	tggccattct	73080
tgacttttga	tggcttatct	ctagaatcaa	gacactttag	atactgagtg	ttggtttctg	73140
tatttttggg	cttcattctt	tttagtaaca	ttaaacagtg	ggagcaactc	cttttttttt	73200
ttttgaagaa	aatcttgaac	ggaagatgct	tccaaagtga	aagtttaatg	caatctcatt	73260
atattaaccc	aatgtgcttt	gtgtttacta	aataggcaga	ggctggattt	gatgcgagag	73320
atgtatgata	gggcagggga	gatggcctcc	agtgcccaag	acgaaagcga	aaccactgtg	73380
actggcagcg	atcccttcta	tgatcggttc	cactggttca	aacttggtgg	gaggtatgtg	73440
atgattttgt	tgatgtcttc	ttttaaaata	atgattagtt	ttaagtgttt	aaaacctgag	73500
cagataatat	aaaaagttaa	gaggaaaaaa	aaaggtgtaa	aaaggcctta	tctataatct	73560
tatccactca	caaccactgc	gtcatittgt	atatttccct	tcagttatat	tcagcaggca	73620
tttttgttct	gactctcaaa	ctttcagcca	aagtatttaa	atcttccctg	actttaaact	73680
tacaaagtgt	tctgtaattg	attaagataa	atttgaagag	aaaagaaac	ttacgcctgt	73740
atcctacaca	gtttaaattc	atagtacagc	ttatcagaca	caatctcttc	gtgatctgat	73800
ttgttttctt	tttttttaaa	aagcaattca	tccactttta	gtcattgcaa	gggtaaatga	73860
attgcacctt	ggtagtcccc	ttcagcctca	tttttttttt	atacattatt	tccagcactt	73920
aatgaaaaca	catagtgcct	ctatgagtc	atattgagtc	aagtgttctt	agaggttttc	73980
aaattttcaa	agcaggggaa	tttaactttg	agcttatgat	ttcagaattt	gaattgtctt	74040
ttcaattgca	gaaaaccctg	tgatgttttt	ttaaagaggct	gggtaaaaag	gagttgtctga	74100
agtcatagta	tgtgtggaa	atgaagtaaa	tcttttccac	tctgtctctga	tagcaaaagta	74160
gtttttcaaa	gagatgaatt	ggccaaacac	atcttaccct	agcaaaagcac	atgctgtctt	74220
gccatatgcc	atttcatcca	ttttggtttc	tgctccaaaa	tgacgtgaga	atttcagtaa	74280
agactgaaag	ccttcttgtt	tcactttgat	gttgagtggg	tcctttgagg	cctggatgct	74340
gatcttttat	gtgaactgga	gaatgtaaac	gtgttatagc	acagtattta	cctaggatgg	74400
tctactcacg	tgtttccctt	gcttttctta	ttctaaggat	tagtccttac	ataacttttt	74460
caacttgccc	ttgttccctt	tcacctgacc	caaagtattt	atgcttattt	tatccttttg	74520
cttttccatt	tgcttttatt	gtgttgattc	cccagtggtt	tctgtcctgt	tagcacctat	74580
tgcttctctg	ttaaccaact	attcctctct	ccttggtctg	gttaattggc	gtcttacctg	74640
gtgtctagct	cccccathtt	ccacggctgt	gtgaacgagc	gccttgccga	ccgcacaccc	74700
tcccccaact	tttccacggc	cgattccgac	atcaactgagc	tggtctgacga	gcagcaagat	74760
gagatggagg	attttgatga	tgaggcattc	gtggatgacg	ccggctctga	cgccaggagc	74820
gaggagggat	cagatctctt	cagtgcaggg	catgacccgt	tttacgaccg	atccccttgg	74880
ttcattttag	tgggaagggt	ggtagaggta	ttgtgagaaa	ggcgaaaagg	gaccagctct	74940
tgctctgaag	gcctccctgc	ttgcacaatt	ttggataacc	ttgcattagc	caattcaact	75000
catgaatgct	ctttttcaag	ttctccaata	cttcaagctc	ttagtcagtg	ctggtctggc	75060
tgagatggtt	ctcaggcatt	gtgtgtgagg	tcctcagctc	actcagcaca	atacgtagca	75120
atatatgaag	actgcatgga	agacaccttg	tcttaaatgt	cccagtaatt	ttgttctgaa	75180
gaagggaacc	ccaaaggaga	aagtaaagct	aagagtcaaa	cgtagtggtg	tcagtgaata	75240
gtattatggt	ctagatgttt	ccttttttaa	ggattgtgat	tgaaaaagca	tatggaggta	75300
acatgagaga	gggggcagag	tgaggccaag	gtttggctga	agaattttgtg	gtactgcagg	75360
agtcagagga	caaaagtaga	ttcaggagta	gagtttaaatt	tattttttgc	ttagtctact	75420
gaatcctaaa	agaatcagg	aaagttaaag	ttctccttta	cctaaaacgg	atgcctgtct	75480
ttggccagtt	tcaactgagc	tatctacgga	tctttggcat	ctgctaaaaat	tttaacctat	75540
agggtttata	atgataaagc	tcagaaaaatc	atctattgac	cccatcagca	gtgaatattg	75600
gccaggcacg	gtggtcacc	cctgtaatcc	tagcactttg	gtcacacact	gtaatcccag	75660
cactttggga	ggctgagggtg	ggtaggataac	ctgaggtcag	gagtttgaga	ccaacgtgac	75720
ccatgaaaac	cgtctctact	aaaaatacaa	aaattaggct	gggtgtggtg	gtcacgctct	75780
gtaatccagc	agtttgggag	gccaaaggcg	ccgggtcacc	tgaggtcagg	agtttgagac	75840
cagcctgccc	aacacgggtga	aaccccgctc	ctactaaaaa	tacaaaaaat	tagccgggca	75900
tggtggcggg	tgctgtaat	cccagctact	tgaggaggctg	aggcaggaga	atcacttgaa	75960
cccggaaagg	agaggttgca	gtgagacaag	attgcacaac	tgactccag	cctgggtgac	76020
aagagcgaaa	ctccgtctca	aaaaaaaaaa	aaaaaaaaatga	atatttgttt	ggtgctatat	76080
gaaactgctt	tgtgttaaga	ctaggtcacc	aaactgaggc	attgtctctg	gctctcagtt	76140
tgtgcatacca	tctttcagtg	atgcagaact	ctctatgcac	atgatcctaa	tactggtgtc	76200
tttgttcac	tccagccttc	atgctgagg	aatttaaatt	acaagttcaa	cagtagagca	76260
tgcatgtaat	taaccaaaaa	gtatgtccag	gtttaaaagt	attaaattta	aatttaattg	76320
tacattagaa	tcacatattg	ttgaatcctt	tataaataag	ggaattatgt	cagtgttaaa	76380
tggtgagta	ccagttatct	aaaaattaaa	ggatagcgag	tctcctgacc	tagtagagac	76440
ataggttcta	aaatttatct	caccagggtgc	agtggctcac	acctgtaatc	tcagcaactc	76500
tgagggtgga	ggccagaagt	ttgagaccag	agtgggcaac	atagttagac	cctgtctctt	76560
tgaaaaattta	aaaaacaatt	agctgggtgt	tgtggtgcat	gcctgtagtc	ccagctcctt	76620
ggaaggctaa	ggcaggagga	taatttgagc	ccaagagttt	gaggctgcag	tgagccatga	76680
ttgcaccact	gcactccagc	ctggggacag	agcaagaccc	caagtctacc	aaaaaatata	76740

aaataaaata	aaacagaatt	tgtagtatgg	tattgggtcac	atgggtcggaa	cctcataact	76800
aaattagaag	atgatcattc	ttatggaaa	ggggaaaaga	aaagcttttt	tgatagaaat	76860
gaaagcctca	agtgtaaactg	taggtcatag	tatttagacag	tctctgggtt	ttccagtggc	76920
aaaggagcat	ttagaattca	gtttttccat	gaataaacat	aaagggatta	tgtggccatg	76980
aatattgttt	tctactggca	tatcgccatg	aatagataac	gtagaatttc	agagaagctg	77040
atttttatta	aaattgagaa	aggatatgtac	aaataatggg	acaataatgc	ttgcttcatt	77100
tgatggctgc	ctgtgttctt	cagcacgttt	ccatcctgtt	ctaagggagt	tggaagttaa	77160
aaccagttt	ctggccataa	actttgtccc	agtgttaatt	acacagtttc	taaaactgtt	77220
cgcttgtgtt	ctataccatt	gcagtgaaga	gtaggcagct	tcattcttacg	taacatgggtg	77280
aggagaagcc	tgacctaat	gtcaggagaa	gccttgggtc	ctaggccatc	tctgtcacta	77340
tgacaagcaa	cttaaagtga	agcctttggg	ctctcaagt	ttaaagtgaat	aacaggtaga	77400
aataaagagc	cataaagatc	tttttcaact	ttcatattct	gtcatctaca	ctagcaaaaa	77460
tgaatatcct	ggtttaagact	tctactaaga	agcatttcta	ttttgaggaa	gttgtgtgga	77520
aactcaagac	aatcccactt	aatgtggaaa	ttactgagg	atacatgtat	tgttttctct	77580
ctagttttat	agttttcttt	cttttaatac	tgctcagatgt	aagtataggt	ataaagcaat	77640
caactctatt	ttgattttca	cataaaatac	ttatggagct	gtggctcagg	aattgatttt	77700
tccatttgtt	taggttccca	agttctaacc	ttgcaagaaa	ggactggggg	aggaagtga	77760
ataatataat	agcaatcagt	aaagtgaagg	atagctttcc	atttttgccc	ttgactacat	77820
ttctcttcta	caaacaaaat	ttacaataat	gcaacatcac	ccagtttgtgt	agtgggtcatt	77880
gtatgtgtaa	ggaaagttaac	tactttcctg	tttttgtgat	tcacaagatt	ggattttctcc	77940
tattggcata	taattgccaa	tataaatcac	ttgtaaaatt	tttagtttct	ggctgggtgc	78000
ggtggctcac	tcccgtaatc	ccagcacttt	gggaggccga	ggtgggcaga	tcagagggtc	78060
aagagttcga	aaccagcctg	gccaacatgg	tgaaacctcg	tctctactaa	gaatacaaat	78120
attagccagg	catggtggtt	tgtgcctgta	atcccagcta	ctcaggaggc	tgaggcagga	78180
gaattgcttg	aaccggggag	ggggaggttg	cagtgcgcgc	agatcgcgcc	actgcacttc	78240
agcctgggca	acagagcaaa	gactctgtct	agggggaaaa	aaaaattagt	ttcttgtgaa	78300
attccattag	taagtttgtt	ttttagccat	gactgcctct	gctaaaaact	tatattgcaa	78360
gtgaaaaaga	gcttgtccct	ttagctattc	tcttatggca	tgctgggatg	ttatgaacca	78420
tccatcataa	tttgggatga	aatgtaaaat	atttaagcta	gtcttctggg	ttaaaagggg	78480
gttaaagatg	atgattctgt	gatctctcta	agtaccttaa	ctgacattct	ttctataagt	78540
agcaccataa	ttttgcta	ttttcttata	tttatgttga	ccagtttgtt	ccactttttg	78600
aagtaacttc	attgaacttt	ctgctaacgt	atcctttcca	tattcccctc	tctcatccct	78660
gttctccttc	agtctgctag	gtcaaagtat	aaggattctc	aaaaatgcag	agtaattatg	78720
aaaattctgt	ttaaaaagct	tcagaataga	atgtttatat	agcattgatt	tgagagctaag	78780
ggctatatcc	tggtaacata	acaagggtct	atctagtga	tgtgtgattg	tgtagtcatg	78840
gagctgatgt	ccctgccata	gaattacaga	attatctaaa	aagggaatc	tataataatg	78900
gccttcaatg	ctcccacact	tgactagacc	attgccatag	gcagaaaggc	aatctcatct	78960
ttactgtgtg	tgcaaacagt	acttttaagt	aatgtgtgct	taataataagc	tttctttaaa	79020
aaaaaaaaag	gtggctcctg	tttttgata	gctattttta	agatagatat	agtttaggaat	79080
ctaaatgtgt	tctatatagt	taatatccat	tatgaggtgg	ctctgaaaaa	tcaacctagt	79140
ataagttgga	tggtttgtct	tttctgtctc	tgttayaacc	catttttcta	gaaaagcttc	79200
cacttctact	taaagataag	aggcaaatca	tttctgtgtt	cccttatgca	catagatgtt	79260
cacttaccaa	atatttatag	agtggcccag	ccctctgcag	acagtgttga	gcaaaaccag	79320
acatggttcc	tgccctcata	gacttatagt	ctgatggatg	acacagacat	aaattaaaata	79380
attgcacaaa	taaatgtaaa	ttatagcagt	gatgaaatag	ttgtgtgtgt	tgctcgagat	79440
cacataacgt	tggtgggacct	gttatgagaa	agtgtgtgtt	gaaataagac	ctgaagatgt	79500
gaaatcaaag	gcgtgaggct	gggtgtggta	gtcacgcct	gtaatcccag	cactttggga	79560
ggctgagggtg	ggtgaatcac	ctgagatcag	gagtttgaga	ccagcctggc	caacatgggtg	79620
aaacccccacc	tcaaaaaaaa	aaaaaaagac	atcaaaggca	tgaggtgtag	gcacagagtg	79680
cagagaggac	tccaggcagg	ggacatgtgc	gatggctctt	taatggccca	ggccttgaaa	79740
tggtgcaagtc	tagctggggt	gcccataata	gaggagggtg	atctgtgtat	gtgttgactt	79800
tgaaagtga	attatccatt	tgatacctta	cgccaaattc	tttgtctttt	aaattctgtt	79860
gcagtagttg	ttgtttgagg	gtcttggtta	gaacttttta	gaacttttca	ttcttacgtt	79920
tttttacacc	agagcttcaa	ctctctgttt	cttagcattt	ctctctctcc	tgctttggag	79980
cagactgaaa	gggaagtggc	attatttcat	ctgatktaat	gttcatttgc	agttattttt	80040
gttatattgtt	tcatttgtat	ttcattttatg	cttagtttat	gttaacattt	gtttttctgt	80100
gatgtattat	gtctgttttc	ataattttca	ttactttgag	ttgaaatccc	aaatacccag	80160
tctatcctcc	tcctctcttt	gttttaaaat	agatttcaga	gcaggaaaag	catcagataa	80220
acatgtatgt	cacttggttt	tgtttgattt	attgatacaa	gtgtctgtct	ttgtcttttg	80280
tcttatgttc	gaagatcagc	atataccaaa	tggaagact	ggaatcatag	taggaactag	80340
attctatttc	cctttcacta	attggaatga	acactctagc	ttgtgacca	ccagttttta	80400
ttgctttttt	aaaaaagtag	tagctacaca	ataaagaaga	aaatacaaat	gtgataggac	80460
tttgctagtt	tgcttgaaat	ataaagtgtt	agagctataa	tttagctttc	acttatgcac	80520

ttctgaatta	ggttttgaac	tctaggagca	aacgcaaacc	ttttaggcat	caactgtcat	80580
cttctccctt	aacattttatc	ttttttcttt	ttttctcttg	agacgaagtc	ttgctcttgt	80640
ccccaggct	ggagtgcatt	ggtgcgatct	cggctcactg	caaccttcac	ctcctgggtt	80700
caagcgattc	tcctgcctcg	gcgccccga	gtagctggga	ttatagggtg	ctgccaccgt	80760
gcctggctaa	ttttgtatt	tttagtagag	atggggtttc	accatggttg	ccaggctggt	80820
ctcgaactcc	tgacctcagg	tgacccgcct	gcctcggcct	cccaaatgc	tgggattaca	80880
gacgtgagcc	accatgccc	gcccttaaca	tttatcttga	tgccattca	gagctcctc	80940
taatcttaga	actcaaaaat	ttcttgctgc	agagcttctc	tgcccatatg	tgtctgctga	81000
cctcaccttg	tcagtcaccc	tgctgaaga	taatagtcat	tttttttttt	tttttttttt	81060
tttttttttt	gagacggagt	ctcactctgt	cgccaggct	ggagtgcagt	ggcgcgatct	81120
cggctcactg	caagctccgc	ctcccgggtt	cacgccattc	tcctgcctca	gcctcccag	81180
tagctgggac	tacaggcgcc	tgccactatg	cctggctaata	ttttgtatt	ttttgtagag	81240
acggagtttc	acogtgttag	ccaggatggt	cttgatctcc	tgacctgtg	atcggcccc	81300
ctcggcctcc	caaagtgtg	ggattacaag	catgagccac	cgcgcctggc	ctcttttttt	81360
ttttttgaga	cagagtctca	ccctgtcacc	caggctagag	tgcaagtgtg	caattttggc	81420
tcaccacaac	ctccacctcc	cgggttcaag	cgattctcct	gctacagcct	cttgagtagc	81480
taggacaaca	ggtgcatgcc	accatgccc	gctgattttt	atatttttag	tagagacagt	81540
gtttcgccat	gttggccagg	ctgggtctga	actcctggcc	tcaggtgatc	cgcctacctc	81600
agcctcccaa	agtgcctgga	ttacgggcat	gagccaccac	tcccagccaa	tatagttatt	81660
tcttaagta	acaataaatg	gccaggcacg	gtggctcacg	cctgtaatcc	cagcactttg	81720
ggaggccgag	gcgggcagat	cacctgaggt	caggagtgtg	aaaccagcct	gaccaacagg	81780
gagaaaccca	gcctactcag	gaggctgaga	caggagaatc	gcttgaaccc	gggagataga	81840
gtttgcagt	agctgagatt	gcgccattgc	actccagcct	gggcaacaag	agtgaactc	81900
catctcaaaa	taaataaata	aataaggtaa	caataaagac	ccataggcca	ggcgcggtg	81960
ctcacgcctg	taatcccagc	actttggaag	gctgttgccg	gcggatcacg	aggtcaggag	82020
atcaagacca	tcctggctaa	catggtgaaa	ccctgtctct	agtaaaaaa	caaaaaaatt	82080
agctgggcat	ggtggcgggc	gccgatagtc	ccaggtaact	gggaggctga	ggcaggagag	82140
tggcgtgaac	ctgggaggca	gagcttgca	tgaggcgaga	ctgcaccact	gcattccagc	82200
ctgggtgaca	gagtgaact	ccatctcaaa	atcagacaaa	aagcaaccac	taaagaccat	82260
atgctgtgat	atggaattta	tttttagagtt	attaactgta	tttcgtccca	tttcctgcc	82320
ttttttgtct	cctggtttct	gttttatcta	ttgggaaaaa	tgaacaaaaa	cctctttata	82380
atttcttaaa	ttaaaagggt	aagctcgcct	gggcacgggtg	gctcacacct	gtaatcccag	82440
cactttggga	ggccgaggcg	ggtggatcac	aaggtcagga	gattgagacc	agcctggcca	82500
acattggtgaa	acccatctc	tattaaaaat	acaaaaatta	gctggctatg	gtggcacgtg	82560
cctgtaatcc	cagctactcg	ggaggctgag	cgaggagaat	cacttgaacc	cgggagccgg	82620
aggttgca	gagctgagat	cggtccactg	tactacagcc	tggcgacaga	gcgagactct	82680
gtctcaaaaa	aaaaaaaaaa	aaaaagggtg	agctcttgaa	tgggctcctc	ctccaacctg	82740
cccagcagc	attcatagag	cagtatcttg	aaagtctgct	tccaggtgtg	actcttctcc	82800
ctacagtagt	taagtttttag	agcatgattt	ggtgtgtgtt	ggtcttcgtc	aacagttctg	82860
ccaagcggtt	gcttggtcat	tatctgtgct	gaaaaggcag	ctggaggcaa	atctgagagg	82920
ctaggactga	gaaggggtct	taagacaagg	ccagaagtca	gctcacagtt	gcaggaagat	82980
gttctcagtg	aatctgaaa	ccagtttatc	tctccatgga	tgttctgaca	ttgtctcct	83040
ggtttctgtc	tttagggcat	ttgtttacct	gagcaatctg	ctgtatccc	tggccctgat	83100
ccacagggtg	gccatcgta	gtgagaaagg	tgaagtgcgg	ggattttctg	gtgtggctgt	83160
acaggccatc	gcaggtaggt	gacctcttc	tgaatgaga	gctgtgagtt	ctttgtctag	83220
agacaaagca	ggctgattct	gcgtgggtca	cgcttatatt	acacatacaa	actgtgggga	83280
gtttgtagt	gtaatatata	gacacatact	ttggagaaa	cctttattgc	tcctttctag	83340
caattgttcc	caagtatcac	aaatagatcc	tgcttagtcc	agactactgc	tactttgcaa	83400
taatggggtg	aagagatttc	cttaaacatg	ttaaagaaca	gctcttgcat	taactatata	83460
aaagtcattg	gtgtgtgtgt	tttttttttt	ttgagacagg	gtctcacttt	gtcgtccagg	83520
ctggagtga	atggtgcaat	catatctcac	tgcatccttc	acctccaggc	tcagccatcc	83580
tcccacctca	gccacctgaa	tagctggaac	tacagatgca	caccaccatc	ccagctaatt	83640
tttaagttta	taaattttat	agagttgggg	tctcattatg	ctgcccagca	tagtctcgat	83700
cgaactcctg	agttcaagtg	atcttctctg	cctggcctcc	caaaatgtta	ggattacggg	83760
tgtgggcat	tgacccggc	cagtcatttg	ccttttattt	tgttttgaga	cggagtgtcg	83820
ctctattgcc	caggctggag	cgcagtggca	caatctcaac	tcactgcaat	ctctgcttcc	83880
tgggttcaag	cgattctcat	gcctcagcct	cctagtataa	tgggattaca	ggtgccacc	83940
atcacacca	ggtaattttt	atatttttag	tagaaacgaa	gtttcaccat	tttgccagg	84000
ctggctgtaa	actcctgacc	tcagatgatc	tgccacacct	ggcctcccaa	agtgtggat	84060
tacaggcgtc	agccactgca	cccagcctgg	cctaaataat	atatcttaac	tgcttcatta	84120
ttacaaaagg	attgttaggc	ctttgtcttg	cttagttgtt	gtttttttgt	ttgtttgttt	84180
ttttctcaga	gtacttgtaa	tagacatttg	tttggtgtgt	gttttgtaat	gcacgggtca	84240
agctgatata	agctgattgt	ggtggctcgt	gcctgtagtc	ccagctactt	gggaggctga	84300

ggcaggattg	cttgagccca	ggagtttgag	gccaacctgg	gcaacatagt	gagaccccat	84360
ctcttaaaaa	acaattgaaa	aaaaaaaaag	attttggtat	tttccatttt	tccatgatcc	84420
agcctatggt	ctgaggaaaag	aatgtgggtca	tttgctgtaa	tctttctgtc	ttaatcccca	84480
aatgctgggc	tgtgtccagt	gacctttcag	cctttccctt	atggaaactt	gtgacacaat	84540
caccagtgtt	gagtatatgg	aagtctcctc	cacaagtaca	atataatatt	ttcaaatact	84600
tgttgggtat	tattgactat	tatcttgcca	gcatggagac	agtcaggatt	tctaggtatg	84660
acatttgaac	aatctcaaga	ataaaactga	ttcttttttt	ttgagacgga	gtctcgctct	84720
gttgcccagg	ctggagtgca	gtggtgcaat	ctcggctcac	tgcaacctcc	gcctcccggy	84780
ttcaagcaat	tctcctgcct	cagcctcctg	agtagctggg	attacagggtg	cccgccacca	84840
cgccccggcta	atttttgtat	ttttagtaga	gacgggggtt	caccatgttg	gtcaggctgg	84900
tctcgaactc	ctgacctcgt	gatctgcccc	cctcagcccc	tcaaagtgt	gggatgacag	84960
gcgtgagcca	ccgcacccgg	cctgaaaccg	attctttatt	gactagtctc	atgtgagctg	85020
aggaagtga	taaacttaca	tgaatagagt	gcgaagcage	ccatgccgtg	ttactacttt	85080
ggatatcattc	tctcatgcca	aagacttgag	tttataatcc	agcaagagct	ttttccctcc	85140
ctccccctgt	gtagtctcac	tcaattcttg	ctaatttttt	tttctgcttt	agcggatgaa	85200
gaagctcctg	attatggctc	tggaaattcga	cagtcaggaa	cagctaaaat	atcttttgat	85260
aatgaatact	ttaatcaggt	gagaaaccgt	caggaagaag	gaaaaccctg	tggaaagaga	85320
aagaatgttc	agcattggag	atcagttctc	ctgtaaaact	gaaaaataca	gggtgctggt	85380
tgccacagag	tagttcaata	gaatatggga	catgtctaac	aattatgaag	gtaaagggtg	85440
atagactcaa	tattcatatt	ttattcattg	cctttcctca	gacagccctc	agtcctatata	85500
attttccagt	cactattatt	tgttggacag	ataaacagaa	gtaaggcaac	tggggaaaaa	85560
tacgttttta	tactacttta	ccatgtatct	gccctgcctt	tttttccaga	gtgacttttc	85620
gtctgttgca	atgactcgtt	ctggtctgtc	cttgaggagg	ttgaggattg	tggaaaggaca	85680
gggtcagagt	tctgaggtca	tcactcctcc	agaagaaatc	agtcgaatta	atgacttggg	85740
tatgtagaca	tagtttactg	tgcttgggga	cattttcgac	aaagggaata	tattgaccat	85800
tatcaaggga	catagtggcc	ttcatcaact	aggaatggaa	agcatgcccc	actccctcct	85860
ctttgcatta	tttgaacat	ctgtgtcttc	atttgaccct	cttttagattt	gaagtcaagc	85920
actttgctgg	atggtaagat	ggtaatggaa	gggttttctg	aagagattgg	caaccacctg	85980
aaactgggca	gtgccttcac	tttccgagta	acagtggtgc	aggccagtgg	aatccctccca	86040
gagtatgcag	atatcttctg	tcagttcaag	taagctgccc	ctttgctctg	cctccagct	86100
agctgctaac	cgaggtgaca	tctcttgtgc	actgtagagt	gctttacatg	tgtgagggat	86160
taacactctt	gagacaaaaga	ctgatcagtc	tctgaaggaa	aatactttca	tcactcctat	86220
gggagtgaga	accagaaaacc	tgtcctggaa	tgaacagggy	ttccctggca	ggaggagcat	86280
gctgcacatc	ctgctgcagg	gctctgctgt	gagcgggtccc	cctttaccag	gaatgtaggg	86340
tctgatcaca	taagcctgtt	gatgagctct	tctagtcctc	ttcatgttga	aatgtcttat	86400
taataagcct	gaaagcaatg	acttatctag	ctagcaatta	cgctatttta	atactcgga	86460
aaaggaaaat	ttaaacaatg	taaagaagga	taaattctac	atcaactgaa	aaaagctagt	86520
atgtagggaa	gatgtgatac	ctctcatttc	tcactctacat	aggacttgct	ctttcccgcg	86580
gccctctcaa	accctgtgct	ctggatgctt	tcttctcaag	actgccccag	gaaaacattt	86640
ttttgtgaat	cttcagtttt	aatacctcca	aacttcaatt	tgtcttttca	gtacttggaa	86700
agggccacag	tacaagagac	tgaagtgtga	ccccccata	cctaccgccc	tcagtttgcc	86760
actggtgcc	ccaagaggct	gctttgggtt	ctgccatccc	tctcccctca	gaatgtgcat	86820
tggccgtccc	atgggtccta	gtgcagttta	taaagctttt	gtataggttt	tatcacagaa	86880
aattgagtca	aatgtgattt	aaattaaatg	tgatcactat	atagtaagcc	actaatatca	86940
tttgctacaa	gtcatagtgt	ttggaatatt	accattgag	cattgatttc	acctatttaa	87000
tgtgttttca	ctcataccct	caaacttagg	gcaggatagt	gttcaccaga	tgaatctgtt	87060
ttttggaact	gttgctatta	ttctaatacag	gcaactgata	cgttttatgg	aaaacatcta	87120
tttggttaac	tgatgtctga	agtgacaata	atagttagtt	gacctttgtc	ctacaatctc	87180
ctaaaaagag	cagcatctct	acctggtgtt	agctctgggg	aaaagggcag	gttgcttctg	87240
ttgaaagcag	cctcaaaggc	gtcctgtgtt	tttatagtcg	ctcctctggc	ctgtgccggg	87300
aaggatggc	tgagagcac	tgtgaatgag	ttcacctgtc	gtcagggaac	cccaggaatg	87360
ctgcggtgcc	agcaatacag	gaatcaggga	tttgagcac	ttggcaaaag	catgctagt	87420
ttctcagacg	ggtgaactaa	tttggaagac	aacaagttca	taytaccagt	ctgtctgtcc	87480
ccccagttga	aaatagtttc	tttttctcca	tttttctctg	atggaagggg	tgtcctctctg	87540
atggattttt	ctatctggca	gtaaatcttt	gtgtgggtc	ctctttaact	gtgcagttct	87600
ttttatttca	gactaataat	aggagagag	agagagagaa	cgataacatc	tcactrtctg	87660
tgaataatac	taaataatagc	ccattgtttt	aacaaatctt	tccagagtat	ttctgtgtg	87720
tcgagacctg	tgggaaaacg	cttgctgtct	gggggagaga	ggcatataaa	tacattatta	87780
cagtgaatac	attgctatta	aggaaaaaaa	cattactagt	aatgtgggaa	cacaaagctg	87840
ccagggttaag	tcaggaaaga	ctgcctggaa	gagtgacat	ttgaatccac	aacagactct	87900
tatgtaacta	tgccatccac	aggctctcgc	actaactgtg	gaagcttgag	tctgctaaat	87960
tttgtctgta	caaaatgaca	ttgtttttat	ttttagtaaa	aagctttttac	aaagggaagag	88020
gagttaattg	aatgtacttc	ttcactctct	ctaaagcatt	cttgcatata	cctccatata	88080

gctaagtgat	caggggtcagt	taattttttca	gtcattttttg	aaataaaatc	taaaatgcc	88140
tctgtgcaat	atggaaaagg	ttcttgacaa	ggaaagtaaa	taacttaaag	aatctgctta	88200
agtgggaatgc	ttctttctgt	aggatccaca	ttgggctctt	gaaagagatt	ctgataaaac	88260
cgtttaaatgc	attgttcacg	agtcttttta	ttttttttat	gaagcttttt	gcatcgccat	88320
gatgaagcat	tctccacgga	gcccccaaa	aacaatggca	gaggaagtcc	cctggccttt	88380
tatcatgtgc	agaatgtaag	tgacatggac	ctttttgcca	aacatatgtt	tttctggtac	88440
cttggccttc	tcagcccggg	aagggtgata	agaagaggta	caagtcactg	gctgtgcagg	88500
gggagagtag	acaatagagg	gcgtggctag	atggttgagg	tcattccacag	cggcctgagc	88560
agcaagcctt	cttgtctgtg	ttacagtggg	cctagtgaat	gtgggttcgt	ctgccaataa	88620
aatcttccat	cagtttgaat	gcaatgagtt	gttttagtgt	ttctgtccta	gaaaaataag	88680
catgcttaat	aagagtttta	ttaaagtga	tgcatgtgcc	tttaaaccct	aagatactta	88740
tggtgctcga	ttgcaaataa	cataaaaaat	atcaaatacc	ttctgtagct	aaattaagat	88800
ttagatctgt	tttccttgca	actgatagta	taattaacac	attaatgcc	tttacaactg	88860
tattcctacg	tggagtaaga	aaacagaccc	ttccctctct	tacgtgaaat	ttcattcaaa	88920
ctctattcca	aagtcattat	cctcatcaca	atagctaata	ttcactgaga	gtttattttg	88980
tgtagtcac	tggtctgaga	gttttttgtg	aatttaactc	ttacctttat	accatagatt	89040
aagataatcc	ctgtgttagc	tgagatgact	gaagcataga	gcaattaaag	tcactcatca	89100
ggctgggcgc	agtgggtcat	acctgtccc	gcactctggg	aggccgaggc	aggtggatca	89160
cttgaggtca	ggagtttgtg	accagcctgg	ccaacatggg	gaaatcccat	ctgagacacg	89220
agaattgctt	gaacccagga	tgggggggtt	gcagtgagtt	gagattgtgc	cactgcactc	89280
cagcctgggc	aagaaagcga	gactcccatc	tcaaaaaaaa	taaaaaagta	ataataaaaa	89340
taaagttact	catcagtggt	acaagaggca	gagataggac	ttaagctcag	agaaatttag	89400
ctctgggtatt	caggatattc	actactacat	gaaactgtct	ctcattctct	tgctaaccat	89460
gtctgctctc	taaagtaaga	gaggaaagga	gaacttcaaa	gatggaacaa	gcaaacacaa	89520
aatatgccat	cttctttttc	atgttggtat	aaatataagc	aagcatgtat	ctgataaact	89580
ttgatcagct	gctgtgcaca	aggattgtgc	ctgacaactg	gggaatgcag	atgaatctca	89640
gatgtcccca	gtgagcttcg	aattacgata	ggggaaataa	aatgcataca	taaacaccag	89700
tcatgcaaa	atgaacatga	gaaaatattc	actaaatgca	gtgggagtgg	agaaagggaa	89760
agttatttct	attaatggaa	agctattttt	agttactttt	tggtggggag	gatagttggg	89820
tggagaaaag	ggataggaag	ctccctgaag	gatgtagtat	aagcaagaaa	aggtgtgtca	89880
aacttggcat	tgtgcctggc	acataggcca	tagtcagtag	atgcatggat	gtgctggaag	89940
ggccttccag	gcaggggttc	agcctggggc	agtccttggc	acagctgagg	agagagcagg	90000
tgaaggagc	agcaggtgag	tgagtgaacc	agaatagtaa	gtgtgggaag	gacaacaggg	90060
gccaagatca	gttgaggcaa	agtatggaag	atcttgaaata	cctgaagctg	ggaaaagtta	90120
ctttagaatt	aaaaacaaca	taggggatcc	cctgtttttt	gttgcaagag	tcctgcatag	90180
taaaataccc	agcagatact	ggacactcag	cagcagtaaa	attggggccag	gcatagtggc	90240
tcatgcctgt	aatccagcac	tttgggagct	gaggcagaag	gatcagttca	gcacatgagt	90300
tcaagaccag	tctgggaaac	atagtgaat	actgtctcta	caaaaaattg	ttttaaatta	90360
gccaggcatg	gtggtgcatg	tctgtagtcc	tagctactcc	ggaggctgag	gcaggaggat	90420
tgcttgagcc	cgggaggggtg	aggccgcag	agccttgttt	gcgtgctaca	gagcaagacc	90480
ttgcctcaaa	aaaaaaaaaa	aaaacccaaa	atacgtacta	aagttctgat	tgaagcaaaa	90540
atgtgtattt	tcttgtaatc	ttttccta	cttgcctggc	tttagattgc	agtggagatc	90600
actgaatcat	ttgtggatta	catcaaaacc	aagcctattg	tatttgaagt	ctttgggcat	90660
tatcagcagc	accacttcca	tctgcaagga	caggagctta	acagggtttg	accagataag	90720
caaacatttt	tgatgggtatt	gttttgattt	ttagtttctc	aattgctgga	aaggaagaaa	90780
tcttttgagg	gattaagata	actaaataag	tattctatat	taccctgttt	cctctttaga	90840
aacttcaact	tcattcttata	gccaccta	cactgatttt	agatagagca	aattaaattt	90900
tgaccagggtg	acctcaaata	ccatcttttt	gtaggaagag	actttgaata	tggtagaaat	90960
ttgaggtaat	aggaatataa	catgaaaatt	gttaaaaatt	gttccta	cccacagctt	91020
tatgaaacag	aagctctaag	tacagtcac	acaactgctg	aggaaagcaa	cagtgggcct	91080
ttctgatgat	ggcagagtgt	tggtgttggc	agaagaattt	ttttatttat	tagctaagac	91140
ttagagggag	acatttttta	aatgagattt	agcctgtttc	actatctagg	attatctcta	91200
ttatctggta	ttatgccaga	gttgaaaatg	gaaaagtctg	cttccttaaa	gggaattaaa	91260
acatgctatt	gataccaaca	tccaactctt	tgtagtcttt	ttctttctct	tgaaaaaaca	91320
gggtttattg	atttctacac	tgcaaatagt	cattacaaaa	agtttttttt	cttttaataa	91380
aattcacaca	aagagaatta	gaaagcgatg	gtaatgacca	gcaagaggaa	taataattac	91440
gttcatctta	atgtctgtgt	gccagttctg	tttacgttaa	tggtggaaaa	ctccaacttg	91500
gaatccagaa	cttcaaactc	gcaaatagaa	tgtcttgaga	taggcacgtg	gaagtcaaac	91560
agtttctctc	ttttccctg	ctttcagaag	ctatacataa	aatgttggtg	ccttctgtac	91620
tgtcacagaa	cttttacgta	catttctact	cctacttggt	aaacccttaa	agagaaggaa	91680
attcaatttg	cagtccagca	caaaggggag	aatttctaaa	ataaataact	ctttatagtc	91740
ttgatctttg	tcttcctttc	tttgagctcc	gcctcagccg	tgccgcgat	tcttccctcc	91800
acccatgcc	ctgtccaagc	caggtgagca	ctcgctccgc	tttttgcatg	atgatctctt	91860

tgtgaataca	tgtcttattc	tgaatgactt	ttcaaataga	agtcattgtt	gaattcctat	91920
tcttctatat	atccacatgt	attcctgctc	ctcttgtatg	tatctagtag	gaactagaat	91980
tcctagttcc	tgatttaaaa	taagaagtaa	aggctgggcg	cggtggctca	tgcctgtaat	92040
cccagcactt	tgaggaggccg	aggcaggtgg	atcacctgag	gtcaggagtt	caaaaacagc	92100
cagaccaaca	tggtgaaacc	ctcgtctcta	ctaaaaatag	aaaattagcc	gggcatgggtg	92160
gcacacgcct	gtaatcccag	ctacttggga	ggctgaggta	ggagaatcgc	ttgaaccggg	92220
gagatggagg	ttgcagtgag	ctgagattgc	acctttgcac	tccagcctgg	gcaacaagag	92280
cgaaactccg	tctcaaaaaca	atataaaaata	aaataaaaata	aagtaataaa	ataagacaag	92340
aagtaagaaa	caataggggg	ccttatctac	aaactctctg	gtcttctatt	tgcagattttt	92400
gataatcact	ctgttcctaa	atctttctgc	tttgcatgta	acttcccata	actgtcaggg	92460
gactaggcat	ttctttgcat	agtaaattta	ttctgtatgt	tcagtgatct	tgttttgatt	92520
ttcttttatt	tctgtccttt	ttcctcattg	cctattcgcc	aaattcccac	cccacctgt	92580
gtcactctc	cacctctgt	attcccac	ttcatctgcc	ttcagaccat	gagagaaaga	92640
ttgagctgat	tcggtttctc	gtgtctggct	atgtgaatgt	gcatatgttg	gacacatttt	92700
ctgagcacgc	cagtgtgctt	gcatcctccg	ctattgccac	cttccaggat	gaggctgacc	92760
cattgccgcc	ttttctcttt	ctgccagttc	ccagctgcc	gagcgagagg	gtacccaaac	92820
gagatgggta	gtagctgaat	tagctccacc	ttagatgtag	aaccatcctc	aggcgcttgt	92880
cctgaagcct	ggctctctgc	atatgaaaca	ctgacatttc	cctttgcaac	ttgagctgaa	92940
tgtagagctc	agcacatttt	caagggtgta	tcacattttc	ttcctagaga	tttgggaagc	93000
agcttctcta	catgtgcctc	tttcacatca	ccatcgcagt	gggggaggcc	tttgcathtt	93060
tttttatgga	ttctttgcaa	tagtattttt	aattttgaag	attttactag	tagttggttt	93120
gttagaatac	ttctggtttt	gctttctgac	ccttagatga	ctgccaggat	cacttacct	93180
atcttctttc	tgactagcat	caccaactca	tgtgaaactc	agctgggctg	aatactgctg	93240
ataggggtaca	cgtgtgaata	gtgctaaatg	attcataaag	ctcctttatg	tacattatgt	93300
tagtggatct	tcatgattgc	ttttcaagat	agtaggctaa	atattaggg	ctacagttta	93360
taaatgagga	gacagactcc	cagagggtta	gcagcttaac	taaagtcaca	tagcatcagt	93420
acagttaaga	atatggagtt	taaatctggc	cggttatggt	ggctcgcacc	tgtaatccca	93480
gcatttgagg	ggctgagtcg	gacagattgc	ttgagctcag	gagttagaga	ccagcctggg	93540
caacatggcg	aaaccctgtc	tctacaaaaa	aatgcaaaaa	ttagccgggc	gcgggtgtgc	93600
acgccttttg	tctcagctac	tcaggaagct	gaggtgggag	gatggtttga	gccagagggg	93660
cagaggttgc	agtgaagccg	gatcacgcc	taaaaaagaa	aaaaaagcat	atggaattta	93720
aatctgaaga	aaatgtgtga	cataagatca	tttacagaat	cctgtattta	taaatgatac	93780
cagtagttgg	tttccaaact	tgctctcaca	ttggtatcac	ctgggaaact	ttaccctca	93840
tgagggaaga	actttgtctt	tccttagag	attgtgattt	aattgatctg	ggatgtgtcc	93900
tggaacttaa	ggctcccagg	tgctcctatg	tgacgggagg	tttgggaacc	actggggtgg	93960
aggatgggtc	ttttcttcag	gacatcagct	ttgattgagc	ccctgtgacg	tgaggctttc	94020
caagcattgt	ctctcccacg	ttagtcctat	gctttacgtg	tcacacaatc	tgagattctg	94080
gaattggaag	gtttcatgag	cacatttatt	ctacacaaaa	gctgttttct	catcattaga	94140
gacagtttgt	tgagaaaaaa	gatactgatt	tctttatttt	cctgtttctc	tgaatgtaga	94200
gataactaat	aacttacaaa	gtttatatgt	gtgtttctct	gggtcttttg	cattgaaatc	94260
tttctctga	ctcttgctgc	ctttcccaag	tatgtcttta	aagactgaca	ttagagaagc	94320
ttgtattttt	gctgagtagt	cttgaccag	atthtgacat	actctaaaaa	cttttaaaag	94380
ttccagccac	caagttaaac	acgatgagca	aaaccagcct	tgccagagc	atgagcaagt	94440
atgacctcct	ggtttggttt	gagatcagtg	aactggagcc	tacaggagag	taagtccaac	94500
ttaataaatt	tttaataaag	gcaaaatgtt	tcaaatagg	aaatgcaatt	ttgaatcttc	94560
ttgtatttgg	gttcatgaat	cattgtcact	gtcaaatcc	agggatggtt	ttgtttttgt	94620
ttttgtttta	atagagatgc	agtcttgta	tgttgaccag	gttggctctg	accactcctg	94680
gcctcaagca	gtcctcctgc	ttcgacctcc	caaagtgtctg	agattacagg	catgagccac	94740
cacacctggc	cgaattccag	ttttgaattg	atthttgggtg	gtcttggtat	ttgaatcaga	94800
tgtttatcag	agaactgact	gttgagcata	ataatttgcc	ttctcgttag	caacagcaaa	94860
atcatggaga	caggcatctt	gttggttttt	gggttttttt	ttgttttttg	ttttttttga	94920
gatggaggct	tgctctgttg	cccaggctgg	agtgcattgg	cgcatctca	gctcactgca	94980
acctctgcct	cccaggttca	agcattctc	ctgcctcagc	ctcctgtgta	gctgggatta	95040
caggtgctgt	ccaccagcc	aggtaattt	ttgtattttt	agtagaggca	gggtttcacc	95100
atgttgccca	ggctggtctc	gaactcccga	cctcaggtga	tccaccgccc	tcagcctccc	95160
aaagtgtctg	gattataggg	gtgagccact	gtgccagca	gcacctgtgt	gtaagggttt	95220
tatcataatg	ggttttatta	gtagtaaagc	aaaaggagca	tattgcatag	aatatggtaa	95280
agacttacat	tactgatgtc	aaggaggaca	tagaagggca	gaatattttc	ctagaccagg	95340
gtttctcaac	ctcagcacta	tttggggctg	gataattctc	tgtgtgtgag	ggctgtcctg	95400
tgcatttaga	atatttttga	gcatccttgg	cccctaccca	caagatgcc	gtagcagctc	95460
cactctattt	gtgacaacca	aaaatgtctt	cagacattgc	caaatatccc	ccaggagggg	95520
catcctcagt	tgagaaccac	tgtcctagat	caaagtgtga	ccatcagctt	actctgtctc	95580
gatgagctga	gttagagtaa	gtgattaatg	taattgttgt	gtataatcat	ctctaaaatg	95640

cttttgtaaa	aacacaaatg	aaagagatct	tactcttagg	acagaggcaa	ggggctgtat	95700
cagttaatat	aaagggtagg	agtaaatgtt	atcctaagt	tctcctatat	gtaattctga	95760
gaaccaccat	catgggtcatc	attgggaaac	acttattaag	ttcctactat	gtgtaggata	95820
ctgtgttaag	tgtgggtatgt	agttgttagt	agtgacagtg	ctttgaggaa	tttgactttt	95880
catagacaga	cttatgatgt	gattacaagt	tgcattatga	gggagaggta	gttacagtaa	95940
ttgtttctac	tgaattaatt	gctttctgaa	tcttaaaaaa	aactatagtc	tgaattatc	96000
tgtgtttctt	atctcctgtc	aattgccatt	gcacttctac	attattttatt	gaatacttaa	96060
tatcaagggg	ctggatgcaa	ccaaaagatg	ttactgtcct	ttgactactc	aagtctagag	96120
aggattgaga	catagatgtg	tggaaaggat	gagtaaatgc	aaatgactgt	atgattaaat	96180
gccagcatgg	gtgagaaaga	catgggatca	gaatttatca	aaacaaagac	caccaggggc	96240
taagccttgc	agatgaagta	gcacttagtt	ttttcttttg	cgtttctatt	ttttagggtat	96300
atcccagctg	tggttgacca	cacagcaggc	ttgccttgcc	aggggacatt	tttgcttcat	96360
caggttactaa	tgagggacca	aaacaggcat	cggagggaac	actgaaaaga	atttagaata	96420
agagacggaa	ttctttctta	ttctctgttt	tcactcttat	ttcagagggt	gggctgggca	96480
ttgtcctgtt	gtcatttttt	ttttttctg	tgtctgtctg	tagatatttc	ttcctcagac	96540
cttggttat	atttatcctt	gcttttatat	gtatttagaa	tagctatttt	gaaaaatatt	96600
ttgtttcatt	caattactct	aatttgtgta	gtccttatgt	cacttttaat	acttttcatt	96660
tcgtgatgtt	tggaagctct	gtgctttcac	accagaagtt	cttaatagga	atagaggatg	96720
aaggtttcaa	tggactaaaa	atctggttat	gaaacatagg	tccgcttttg	gtaaatcaaa	96780
agtaggttca	gtcagtgttc	aaaagtactt	aagaccaag	tgggcatagg	ggctagggat	96840
accatacatt	tgggacatac	aatcagggca	tggttgaata	cttattatac	aaggggctgg	96900
gatgcagcca	cataatctcc	tgtgacagat	tatggttatt	cttaaaaacca	tagagaaaaa	96960
gaatttttta	aaaaagaaaa	gaaaaccata	tcagaacctg	actgttcctt	agtcctgac	97020
acatctccct	ttgttttgaa	taacaatgtg	gatagcattt	tccgtctttc	cacacctctc	97080
cttctgtctc	atcccccatt	ccatgccata	caggccagaa	aacagtagac	gagaggagat	97140
aatagattgc	ttcagctaaa	ttgcaaccct	gcttcattac	ctagggcatc	cagcgaagga	97200
tcacagtgc	cattatccat	gagaagggga	gcgagctcca	ttggaaagat	gttcgtgaac	97260
tgggtgtagg	tgagtacgtt	tcatacagcca	aggatagaac	caggacttac	agagattttt	97320
tttttttttg	agatagggtc	ttgctctgtc	acccaggctg	gagtacagtg	atgagatcat	97380
agctccctgt	agcttcaa	ctcctgggct	caagcaatcc	tcctgcctca	ggttcccag	97440
tagctgggac	cacaagtgtg	aggcactgtg	ccctgttaga	gatgttattc	ttccctgttt	97500
gcggtacatg	gaatgtgtag	aagaccatat	tcggaaacac	agactatgga	gatgcagagg	97560
ccttgagttt	gaaactctgt	attcctgttt	cttcgctctg	tcattagtga	tatgttatcc	97620
aactaactaa	ctcagttacc	agggaatggt	tctgtggtag	agtctgtgag	agggcataaa	97680
agaaatctct	gctggctggg	tgtgggtggt	catgctgtga	atcccagcac	ttttggaggc	97740
tgagatgggt	ggattacttg	agcccagggg	tttgtgcctg	gccaaagtag	caaaatccca	97800
tctctacaaa	aaaatatgaa	aattagctgg	gcatggtggc	acgcacctgt	agtcttggt	97860
actcgggagg	ctgaggtggg	aggatcgctt	gaacttggga	ggtggagggt	gcagtgcgct	97920
gagatggcac	cactgcattc	aagcctggac	aacagagtgg	gactccgtct	caaaaaaaaa	97980
aagaaaagaa	aaatctctgc	taattgtgag	cttgcathtt	agttagggag	acagcatata	98040
tatgtgcaga	atgctttgat	gaataaatac	caaaatgagg	acaaaggaaa	atctgtgcta	98100
agagagttaa	cagtgaatac	aagccaggag	gactgaatag	attcaagaaa	aattagaggg	98160
tttttttttt	ttttccagac	acaagcattt	tgagtaagag	agtcaggaa	atgtctaaca	98220
tgcaaatgac	aagaaacccc	atggccagag	tggagaaagt	aaattaagga	gtattggcta	98280
gatagtgtaa	actttagaat	gacagtccga	ggcaagagtt	tggaaacttt	gttgccatgg	98340
acctcttttg	cagtaacgaa	gccttagaat	tatgttttca	aatgtgagaa	ataaagtatt	98400
taggattaca	aaggaaaccg	ataatattga	aatacagttg	ttaaaacatt	aagctcttct	98460
ttactgatgc	atgaaataac	aagatttagc	agttatttcta	attaccataa	tttccaagta	98520
gtgatcaatg	aaaaagatat	cttgtgtcca	ggcacgggtg	ctcacgcctg	taatcccagc	98580
actttaggag	gctgagatgg	gcggatcaca	aggccaagag	attgagacca	tcttggccaa	98640
catggtgaaa	ccccgtctct	actaaaaata	caaaaattag	ctggttgtgg	tggcgggcgc	98700
ctgtagtccc	aggtactcgg	aaggctgaga	caggagaatc	atttgaacct	gggagacaga	98760
ggttgacagt	agccaagaca	gcagagtggg	actctgttta	aaaataaata	aataggccgg	98820
gcgtggtggc	tcagtccctgt	aatcccagca	ctttgggagg	ccaagatggg	cggatcacga	98880
ggtcaggaga	tcgagaccat	cctggctaac	acggtgaaac	cccgtctcta	ctaaaaatc	98940
aaaaaaatta	gccaggcgtg	gtggtgggcg	ccttagtcc	caactgctcg	ggaggctgag	99000
gcaggagaat	ggcatgaacc	tggaggcgga	gcttgacagt	agccacgatt	gtgccactgc	99060
actccagcct	gggcaacaga	gcaatactcc	atctcaaaaa	aatatataaa	taataaata	99120
agatattttg	cacatggaaa	catctgtggt	ttctactggg	gacagagtca	cgggtactgg	99180
ccaattcttc	tgtggtttat	tgccttcctt	tgtgattgaa	ggagatgcca	atttttagtt	99240
aaaggtttgt	gaaaaataag	atgtaatttt	tttcccctac	caagctcaag	gaccctctga	99300
attctgtcca	cagagtgcct	gtgtctctct	ggaccagatt	aagaaccttg	gggccaggca	99360
cagtggtcca	cgctgtaaat	cccagcactt	tgggaggcca	aggcaggcgg	atcacttaag	99420

ttcaggaggtt	tgagactatc	ctagacaaca	tggcaaaacc	ctgtctctct	acaaataagc	99480
tgggtgtggt	ggctcatgcc	tatagtctca	actactcagg	aggctgaggc	tggaagcttg	99540
cttgaacctg	agaagtcgag	gctgcagtg	accatgatca	tgccactgca	tgacagaaca	99600
agacctgtc	aaaaaaaaaa	aaaaaaaaaa	tcgttggaag	ggagattatg	ttctctcttg	99660
tctggggcag	cagtggtagt	agcttttcaa	tatttttcat	aataaatttt	tcttttttta	99720
atttagtttt	ttttttaact	ttattttaca	agtactgaat	gtcctttgac	aaaataatgt	99780
ttatatggaa	ccctaatacc	gaaacagaag	agatgctttc	tgaccagaga	aggagaggga	99840
ggatgcagag	gcagcagtta	ctatggcttc	tctgaaggat	cctatgatat	atttttagtt	99900
tgttttatta	acaaactctg	atgtagtatt	taccacatgc	cagggtgcttt	tctgaatgct	99960
ccaaaaatac	tagtttgttt	aatcctcata	gcaacctat	atggaaggaa	ttgttattct	100020
cgcttgacat	atgagggaac	tgaggctcag	agaggttgag	ccttaggctc	aaagcctcac	100080
aggtggtagg	agttggaaac	caggcagtc	agcactggag	gccaagtctc	taacgtctat	100140
gcaatgctgc	atccgctagt	gttaacttga	aaacagagtg	taggattttg	agcaagaaag	100200
gggtgtaata	aagtcagcat	ggccagggtat	ggtggctcat	gcctgtaatc	ccagcacttt	100260
gggaggccaa	ggcaggagga	tcgtttgagc	tcaggagttc	gagaccagcc	tgagcgacat	100320
agtggggccc	cgtctctaca	aaaaaataaa	ataaaataaa	atagctggat	gtggtgctgc	100380
aggcctaaag	tcccagttac	tcagggtggct	taggtgggag	gatcacttca	gccaggaat	100440
tcaaggctgc	agtgaagct	gatcaaacca	ctgctctcca	gcctgggaga	cagaccaaga	100500
cattgtttca	aaaaaaaaaa	cacaaaaaaa	caagaactg	tcagtcagtg	gcttcagaag	100560
ttgctgttag	gtgagagtgg	aatggattgg	aacggggagg	tgtgaatagc	aggtcagaag	100620
ttgctgttag	gtgagagtgg	aatggattgg	aacaggggag	gtgtgaatag	caggtcagaa	100680
gttgctgtta	ggtgagagt	gaatggattg	gaacggggga	ggtgtgaata	gcaggtcaga	100740
agttgctggt	aggtgagagt	ggaatggatt	ggaacgggga	ggtgtgaata	gcaggtcaga	100800
agttgctggt	aggtgagagt	ggaatggatt	ggaacggggg	aggtgtgaat	agcagggaga	100860
ccagttaaat	ggctgttgca	ggttccagac	cagagaggag	gtggcctttt	gtgatcaggg	100920
caatgatatt	gtgaataaaa	gttgggatga	atgtgggaaa	ctctttgaat	aaagaatcaa	100980
tatggtttca	tagttaactc	aatgtgggaa	gtaatggagt	cagaccgtgg	aggaccctac	101040
aggtaaccaa	atgacccttc	cagctcctga	acttacacag	agattttaaa	gtctcttttg	101100
ggcttaatga	attcatgttt	tgttttcaaa	tgtgtccgtg	ctctgttttt	tttatccctt	101160
cttttaggtc	gtattcggaa	taagcctgag	gtggatgaag	ctgcagttga	tgccatcctc	101220
tccttaataa	ttatttctgc	caagtacctg	aagtcttccc	acaactctag	caggtgggac	101280
accagagca	gtgtgaagaa	gtccacactt	gcaggcggtta	attggtacac	cgtaggtgct	101340
cttattcatt	aatgactccc	agttcggaca	aagaaattaa	ctcccttctc	ccttctagct	101400
tcaaaaatct	ctttattttc	tcacctgcct	gctgtactct	ccaaaaagaa	agaaagaaag	101460
tattgcagat	atttgtatgt	gatcagttac	tcttagagaa	tggaaagtga	cctgtcccat	101520
gtgaagtttg	aatagatgta	acaagtgata	aatgaaaatt	ggagaaagaa	aacagtatat	101580
tccccaagga	tttaaaagta	craattaatt	attgccaaag	ttaatttttt	tctgtgaaa	101640
tgggtgctgt	ggagagaaag	gtttctctca	ctccccaagt	atcatggaaa	tgtgccctct	101700
gagataaaa	gaagcctctt	gtgttaagtc	tttgctgctg	ctgactatag	gtccatttat	101760
ccttagtatt	tgtttttcat	ttatgcacaa	ctttgttttt	taatatgaac	tcttatgacc	101820
aagattggct	gccataaata	atagatttta	ctttgttttt	tattttaaat	gtttcactta	101880
aaattatctt	ttataatcag	gtgataacag	acagttcgtg	aagtacatag	ccattctgct	101940
ttcctttgta	agactagAAC	taaaccagct	gggatcctct	tacctcacc	aggaagggag	102000
ttctcattgt	ctcagaggct	ccgtctctat	atcctagggg	ctgggaaaca	ggagcggctg	102060
cctcttctaa	tgtgggctga	gcacactgcc	ttcccaccac	aggactttgc	tgtgcttcag	102120
tctgtcctcg	aaaaacttgt	ttacatccat	gttgagagga	gggagggaag	gaataagacg	102180
ataggcagaa	atggaatgaa	aggtttgagc	caggattaat	ttctttctgt	tgttttcttg	102240
tattatagtt	taccatagga	aagagaaaaa	agaaacaaan	ttctgtttgc	tggcaggggc	102300
tgtggagaaa	gagggagga	gtgagagggt	acatatgtat	gtctggggag	ggggagtggc	102360
agaaataaag	aggggtgctt	tgactatcca	tgggcaaaat	cacgagcaat	actatttccc	102420
tgaagtcagc	agtaattctg	catagctttc	tgttttctgc	caatgttcag	ccgggtttta	102480
ccaccacatt	ttctttgtag	ctatgaaaaa	tgtattgtaa	atttccctag	ggtacttagt	102540
aaaatataca	gggaaagatg	ggcttttggt	gaaatactgg	ttatttctaat	tttgcttcca	102600
ttagagttaac	tttaagaagt	cctttaattc	acgccaaca	acaggtcagg	tgctacaaga	102660
aatgaatttg	gctttgttag	aagcagctta	tcttgtagt	tagaatgagt	tcatcaaaaa	102720
aggctgaagt	caagaataac	aacaacaaca	aaaaacacct	taaaattttc	tgttctcatt	102780
taaactatat	aaatctacat	taatttatca	aacttttgct	ggaaagccat	ggccttttct	102840
ttatattata	gatcatttac	tggagtctcg	attgtctttc	ttgcaaaacc	acaccataaa	102900
ttcatcatct	atgactttca	gtttggtgga	gcaagaatta	gaaacctgtg	aagaaaatct	102960
aagttaaaaa	tcctttaaat	tttttatcat	ttattaacat	taatcattta	aacattttat	103020
cattttcccc	tttcttttcc	atttgtctca	ccaaaactta	acagttattt	ttaaacaact	103080
caccattttc	taaccttttg	aaaacattca	acttagtttt	cacaggaaca	aaaattattt	103140
atgttttcat	atttcattga	tttgccta	gtttaatgaa	gttagataat	aataataagt	103200

acaactggca	ttaacagatt	tgggaccaga	tactactgac	atttgataag	cactcaactc	103260
ttcggagcga	agtgtctggg	caaatgagac	atctttgtag	tttagaatgc	tcaaggagaa	103320
ctagccatca	atctgtatat	tttagggata	tggaaaacat	aggtaaatat	ggtaagtggc	103380
aaaatggagt	ttgggttaaga	gaactttcgt	tctgtttgtc	taagggttaa	tgtttaaaagt	103440
tgaacttacc	tgcaatttgg	gtatattatg	tgtgtatctt	agagggcttt	gctaatacat	103500
tgatagtggg	gtataattga	caagggacca	ttcattatcc	cttttcttcc	cagtccctaca	103560
tttatgttaa	tgcatgtgact	tggttttccc	ttaactccac	agcaacttcc	tattattctc	103620
aataactaagt	caccactgaa	aggggtactg	atgggtgggtg	tctccacttt	ccctcttatg	103680
tctggcattt	tccttagttg	gaagtctcag	gagagggacc	catgttattg	taagtccctc	103740
atggaattgt	gttccatggg	attctgtgtt	aattgtctgt	ccacttgaat	tcactttgct	103800
tatcagcaaa	ttttgtcttc	tttatatata	tattttttct	gattctctcc	cttatttctt	103860
ttgttttttt	gctgcactaa	ctaaagcctt	ttctatgttt	ttgcaggctc	tttcttgata	103920
aggatattcc	taggtattgt	ctgtttgctt	ttttgcacgt	aggattctag	tttatcagaa	103980
tgctttctgt	cttatgttgc	tttgccattt	taattttgtat	gtccccctcc	catgtaaac	104040
tactgtacct	aggatagagt	agaccctgtt	ctgtacccta	agaagcgtat	agcttcttaa	104100
atgattttta	gagtttcaaa	ttattttttc	attccatttg	ctaaacttgt	cagttttatt	104160
cttagactct	ttcaagtttt	cctcttctac	attacaaaagt	agaatgtttc	tcccttcatt	104220
tcatacagaa	tctcaggttt	tgttttctgt	tttctctttt	atctgtcttc	tctccgtcaa	104280
aagcatgaag	ggttttattgc	tttataaaac	ctgagtcctat	tgcatgtctg	caaaactgtt	104340
cactttcttc	tctgttttgt	cctgtgtggt	aatagagaca	aattttggcta	yttggatagc	104400
tgagtctatg	agaaatattt	tatatatatt	ttttaacatt	tatctttctc	ctgtcagggc	104460
tgcttgtttt	atctgtttat	tttcttcttt	tggaatagga	aaaatcctgg	tttggtatta	104520
aagggaagttt	tttaaaaaat	ctgtctcatt	tgggaagaa	aattgaagga	agtatacat	104580
gacaatataa	cctaaaaatgt	taaaaaaaat	acaacataaa	cgggctgggc	tcagtggctc	104640
acacctgtaa	tcccagcact	ttgggaggcc	aaggcgggtg	gatcacgagg	tcaggagttc	104700
aagaccagcc	tggccaagat	gggtgaaacc	catctttact	aaaaatacaa	aaattagcct	104760
ggcatgatgg	tgggcgcctg	taatcccagc	tactcaggag	gctgaggcag	aggatcgctt	104820
gaacccgga	ggcggaaggt	gcagtgaacc	gagatcgccg	cactgcattc	cagcctgggc	104880
gacagagcga	gactctgtct	cgaaaaaaa	atacataaac	atatttatat	atttgactga	104940
tgttactgat	ataatgattt	tatccctcaa	tttgacccc	attcagttta	catttcagag	105000
attctcctaa	tgaactcaaa	agggttcac	agatgtgacc	atgatataag	aagtggtgctt	105060
ctgttttggt	ttcttgagac	gggtgtgtgc	ttgttgccc	aggctagagt	gtagtagcta	105120
tacctgcaga	tgctttaaaa	aaagaaaaaa	aatgctggcc	gggcacgggtg	gctcacacc	105180
gtaatcccag	cactttggga	gaccgaggcg	ggcgatcat	gaggtcaaga	gatcgagacc	105240
atcctggcca	atatgggtgaa	gccccatgtc	tactaaaaat	agaaaaatta	gctgggcgtg	105300
gtagcatgtg	cctgtagtcc	cagctactca	ggaaggctgag	gcaggagaat	cgcttgaaac	105360
tgggaggcgg	aggttgacgt	gagccaagat	ctgtccacta	cactccagcc	tggcgacaga	105420
gctagactcc	gtctctaaat	aaataaataa	ataataaata	aaataaagtg	cggtagcatg	105480
atcatagctc	actgcaacct	tgaattcctg	ggctcaagca	gtcctccctc	ctcagcctcc	105540
cgaaatagggt	ggactccaga	cacgtgccac	catgcctagc	taattttaaa	tttttttgta	105600
gagacagggt	ctcgctatat	tgcccacgct	ggcctcaa	tcctggcctc	aagtgatctt	105660
cccactttta	tctcccatag	tgctgggtga	acaggtgtga	gctaccacaa	ctgaccagaa	105720
gttggttttt	aattgttcat	aaataataaa	ttcaaaggaa	tatatccttc	aaagcagtca	105780
ccttgaaacc	tgtttttcc	ctgatgtgtc	tgctgttcag	aacatcttga	gaaatccatt	105840
tggaattact	tctcaccacc	ttaggaggca	tacaggaaaa	tatgtcttgg	gattcagccc	105900
tgtttgtaac	cctaaattct	attgccagct	tgattaaatt	gacatttctc	cccagtcttg	105960
gttgattttc	ttacaaccac	ctctcaagaa	ttgccagtgt	aatggctcac	acctgtaata	106020
ctagcacttt	cctactaagg	taggaagatc	acttgagccc	aagagttcaa	gacctgcctg	106080
ggcaatgtag	aacttgtctc	taaagaaaaa	aaaaaattaa	ttagccgggt	gtggtggctc	106140
gtgcctatat	tcccagctac	tcaggaggct	gaggtgggaa	gatcacttga	gcccaggagt	106200
tagattgcac	cagtgcactc	cagcctgggc	aacagagcaa	gacccttgct	taaaaataaa	106260
ataggccagg	cgagtggtct	cacacctgta	atcccagcac	tttgggaggc	cgaggcgggt	106320
ggatcacctg	aggtcaggag	ttcgagacca	gcctggccaa	catggtgaaa	ccccatctct	106380
actaaaaata	aaaaaattag	ccagacttgg	tggtgggcgc	ctgtaatccc	agctacttgg	106440
gaggtcagg	caggagaatc	gcttaaaccc	gggaggcgga	ggttgacgtg	agccaagatc	106500
gcaccattgc	actccagcct	gggtgacaag	agtgaagctc	tgtctcagaa	aaataaataa	106560
ataaaatata	tattttttta	aattgaatta	ccacctttgg	tgggtattct	aacagaactt	106620
gctgcagcaa	gtccaaagca	aaaaaaaaaa	aaaaagaaag	aaactttttt	tgacaaaaga	106680
aaaaaatgaa	agaaaaat	ttccctcaat	acaggaagac	tttagtgag	tcaaaatttc	106740
ctttgtgtta	accaggtttc	tgcaagtga	attttctctt	tatgttaagt	gaaagcttag	106800
cctgaaaaca	gaaatgtgaa	gaggcttttt	tagacaagtt	caaaacattt	gcctcagttg	106860
aaagagtcag	tgcatgaaa	cctcagcagt	ggtcctggga	ggtgccacgt	tatgggttgg	106920
aacttaatga	aggcagcagc	attgatagaa	atttctgtca	aatatcacca	aagtcataaa	106980

tgccgtcact	ctaagaagag	gggtttat	taactttagca	tggatgtatc	cagcctctga	107040
aatgggacaa	ataagccccg	ccccttctag	gtacttttgac	aaaaacagga	acacaataaa	107100
agagcagacc	tctggaggct	gaggcaggag	aatcacttga	acccggcatg	aggaggttgc	107160
agttagccaa	gatcatgccca	ccctactcca	gcctgggtga	caaaagcaag	actctgtctc	107220
aaaaaaaaaa	aaaaaaaaaca	gatctcacaa	tgatacagtc	agttctcaag	tggcatctgt	107280
tgaagcagtg	gatgtcgtca	cccagtatta	gaatatgtct	taaaccaaat	taaaacttta	107340
ctttacacag	agttttatgg	acacatgaag	ggtttctctc	tgtgccagag	gagcccaaga	107400
actatagaaa	ggccttcatt	gtccatcatt	tctgttctctg	tttgggtcac	ttggtggctt	107460
tgtctttgac	ttgttagttg	tttctctatc	ttctgtgtct	ttgtagagaa	ttttcttgac	107520
ccacactcac	tgcctcaaag	actagacatt	tgagagagta	ttaagtactt	gaagaagact	107580
tagatctaga	agcagctatg	ttagatgatc	agacctcata	cactgccact	tggagatga	107640
ggtgtggtca	tggggttcag	tttcagaaat	gtgttctctc	tgttgatcgt	ttctggagga	107700
agaaacatgg	ggatgtgttt	agagggagga	tatgccagct	acttatcata	ggaagtctag	107760
aaaaggcagc	catagttagg	tcactatttg	aaaaggagta	agaaggaatg	tattgggcag	107820
gaaagagtgt	gcatagactc	ttctaggtga	atcaggtgag	ctgtgttcta	tctgcggaga	107880
actagtgtg	gtggggcaga	ccaggcatct	tcccctccac	ttgtcatagg	ggcttgagaa	107940
gggaaacatg	gggcagaatt	gggaatcaat	ctgtaacact	gcttttaaac	taagtgtctg	108000
atttcacatt	agacttgacg	aactggaaga	tttaatgtca	attagagtta	atgttgtttt	108060
accagtagaa	aaaaaacgat	attagaacaa	taacaacaaa	atagtttgtg	ggtatctatt	108120
aaattaagtt	tccatttggt	acataaatgt	aagattaatt	tttcaaatgg	cttttttttt	108180
taagctaaca	aatggtat	gtctggctct	agtgttaaca	atttattctc	ttttactgaa	108240
acatcagtta	agaataacct	tgtattcttt	tatttaaaat	aaggctctcg	gccggcacag	108300
tggctcacac	ctataatccc	agcactttgg	gagggcaagg	tgggcagagc	gcttgaggtc	108360
aggagtttga	gaccagcctg	gccaacatgg	tgaaacccct	cctctactaa	aaatagaaaa	108420
attagctggc	gtagtgtg	gtgcctataa	tcccagctac	tcaggaggct	gagtcaagag	108480
aatcgcttga	acccaagagg	cagaggttgc	agtgaagctga	gattgcgcca	ctgcactcca	108540
gcctgggtga	cagagtga	ctccgtctca	aaaataaaat	aaaataaaaa	aataataaaa	108600
taaaataaaa	taatgctctc	tatacttaaa	cccatctacc	cagtttctac	cattaatccc	108660
ttcagatata	tctgtggtgt	ttctgaggcc	cagcagcact	ggattgaatg	atgttttagcc	108720
ctcttgat	gtgaagattg	aagctgacaa	gttaccgaagc	attcacttcc	tgctgtcatc	108780
attaactgac	aggagagctt	acatagtctt	tctttcttct	ttcagtttct	gtccatactc	108840
acttatccat	ttcatcttaa	tccctttgat	tgaccttact	atttttatgt	ttataggcag	108900
aggcaggctt	gtcaaaacaa	aattaaanaa	agcacaactg	gccaggcaca	gtggttcacg	108960
cctgtaatcc	cagcactttg	ggaggccgag	gtgggcagat	catctgaggt	caggagttcg	109020
agaccagcct	gaccaacatg	gtgaaaccct	gtctctacta	aaaatacaaa	aattagccag	109080
gcgtggtggc	gggcgcctgt	aatcccagct	actcggaagg	ctgaggcagg	agaatcactt	109140
gaacccggga	ggcagaggtt	gcagtgatcc	gagattgtgc	cactgcactc	cagtatgggc	109200
aacagagcaa	aattctgtct	caaaaaaa	aaaaaaaag	cacaactaag	tgttccttaa	109260
tgatgggtg	ggttgggagc	aagaggatgg	ggacctatct	tttctccta	ggagataagg	109320
gttctctct	gttatttagag	tttgataaac	tgaaagtgtc	tggtgggtg	cgatggttc	109380
tttaggatga	ttgacagtgg	cagtacctgg	actaacgtga	cttttagctc	ttttggatga	109440
ttccctcttg	tctttctgac	cttaggacct	tctaccgctt	tgaggctgtg	tgggtagact	109500
ctctgcataa	ctcccttctt	ctgaaccgag	tgacacccta	tggagaaaa	atctacatga	109560
ccttgctggc	ctacctagag	gtgaggagac	ttggaacttc	agttgatgcc	aacagtcagc	109620
cctgaacagt	gtgcaggtta	cagcaggaag	tggtcagcga	agattccact	tgttttgtcc	109680
tctttcttcc	tccacatcct	taagtttttc	attttattat	acattgctaa	ctcttctgat	109740
tatattagac	tctattctctg	ggttctgttt	actcaaaaa	tcacatgact	cattgcagct	109800
tcagctttag	tagcctaatac	accattgttt	gaagaatgtg	tgatttttaa	gaaaatgtca	109860
cctctgtgtt	ttgtagtgtg	ggacagagt	taatagtgtg	taatagccca	gaccaagac	109920
ccagcctgtc	ggttcaaate	cctgtgtgtc	cacttacttc	ctctatgtcc	ttggagaaac	109980
tctctttgcc	tcagtttcct	cacctatcag	gagaaacaac	atttacctca	gataaaaaata	110040
aatgcactc	taaggagtac	tacctactct	tgacacatatt	atccattggg	tggctgttga	110100
aataattggt	ggagctaaaa	tccttcagtt	ctcacaactg	ggactcgctt	tccaaataacg	110160
ttcgtgtata	gactctagtc	acagtactct	gcctgcactt	catcagcacc	taccctgtct	110220
gctttcagct	ggatcattgc	atccagccgg	ctgtcatcac	caaggatgtg	tgcatggtct	110280
tctactcccg	agatgccaag	atctaccac	cagctctct	gcgtagcctc	tttggcagcg	110340
gctactcaaa	gtcaccagat	tcgtaagttt	ttcacacaag	ttagcttcca	gtgtgtttgt	110400
tcagtacaac	agaatcatta	gtccttaagt	attctcggt	tactgtctca	cggttagttt	110460
acagtttatg	aaccattttg	gtaactgaca	cctggaatgt	actgacttgc	atgctgacat	110520
ttaataaaaat	gttgatggat	ggacgagtta	tgagggaagg	aaggaatatt	acattgtgaa	110580
cacgtctaaa	acaattttgc	aaagactaaa	tttgcaaga	cttttaaatg	tcatcatcat	110640
tatttctaac	agagtccat	tgaatttttc	ttctttttta	gatttggtat	gcatatttaa	110700
ggcatacaaa	atatgtcagg	aataatacag	tgaaggccat	acacagtggc	tcatgtctgt	110760

aatcccagca	cggtgggagg	ctgaggacag	agaattgctt	aagcctagga	gtttgagacc	110820
agcctgggca	acacagaccc	catctccaca	gaaaatttta	aaattaccca	ggtgtggtga	110880
cacacacctg	tagtcccagc	tactcaggag	gctgagggtg	gaggatcact	tgagcttggtg	110940
aggttgaggt	tgcatgagc	catgatagag	ccactgcact	ccagcctggg	tgaaacagtg	111000
agaccctgtc	tcaaaattta	aaaaaaagaa	aagaaaaagg	aaaaaaaaaa	aaaagaaaga	111060
aacaatacaa	tgaataacctg	tgtatcacccg	cccagcttaa	aaataagtg	tactaataca	111120
atagaaggta	ttataacctc	tactactcac	aatccccaga	agttgttgat	tatcatttca	111180
ttctagtcac	tgtttccctc	atatcgtctc	tcttatagta	agaaaagaca	aatttttagt	111240
tttcacattt	atatttttaa	ttttctagtt	tatcattcaa	ttttattcac	cctattgaaa	111300
taaagtgttt	aaaagataga	gatgagggtg	ggcatggtgg	cttatgccta	taatctcagc	111360
actttttgat	gccaagggtg	aaggatcact	tgaggccagg	agttcaagac	cagcctgggc	111420
aacataagga	gaccctatct	ctacaaaaaa	acttaaaaat	tagctgggta	tagtgggtga	111480
tgctgaagt	cccaactact	caggaggctg	aggcagaagg	atcacattag	tccaggagtt	111540
cgaggctgca	gtaagtgtg	attacatcac	tgactccag	cctgggtgac	acagggaaac	111600
cctatctata	aaaaaattaa	attgagatga	aagcggactc	ctctaggaaa	aggtatgcta	111660
cttaggttca	aaataagaaa	agagcgtgga	caaataagaa	atgcctggaa	ttatatattga	111720
gtccctgcag	agaagacact	attttaagta	gcagaaattg	aaattttcct	gtttttgtgc	111780
tttaagagat	taaaacaattg	ttttattttc	ttcaaatagg	aatcgagtca	ctggcattta	111840
cgaactcagc	ttatgcaaaa	tgtcagacac	aggtagtcca	ggtaagctct	tgtggattga	111900
ggaggtgata	gttatctttg	tgtatgtttc	aagtatcctg	tatttaggct	gggcacgggtg	111960
gcacacacct	gtaatcccag	cgctttggga	ggccaaggcg	ggtggatcac	ctgagtcag	112020
gagttcaaga	gcagcctgac	caacatgggtg	agccccct	ttctaccaa	aatacaaaaa	112080
ttagccagg	gtggtggcgc	gtgcctgtaa	tccagctac	tcaggaggct	gaggcatgag	112140
aattgcttga	acctgggagg	cagaggttac	agtgaaccga	gattgtgcca	ctgcactgca	112200
gcctaggtga	cagagtgaga	ctcagtcctca	aaaaagaaaa	agaaaaaaa	atcaaattat	112260
taagtcagat	ttgttctgta	ttctttttta	tgtgttttta	aactgctctt	gttctcttag	112320
ctaggcggag	ttaagcagca	cttagcattt	ctatgtttgt	aaaccgaaac	atgaacaaaa	112380
gttaagggtt	tctaatagca	tttgattaat	gtgagtttga	gtcattttac	agaatataca	112440
tggaatacaa	gtcagggccc	acaggcaaga	cttagataag	ttaagaagac	tttgagtatg	112500
aatccaagta	gcacctaat	attatactat	ccagatagtt	tcttgaggaa	ttttttta	112560
taatataaat	tcatgtat	aatatgtaaa	accagcataa	tactaaaaaa	aattaacagg	112620
aaataataga	ggctaaagtt	ctctgtcatc	tgtcagccag	tgagctactg	agtgtgtgta	112680
tgtgtgtgtg	tattcccttc	cgctttttta	aaaattttta	tttttaggga	caggatcttt	112740
ttttttttt	ttttttttt	agacagagtc	tcgctgtgtc	acccaggctg	gagtggtgtg	112800
gcgccatctt	ggctcactgc	aacctctgcc	tcttgggttc	aagtgtattct	cctgcctcag	112860
cctcctgagt	agctgggttt	gcaagcacc	accaccatgc	ccagctaatt	tttgtatttt	112920
tagtagagat	gggttttcac	tatgttggtc	aggtgtgtct	cgaactcctg	acctcagatg	112980
atagcccacc	tcagcctccc	aaagtgtctg	gattacaggc	gtgagccatc	acgcccggcc	113040
ccatagtgtg	tagactacta	cagactat	aatatttctg	catttttcacc	taacatcata	113100
attttatcat	tttcttaaat	tgcattcccag	tcttcataat	tatcaaagaa	tgtaaaactc	113160
agtttttaaa	aacgttcagt	tgttttgcta	tttaaacaaa	atcccatata	tcttcagacc	113220
aaactgtctt	agtttgctac	gtataagaat	tattaattcg	tggccaggcg	cggtgggtca	113280
cgctgtaat	tccagcactt	tgggaggctg	aggcaggtg	atcacgaggt	caggagatcg	113340
agaccatcct	ggctaacaca	gtgaaacccg	gtctctacta	aaaatacaaaa	aaattagctg	113400
ggcatggtgg	cgggcacctg	tagtctcagc	tactcgggag	gctgaggcag	gagaatggca	113460
tgaaccagg	aggcggagct	tgcatggggc	tgagatcacg	ccactgcact	ccagcctggg	113520
caacagagcg	agactccatc	tcaaaaaaaa	aaaaaaaaga	attattaatt	cgttttgacc	113580
taaaacttga	attgaatttt	ttttctgaaa	caaaaacttg	tttctctctg	tcttaggtat	113640
gcagagaagg	agaagaaaaa	tcttagatac	gtcagtgga	tatgtgcggg	gagaagagaa	113700
cttagcaggc	tggcggcccc	gtggagacag	cctcatcctt	gagcaccagt	gggagctgga	113760
gaagctggag	ctcctacatg	aggtatccag	gggcagggtt	gttcagatgc	agaaactctc	113820
ggacaagatt	gccaaagtat	cagtcttctt	ccccgtgct	gcatgtgatc	atagcttttc	113880
acttttctct	cctgaggtct	taacgagctt	tgtgtttgct	atagcagtag	tattgatctt	113940
ctcaggtgga	aaaaacccgc	cactttttgc	tgctgcgtga	gagacttggt	gacagcatcc	114000
ccaaatccct	gagcgactcg	ttatccccc	gcctcagcag	tgggaccctc	agcacctcca	114060
ccagtatctc	ctctcagatc	tcaaccacta	cctttgaaag	cgccatcaca	cctagcgaga	114120
gcagtggtga	tgattcagga	gacatcgaaa	gcctggtgga	ccgagagaaa	gagctggcta	114180
ccaaggtgtg	aatcccttcc	tctttgctga	acgtcttccc	acaaggctcc	acaaactagc	114240
ctctcggttt	attcattttc	aacacctttg	ttcgaggtgt	ttgaaggcct	gtgataatac	114300
tgtgaatgta	gaaataaaaa	gacgcagttc	ctaccctcaa	caagtttaca	gggccaggca	114360
cagtgcctga	tgctatttat	cccaccactt	tgggaggctg	aagcaaagg	attgcttgag	114420
ccagggagtt	aaagacaagc	ccaggctaac	atagttagac	cttgtctcta	caaaaacaaa	114480
aattaggcca	ggcatgggtg	ctcacacctg	taatcccagc	actttgcagg	gctgaggcag	114540

gtggattacc	tgaggctctgg	agttcagagc	cagcctggcc	aacatggtga	aaccctgtct	114600
ctactaaaaa	tacaaaaatt	agccaggcgt	ggtggcgagc	gcctgtaatc	ccagctactc	114660
aggaggctga	ggcaggagaa	ttgcttgaac	ccaggaggcg	gagattgcag	tgagccgaga	114720
tagctccact	gtgctccagc	ctgagcgaca	gagtgacact	ccatctcaaa	aaataaaata	114780
aaataataaa	aaattaaaaa	attaaaaaat	ataaagggtt	cagaatatta	ggtcacaagc	114840
atataataga	gatgtgtgtt	tcaatgctat	gggagtcaga	ggagaagtgt	ctatggtggg	114900
gagctcatga	gctgaccttg	aaaggatgcg	caggagggag	gcagcagtg	cgacacctgc	114960
gtgatcagga	gggaggcagt	ggtgtcctgc	gtgatcagga	gggaggcagc	ggtgtcctgc	115020
gtgatcagga	gggaggcagc	ggtgtcctgc	gtgatcagga	gggaggcagt	ggtgtcctgc	115080
atgatcagtc	ttggggaatt	cacgtgggac	tggaacagga	ggtgtggagt	gaagaatcct	115140
gtatactgca	ctgaagagtt	tagagtggat	gatgggagca	aggcagaaat	atgttaaaca	115200
ggtattttgt	ttttagaaag	atcttctggc	aggggcgggt	gctcacgcct	gtaatccac	115260
cactttgaga	ggccgaggca	ggtggatcac	aaggctcagga	gattgagacc	atcctggcca	115320
acatggtgaa	accccggtgc	tactaaaaat	acaaaaatta	gccgaacgtg	gtggcggtgc	115380
cctgtagtcc	cagctactaa	ggaggctgag	gcaggagaat	cgcttgaacc	caggtggcgg	115440
aggctgcagt	gagccgagat	cgcgccactg	cactccagcc	tgggcaacag	agtgcagcac	115500
catctcaaaa	aatatatata	tttctctggc	agcaataaag	agaatggatt	gatggtgaag	115560
agatacactg	aggaaggag	aattctttat	ctttattaca	gtaatctgat	gagaaatgat	115620
aaaagctttt	taaaatttta	ttttttttga	gatggagtct	cactcctgtc	acacaggctg	115680
ggatgcagtg	gcatgatctt	ggctcactgc	aacctccact	tcccagggtc	aagcgattct	115740
ccttcctcag	cctcccaagt	agcttggagt	acaggcatgt	gccaccacac	ctggctaatt	115800
tttgtatttt	tagtagagac	agagtttcac	catggtggcc	aggctggtct	tgaactcctg	115860
acctcaggtg	atccacccgc	ctcagcctcc	caaagtgcga	ggattacagg	tgtgtggacg	115920
cccagccaga	taaaagtgtt	ttaaagaagg	cagaggtggc	tggtgtcggt	ggcacacgcc	115980
tgtaatccca	gcactttggg	aggctgaggt	gggaggattg	tgaggtcaag	agatcactat	116040
gatgttttta	ttttttattt	ttatctttat	taaatttttt	ttttttttga	gatggagtct	116100
tgctctgtgg	cccacgctgg	agtgcagtgg	tgcatctcgc	gctcactgca	gccttcactt	116160
cccaggttca	agcggttcct	ctgccttagc	ctcctgagta	gctgggatta	caggtgtgca	116220
ccaccatgcc	cagctcattt	tttgtagggt	ttttttttgt	ttgtttgttt	gtttttttga	116280
gaggggatgg	agtcttactc	tgctgccagg	ctagagtgcga	gtggtatgat	ctcagctcac	116340
tgcaacctcc	acttcccggg	ttcaagtgat	tctccttcc	cagcctcccg	agtagctggg	116400
attacagggtg	cccaccatga	cgcccagcta	atcttcttat	tttttagtagt	gatggggttt	116460
caccatgttg	gccaggatga	tctcgatctc	ctgacctcgt	gatccacctg	cctcagcctc	116520
ccaaagtgtc	gggattacag	gtatgagcca	tcacacccag	ccagaacat	ggtgttttta	116580
gtaattcact	ttggtaatct	tgagaaagg	ttttttaagc	cacttgcttt	atctttatta	116640
gacatacata	gaacttccct	tactatatac	tttctagagc	tgtttgggca	116700	
acatcgagc	tgactgtgg	gtttaggaca	tgttatttcc	cctcctgcct	tgtatccagg	116760
gcttcattta	gtcattgagg	ttttgtttg	tttgttttta	aatttactga	tttattagaa	116820
aggacagtgc	aaaggagacc	aatgaacggc	agatggacga	atgcataggg	agaggagatg	116880
tggaagcttc	atgccctccc	cggaactca	gcccttcagg	agcctccacg	tggtcagcta	116940
tcccaaggct	cctggctgtt	gacattttat	gttcagcttg	cacttctctt	tctcttcttt	117000
agtgcctgca	acttctcacc	cacactttca	acagagaatt	cagccagggtg	cacggcagcg	117060
tcagtgactg	taagggtgagc	acattgactg	taatttttag	ccagtatgtt	gataactgat	117120
ttctccacag	cagcccagat	tacctattcc	tggtttttgc	tgcttttaag	caactgttca	117180
tgggcatagc	attgtatttg	aaaatttatg	acatactgct	ctggtattca	ttctaatttt	117240
tcagagtccg	aacactgact	tctgaagata	aaagtactcc	tttgtgtctc	ttagagtgat	117300
tatcagatgg	gaaacatttt	ggctttttca	tgactccttt	ggaggagaat	attctatggg	117360
gaggtgggtat	gttattcttt	gccagggtac	aaggaaaccc	tgagggttct	ggtggcataa	117420
agttttatrr	acttcaacaa	agagtgaagt	aaacacttca	gagaatttct	gtgttattca	117480
ctcaattcta	gtcagcttga	ccttaaacc	tcccggtca	ttcctgcctt	gggtgctttg	117540
cacacctact	atcttcttgc	ctggaattgc	ctttctccag	gtattgccat	ggttggctcc	117600
cttacctctt	tcatttttct	actccattat	cacctcctgt	gagtcgactc	tgttaacacc	117660
tgataataaa	tctgacagct	gtctgaaact	ttctacctac	ccctctttcc	tgctttattt	117720
tttgctatag	cactaattac	tacctgatgc	tttttaag	gtttgttgag	tggtttattg	117780
tctgtcttaa	gctccaggga	ggcagatcct	ctgccatatt	attagaacaa	tgccatgtac	117840
atggaagggtg	caccactgta	ataccactta	tattaattca	ggggctgggt	gtggtgggtg	117900
atacctgtaa	tcccagcact	ttgggaggcc	aagatgagg	gatcgcttga	ggccaatagg	117960
ttgagaccag	cctgggcaat	atagcaciaa	tacctatct	ctacaaaaaa	atttttgaaa	118020
aattagccaa	atgtggtggt	acacacctat	agtcctagct	actcaggagg	ccaaggcagg	118080
attgcctgag	cccaggagtt	tgaggctgca	gtgaggtatg	attgtgccat	tgacccccag	118140
cctgtgtgac	agtatttttt	tgtctcttaa	tgataacgga	ctggggggct	ggctgcgggtg	118200
gctcatgcct	gtaatcccaa	cactttggga	ggccaaggcg	ggaagatcac	ctgaggtcgg	118260
gagtttgcga	ccagcctgac	caacatggag	aaaccccatc	tctactaaaa	atacaaaatt	118320

```

agccggatat ggtggcgcat gcctgtaata ccagctactc aggaggctga ggcaggagaa 118380
tcgcttgaac ctgggaggtg gaggttgagc tgagccaaga tcgtgccatt gcactccagc 118440
ctgggcaaca agagtgaac tccgcctcaa aaagaaaaca aaaaacaaaa aaacccccaa 118500
aataatgggc caggcatggc ggctcatgcc tgtaatccca acactttggg aggccgaggt 118560
gggcgatca cctgaggttg ggagtttgag accagcctga ccaacatgga gaaaccctgt 118620
ctctgctaaa aatacaaaat tagccgggcg tgggtggcaca tgcctgtaat cccagctact 118680
caggaggctg tggcagaatt gcttgaacct gggaggtgga ggttgcggtg agccaagatc 118740
gcaccactgc actccagtct ggggtgacaga gcgagacttc gtctctccaa aaataaataa 118800
agataatgca ggagaggcca gtgacagtgg ctacacatctg taatcctgcc agcttgggag 118860
gtcaaggctg atggatctct tgaggccaag agttcaagac gagcctgggc aacatgacaa 118920
aaaccccatc tccacaaaaa ttagttgggt gatcacttga gtcctggagg ctgaggctac 119040
tcccagctac tcaggaagct gagatgggag cctgactaca gcctggtaga catagcaaga ccttatctcg 119100
agtgagccaa catcgacca ctgcactaca cctgtctctt aaaattttac acaagatgaa 119160
aaaaagaaaa taataataat aataataaga agaagaagaa gaagaagaa acaataataa 119220
accttatctc gagaaagaaa ataataataa taataataat aataatgaca acaataataa 119280
taatgcagga gggcagtagg aacataaaaa accaaccttg ccccttttat aaatgtcttg 119340
attcctgggtg tggcacttag gccgtaaaat atttgtaggc ttgttttttt tcttttttga 119400
aaatattttc accagttgtg gtggtacacg cttgtagtcc caggtagtca ggaagccagg 119460
gcacaggagg atcgctggag cccaggagtt ggaggccaca ggaagctagg attgcatcac 119520
tgccctccag cctgggcaat agagcaaggc cttgtctctt aaaattttac ctagtattt 119580
atttatgtat aaattatgta taggcttgtt ttcttaaatg accttctgaa acaagatgta 119640
tcaaagttag gttctggttg ttgaaactcc ctattgttac ttagaggcca gcttttttca 119700
gctgaatgta acttgtagtg ttcggtttgc ttccagttgt ctgatattct tccaattgga 119760
cggtatccct ctgagtcag tttcagcagt gccaccctca ctccctcctc cacctgtccc 119820
tctctggtag actctaggag caactctctg gatcagaagt aagtaccag atttactga 119880
gagaagtcaa tctaagaacc aaggtaaatg tcaaccttcc tctagctcaa tggttcttga 119940
ctgaaggcag caccttctcc ctaggcagca tttggaaatg tgtgggagcc tttttggcta 120000
tcacagttag tggagaggtg ctagtacttg gcatctcttc acatcatagc ttactaacct 120060
tctgcatcat caactcatca tgactgtctt cttcatccct ttctgctagt tttatcctcc 120120
attcctaccc ctgcttgtct cccagtcatt aataccagac ataggaagtg ctatcttggg 120180
aaaaatattt tgaaactgtt ttggaaaaaa atattttgat tccacagcaa attatgtagg 120240
taagaaaaat aaagacaagg ccaggtgagg tggctcccag cactttcggg tgccaaggca 120300
ggaggattgc ttgaggctgg gagttcgaga ccaggctggg caacatagcg atatgtgtc 120360
tctacaaaaa ataaataaga taccagggcg tgggtgctg tgcctgtggt cccagctact 120420
cgggaggctg aagtgggagg atcgcttgaa tccaggagtt caaggttgca gtgaggtttt 120480
attattgcac cactgcattc caccctgggc gacagagcaa gaccctgtcc ctataaatcc 120540
taaaaaaag acaaatgtag tctagagtcc atgattcatc atcataatcc gttctttgca 120600
acacaccctc ttgtgcatct ccataaatta tgctcacctg gttatcatcc acctgtctat 120660
cttattgtac caacactgac acaggcaaag gtggtggag aaaataacac aaccaggtaa 120720
tttcattaaa tttaggatgc cacagactgt agggactatt ctgtgccact aagagaagaa 120780
gaaaaagcac tgacaactaa cttaagcatg ccatcaatta taagatgaat cccaatttaa 120840
gaaatggtgg ctgagtgcag tggctcatc ctgtaatccc agcactttca ggggctgagg 120900
tgaggaggatt gcttcagccc acaagtttga gaccagcctg ggcaacatag ggagacccc 120960
atctctacaa aaaataaata aaatagttgg acatgatggc atgcgcctgc ggttccagct 121020
actcaggagg ctacatggg aggattgctt gagcccgga ggttgaggct gcaatgagcc 121080
atggttgcac cattgcactc cagcctgggt gacagagcaa gaaatcagcc aggcataagg 121140
gcatgtacct gtaatccag ctactgggga ggctgaggca ggaagattgg ttgaacctgg 121162
gaggcggagg ttgcagttag cc

```

<210> 2
 <211> 10884
 <212> DNA
 <213> Homo sapiens

<220>
 <221> exon
 <222> 1601..1750
 <223> exon

<220>
 <221> exon
 <222> 2139..2331
 <223> exon

<220>
<221> exon
<222> 2540..2658
<223> exon

<220>
<221> exon
<222> 3830..8884
<223> exon

<220>
<221> misc_feature
<222> 8885..10884
<223> 3'regulatory region

<220>
<221> allele
<222> 2397
<223> 12-602-196 : polymorphic base C or T

<220>
<221> allele
<222> 2551
<223> 12-602-350 : polymorphic base A or C

<220>
<221> allele
<222> 4500
<223> 12-587-379 : polymorphic base A or C

<220>
<221> allele
<222> 6119
<223> 12-596-124 : polymorphic base A or G

<220>
<221> allele
<222> 10130
<223> 12-808-52 : polymorphic base A or G

<220>
<221> allele
<222> 10153
<223> 12-808-75 : polymorphic base G or C

<220>
<221> primer_bind
<222> 2203..2221
<223> 12-602.pu

<220>
<221> primer_bind
<222> 2600..2620
<223> 12-602.rp complement

<220>
<221> primer_bind
<222> 4479..4499
<223> 12-587.rp

<220>
<221> primer_bind

<222> 4858..4878
<223> 12-587.pu complement

<220>
<221> primer_bind
<222> 5996..6015
<223> 12-596.pu

<220>
<221> primer_bind
<222> 6423..6443
<223> 12-596.rp complement

<220>
<221> primer_bind
<222> 10079..10098
<223> 12-808.pu

<220>
<221> primer_bind
<222> 10523..10543
<223> 12-808.rp complement

<220>
<221> primer_bind
<222> 2378..2396
<223> 12-602-196.mis

<220>
<221> primer_bind
<222> 2398..2416
<223> 12-602-196.mis complement

<220>
<221> primer_bind
<222> 2532..2550
<223> 12-602-350.mis

<220>
<221> primer_bind
<222> 2552..2570
<223> 12-602-350.mis complement

<220>
<221> primer_bind
<222> 4481..4499
<223> 12-587-379.mis

<220>
<221> primer_bind
<222> 4501..4519
<223> 12-587-379.mis complement

<220>
<221> primer_bind
<222> 6100..6118
<223> 12-596-124.mis

<220>
<221> primer_bind
<222> 6120..6138
<223> 12-596-124.mis complement

<220>
 <221> primer_bind
 <222> 10111..10129
 <223> 12-808-52.mis

<220>
 <221> primer_bind
 <222> 10131..10149
 <223> 12-808-52.mis complement

<220>
 <221> primer_bind
 <222> 10134..10152
 <223> 12-808-75.mis

<220>
 <221> primer_bind
 <222> 10154..10172
 <223> 12-808-75.mis complement

<220>
 <221> misc_binding
 <222> 2385..2409
 <223> 12-602-196.probe

<220>
 <221> misc_binding
 <222> 2539..2563
 <223> 12-602-350.probe

<220>
 <221> misc_binding
 <222> 4488..4512
 <223> 12-587-379.probe

<220>
 <221> misc_binding
 <222> 6107..6131
 <223> 12-596-124.probe

<220>
 <221> misc_binding
 <222> 10118..10142
 <223> 12-808-52.probe

<220>
 <221> misc_binding
 <222> 10141..10165
 <223> 12-808-75.probe

<400> 2

gcctcccagg	ttcacgccat	tctcctgctt	cagcctcccg	agtagctggg	actacaggtg	60
cccgccacca	caccagcta	attttttatt	ttattttatt	tattttattta	cttttgagat	120
ggagttttgc	tctcgttgcc	cacgccagag	tgtagtggtg	caatcttggc	tactgcaac	180
ctccaccttc	cagcttcaag	tgattctcct	gcctcagcct	ccaaagtagc	tgggattaca	240
ggcaccgcc	accatgccca	gctatttttt	tttttatttt	tagtagagat	cgggtttcac	300
cattttggtc	aggctagtct	caaactgctg	accttatgat	ccaccgcct	cgacctcca	360
aagtgtctgg	attacaggcg	tgagccactg	caccggcca	caccagcta	attttttgta	420
tttttagtag	agatggggtt	tcaccgtggt	agccaggatg	gtcttggtct	cctgacctca	480
tgatctgcc	gcctcagcct	cccaaagtgc	tgggattaca	ggcgtgagcc	actgcgccgg	540
cctttttttt	ttttttttga	gacagagtct	cactctgtcg	cccaggctgg	agtgtgcaa	600
cctctgcctc	ccgggttcaa	gtaattctcg	tgccctagcc	tcctgggtag	ctgggattac	660
aggggcctgc	caccacgccc	agctaatttt	tgtattttta	gtagagacgg	ggtttcgcca	720

tggttggtcag	gctggtctca	aactcctgac	ctcaggtgat	ctgcctacct	cggcctccca	780
aagtgtctggg	attacaggtg	tgagccactg	cacctggcta	acttgatttc	aaaaaaaaaa	840
aagaatgaag	tatgagaaaa	agttaaaatg	ggcttttttag	atttagtgag	atacaatgct	900
atgttactct	acaaagatat	ctagcaatac	agagtatagg	aattttatttc	tcacacagta	960
gtcaagaggg	ctggtccagg	tccaggtcag	tggtgcaggt	cttctctgca	cagccgttca	1020
ggaaccctgg	gtcttcccat	ttgtttgctt	ttctcgccat	tttctaaggc	agtgggcctc	1080
tgagacttca	gtgtacatca	tcatcccctg	ggaggcttgt	tgaagccccc	atggctcaac	1140
cccacccaaa	gtttctcggt	ccgtaggtct	ggcttggagt	ctaaaaattt	gagtttttga	1200
acagtttccc	aggtgaagct	gatgctgctg	gctgggagca	cactttgaga	aacatcgctt	1260
ttggtgtggc	catgggacta	aaaagaaaca	tataaaggag	tagagttcct	attccagata	1320
atgtgttcag	agggctttaa	tataggcaga	gagatttgtt	gacccagtg	ccatcaatga	1380
tcaatttctt	ttcataactt	cagaaggatg	acaccagttg	aacgcagagt	taactttctg	1440
aagccagctt	gtctcctcag	ctgagctctc	tgcaactcaa	gtttgcagct	ggggttttag	1500
gagggccatt	gtgttctctc	cagtgaacaa	gtactcagta	tgccctgatt	gtaactgatt	1560
ctcttggtac	ccttttcttt	ctaactctct	tatttttaag	gacccagaa	gccaattccc	1620
gggcctctag	tccctgcccc	gaatttgaac	agtttcagat	tgtcccagct	gtggaaacac	1680
catatttggc	ccgagcagga	aaaaacgaat	ttctcaatct	tgttccagat	attgaagaaa	1740
ttagaccaag	gtgagtacta	tattgagcag	gaatgccagc	tataaaaaac	aaatccacag	1800
gaagaagtga	ctggccagct	cttccctgtg	aggctgttac	tgtagtctga	tgcaatctgt	1860
actgtagtct	gatgcaagtt	gagtcgtggg	tgagagggcc	agcctgggtt	tctccagttg	1920
ggtctcataa	tgcatctgca	ccctggcgca	gcatgttgcc	atggcacggt	ttcgcttttg	1980
ccgtattact	ttggcagata	agattctagg	ccaaataggt	catttgcctg	tttcatcgaa	2040
tctaaggact	tggectaagt	gtggcgtatt	ttgatgtgcc	tcagcacgat	tattttcttt	2100
ttgtgagaaa	ctaacttttg	tcatattgtc	gttttttagc	cagtggctct	taagaaagga	2160
taccttcatt	tcaaggagcc	tctttacagt	aactgggcta	aacattttgt	tgtcgtccgt	2220
cggccttatg	tcttcatcta	taacagtgac	aaagaccctg	tggagcgtgg	aatcattaac	2280
ctgtccacag	cacaggtgga	gtacagtga	gaccagcagg	ccatggtgaa	ggtccgtcct	2340
gccctgcctt	ggtttcttat	tgccacgtgt	gccctctct	tttgatttct	gcactcyctg	2400
ttacgtgggtg	tgaaatttcc	ctattctctc	tggttttgaa	cttcccttgt	gaagtgtctat	2460
atgccttggg	aatatggcaa	ggcagtcctc	attgctgtct	ctgtagtaac	tttcttgtct	2520
acctgcattt	ttctttcaga	caccaaacac	mtttgctgtc	tgacacaaag	accgtggggt	2580
ccttttgcag	gccctcaatg	acaaagacat	gaacgactgg	ttgtatgcct	tcaaccact	2640
tctagctggc	acaatacggg	aagaagtgtt	gtttgtgttg	ttgttgtttt	tgagacggag	2700
tctcactttt	tctccctgaa	gtgcagtggc	ttggtcttgg	ctcactgcaa	cctccgcccc	2760
ctggttcaag	cgattctcct	gactcagcct	ccccagtagc	tgagactaca	ggcacgtgcc	2820
accattgcct	ggctaatttt	cgtatttttt	gtagagacgg	ggttttacca	caatggccag	2880
gctggtctca	aactcctgac	ctcaagtaat	ccaccacct	cagcctccca	aagtgtctgg	2940
attacaggcc	tgagccactg	caccagcct	agaagtgtgt	ctgtgttttc	tttctctctt	3000
tggtcttttg	cttcaggaat	atcaggataa	ataggaaagg	aagaattttt	cttttctttt	3060
ttcttttctt	gttttttttt	tttttttttt	tttttggaa	ggagtgtttt	ctcttgtgtg	3120
gcccaggctg	gagtgcagtg	gcatgatata	ggctcacccg	aacctccacc	tcccagggtc	3180
aagccattct	cctgcctcag	tctcctgagt	agctaggatt	acaggtgccc	gccaccacac	3240
ccagctaatt	tttttttttg	tatttttagt	agagatgggg	tttcaccatg	ttggccaggc	3300
tggtcttgaa	ctcctgacct	caggtgatcc	acctcctc	gcctcccaaa	gtgctgggat	3360
tacaagcgtg	agccaccaca	tctggcttga	aaggaagtat	ttttcaaaat	aaatttaaaa	3420
gagaggaagg	acaggagtgt	aaattgggtg	tggggtggta	atgaggcctc	tgcatcttgt	3480
aaggcattgt	catgttctga	accaccagg	atactctggt	cacacatgtg	caccaattat	3540
tttccctctt	tgtctttttg	ttctcaagag	ttctaaagat	ggggctcagg	ggagagagta	3600
aagggaggac	agtgagaacc	ttaatggtct	cccaacctcc	agccctgtga	ttcagcctca	3660
cgttctgagg	gcttgtgtca	caggggtccc	gagtagacgg	cttagaggac	taggcctgcg	3720
gtgccaataa	ggaagatgct	ccaggaggga	gcctcagagg	gagaggactg	tgaggggaca	3780
agacttgtac	ccagacttct	aatectacct	ccccgtttgt	cccccatagg	tcaaagcttt	3840
cccgagatg	cccgagccag	tcgaaatact	aagtgaactc	gccgagtgcc	ctcactcgcc	3900
ttcgagagat	aaagaaaagc	ttacctctca	ttctcttttg	tgattcttga	cgggtgactct	3960
tgtatgtaat	cctgtggctt	aactacttct	ccctccttgt	ccagcacttt	tctagctctc	4020
ccgttcccca	tctccattgc	tctgtactct	tttctttttt	cttgtgtctga	gaatctcgtt	4080
agtagcatgt	ggcctaacaa	aaggaaaaaa	tgttttttaa	cacacacaca	cacacacaca	4140
cacacacaca	catacacaga	caaaaacaca	aaaactctga	ggggatctgg	tgaatctcca	4200
aattattgtg	ggtgtacttt	ggcttccttt	tgtatgatag	gtccccatca	tgaccacctc	4260
tgtgtctgtg	gctgctgtca	ccaggcacct	ttgtttttca	agacaacata	cttttttttt	4320
cttttctctg	tttgtgatat	cactttaatt	ttcttgggtg	ggcttagaga	ctaagggagg	4380
agacatctgg	ccttttttaga	acctgagagg	aaaaaaagag	tctttttttc	ccctctgtct	4440
ctttttgcca	tggctaattcc	ctgcatttcc	attcagggaa	aaggtggtag	tgagcatagm	4500

actgcaacag	ttatattctg	agtcaaagtt	ggggcttttt	acggcataat	tatggaattt	4560
ttattttactg	gtagagagga	gacgagaggc	tttttcagtg	ggcctgggac	agtggctgct	4620
cttgacttttg	tgtgaaggga	aatgccaagg	atgcttctgg	tggacttcag	gggaccccag	4680
ggtttggccg	tgggccgtga	tggcagcagg	cgggtgggatg	cttgtagctc	ctcacagcag	4740
gattcctgcc	cactgttttt	tctctgttgg	gaggggaagct	cttttctagg	agtgtctcag	4800
ttctgctttt	ggcattagtg	atgggtgggtg	tacagttgga	attagtgccca	tgtcatcacac	4860
aaatgttcca	caaggcggga	gtgtttcact	ttctgggtgat	aaacttgatg	gtcattgtta	4920
tgattaagat	aatgccgggc	aggccgggca	cagtggctca	cgcctgtaat	ccaagcactt	4980
ggggaggccg	aggcgggcag	atcacgagat	caggagtcca	agaccagcct	ggccaatatg	5040
atgaaacccc	gtctctacta	aaaatacaaa	attagtgcggg	tatggtggca	catgcctgta	5100
attccagctg	cttgggagcc	tgaggcagga	gaactgcttg	aaccaggag	gcagaggttg	5160
cagtgaacca	agatcgcgct	attgcactcc	agcctgggtg	acagagcaag	actctgcctc	5220
agaaaaaaa	aataataata	atgctgggta	gtgaccttgt	gattgttaca	gctccctttg	5280
atcaaagaaa	tatagctttc	aggcataaac	cttgaagtct	ccctctgaat	ccagcagttg	5340
ttttcattga	tgtttgtcag	gttgaagatg	ctttcagtg	tgctctctat	actcataaat	5400
aagcaaatgt	ggcaggcttt	gctttctgga	tcccaggatt	aaaactaacc	gtgaccacta	5460
ctccaaacaa	aacacaatat	gcctaggggc	acggatgaac	gtccagggag	cccggggccc	5520
aggctttgtt	gcgtgttccc	tgtctctctc	catctggtgt	ggaaacactg	cccaggggaga	5580
aaggaggag	ctcactgtgg	acagtcttct	ttccttctga	cagaccaggt	catctggctt	5640
ccgagatcat	cagagaagat	aagtctgtct	ctttcagctg	ccagtaagtt	ttccaggatg	5700
agaggggaaa	aagaaagcct	ccagtgaact	cagttgcttt	gccagttgtc	ttgggattgt	5760
tttacaccat	cctttacttc	ccttgctcag	acctctctgt	ttcaccattg	ctcaggcatt	5820
caggaaagta	tctgtctact	cccacttggt	gagtcctcgg	ccttgagggt	gctgactctc	5880
aggcgttagg	cagctggatg	acttcccgtc	tcacgcagca	aaggccaggg	gcttgccgct	5940
ctctgcagag	ttgttgctag	ggagacttgt	gtcatcatcc	acaaccttgt	ttctcacttc	6000
ctggttgggc	tcatctctga	agaacaggtc	tcccagcttc	gctccttatc	actgcattgt	6060
gaagaggagg	aaaagtgaat	cacggagaga	gaaaggaaag	gatagaatca	caggctgcrt	6120
ctgcacctga	aaagtgaccc	gcggaaactc	tatggcggat	ttttttttta	actttcttct	6180
tcctgttaaa	acataggtca	ctaactgtga	tgttaattgt	tttctaagtg	gtatgtgaga	6240
ttttctaata	tagttagaag	tttcattgtc	tgatggacac	aatatgccct	tccggttcta	6300
ttcaaaccag	caggatctgt	cggtgcttag	agatggctgc	ctggactgga	atcaaactta	6360
atttcaggga	aatgaagatg	gaatttgaag	gtcactttta	aaattaagtc	attgatgctg	6420
ctgttacaga	gtgtgacaga	ggatccatgt	ctgtgacaca	ggacggtggg	aagcctgaga	6480
gagagtgaaa	ttatgtgata	cactgaaatg	acttttgttt	ttcttctaac	tcatacaaaa	6540
ctggtttgga	aagtctttgc	tttggaagcg	tcagacatta	gaacaggcca	aactggactg	6600
tctgttcata	gcgtgcctga	ataagaaggc	ctcttaggga	gccagaggga	gcagagtggg	6660
cgtgtcctgc	gtgctcttca	ccctctgggg	cgccctgct	gcggctggca	ggtgcagaca	6720
gcctttgctg	gtccccagca	cgtccagggt	gggtgctccc	ttgcccagca	gaaccatccc	6780
cactgtgagg	ctgtgagaga	tttgtggcag	gaactgttta	tgaggctcta	gttgttgcctg	6840
ttgtggcggg	aaagttaaga	aacatagccc	ttaaggaaac	cacctttatg	tattttctta	6900
aagcacgcct	ttaaataagc	aaaaacttta	aaaggcagga	aagagaattc	ttaggcaaat	6960
tcagagaaat	aagtgtagtg	taataactaat	cacctctctc	tctgtctctc	atcctccttt	7020
ctcccatcaa	agcaaaatat	ggcctcacca	ccagccccc	atcagtgtctc	agacctctc	7080
tgtgtctgtg	tgccctcctg	ggagtcaatc	agcgtcagg	ccaggactgt	gcaggggccag	7140
ccagcccatg	cgctagtcag	gagcacaggc	aagggggtgt	tgtggcagtg	gccggggcacc	7200
tgagccccag	ctcgttggtta	aacgtgtctga	cggcaagggg	caatggagtg	agtttcccaa	7260
ctaagaaacc	actatttatat	attttttccc	ttcagtcaca	tagacttcag	acaactctcc	7320
tattttttat	ggatttttca	gctcatttca	gatgaaggaa	ctaagtcatt	gtgaactgtc	7380
tcttgagatc	taaaaacaag	atgacttttc	ctggcacata	ttccaaagca	aagactttgt	7440
tgcctgctgc	ttattgtcta	atttacaggg	atattttaatt	ttgtcaggtc	tatgtatatt	7500
tatccagcta	tacttacttg	cacagtggat	tggagagaaa	ggattctcca	gtgtgcacac	7560
tcacgggtac	tctttctgca	tttccctcgt	gctgtgtccc	gctcgggttc	caatggacag	7620
tatcagggtc	tgtttgactt	aggtctttca	gttttccttt	cggttccctt	ttaaaaaatgt	7680
gattgttaac	ctgcctcttg	aaagattcaa	ccgggtgtgg	tggtcacgc	ctgtaatccc	7740
agcacattgg	gaggccaagg	caggtggatc	acctgaggtc	agaagtttga	gaccagcctg	7800
gccaacatgg	tgaaccccg	tctctactaa	aaatacaaaa	agtagccagg	cgtggtggcg	7860
tgtgcctgta	gtcccagcta	cttgggaggg	tgaggcagga	gaatagcttg	aacctgggag	7920
gtggaggctg	cagtgaagtg	agactgcacc	attgcactcc	agcctgggtg	acaaagcagg	7980
actccgtctc	aaaaaaaaaag	aaaagattca	tgatgtctgt	gctcccagaa	ggtttgctgg	8040
atgtgtttac	ataggactct	aacttgtgtg	cactacagtt	gttcaccagg	gccagtgatt	8100
cacccacgtg	tgtggccaga	ccatgactgt	gtagcaggaa	tgttttaatt	tgtgttccct	8160
tagtaaatg	aaatatcagc	tgagagatta	tttgtctgtg	ttattcaaaa	ggccatttat	8220
gaagttagta	tttgagcccc	ataagatctt	taaaaagcct	ccaatcattt	aaaggaagaa	8280

atcagagttg	ctataaaatt	cagtaaaaag	ctcatagcca	aacggctgtg	ctcagatgga	8340
aagtctgagc	tgaggttggt	ctcttgccaa	accgtggctg	ttgtgtgttg	ttcttcatgt	8400
cttcgagttc	atTTTTtttc	attctgccta	ttctggcatc	agctcacttg	aggagtccct	8460
cagccttctt	gtatttaagg	catcgtctta	gactttgttg	ctctaaagta	cctgtctgtt	8520
gagatttcaa	gtctcttgtc	accatcctca	cacatgacaa	caaaacccat	aatgcataag	8580
tggccttttt	gaaccaagac	tttgcaaaact	gatctctccc	ccgtgaagga	gttgagcaca	8640
ttagcaacaa	tgtacattaa	ttttggattt	tcattttcat	gttttatttt	gtaaatatta	8700
tctgatgttt	ggagcttgag	tatacagact	gtaaatatag	ttcttgtatt	tgtactaatt	8760
ctgattcttt	tgctgtatag	ccttagatgt	gcaatgcaga	cactatctaa	ctgtgtgttg	8820
taaccttgcg	tcacggagct	gttagtgaac	gaggtaaaaa	taataaaggt	acagccagtg	8880
catcagaagg	ttctcgatgt	gcattttattc	ttgcaccctt	tgaaaggtaa	ttgcacaaag	8940
atttcttttc	ttgatttgca	aagataatac	ctttctaaga	cagaagtcac	gatatcatcg	9000
cttaagtatt	cctttctgat	attcaaaate	gtggttttta	tataaccaag	aaagctaata	9060
tggtgctgtt	tctttgaaag	tacttttctc	ccaaagatgt	aagtggttat	tttagctaga	9120
agatttggtg	tttttctcct	gaagcctcct	tgtatgcctt	ccaataattc	tgtagtgttg	9180
tgttggcttc	tttttttttt	tttttttctt	tttgagacag	agtcttgctc	tgtcacccat	9240
gctggagtgc	agtgatgtga	tctcgggtca	ctgcaacctc	tgctcctggg	gttcaagcga	9300
ttctcctgcc	tcagactcct	gagtagctgg	gattacaggc	gtgcgccacc	atgcgtggct	9360
aatttttgta	ttattattat	tattattatt	attattattt	tatttattta	tttatttatt	9420
ttgagatgga	gtctcactct	gtcgtccagg	ctggagtaca	gtggcatgat	ctcggctcac	9480
tgcaactttc	cacctccogg	gttcaagcga	ttctccctgc	ctcagcctcc	caagtagctg	9540
ggattacagg	catccatcac	gcctggctga	ttttattttt	agtagagagg	ggtttcactg	9600
tgttggccag	gctggtctca	agctcctgac	cttgtgatcc	acccgcctca	gcctcccaaa	9660
gtgctaggat	tacaggcttg	agccaccgtg	cccaacctat	aattattttt	ttgagatgga	9720
gtttgactct	tgtcacccat	gctggtgtgc	aatggcacca	tctcagctca	ctgcaacctc	9780
tgctcccca	gttcaaacaa	ttctcttttc	tcagcctccc	aggtagttgc	aattacaggt	9840
gcccgccacc	acacctggct	aattttttaa	aatattttta	gtagagacgg	ggtttcagca	9900
cattggccag	actggtctca	aactcctgac	ctcaggtgat	ccgcccacct	cggcctccca	9960
aagtgtctgg	attacaggcg	tgagccaccg	tgcttgccca	tggtgggttt	ttttgggttg	10020
ggaagggtaa	atggggattt	gaaaagtga	acatgtcaat	ttaatttaac	aagccctcat	10080
cacgagcaga	gcatgagcag	gggccactgt	cctgggtgtt	tgggggagaar	cagaagaagg	10140
gagggaggag	casccctcag	gttaaaagca	atgatttgtt	ccaaccaata	tttgagtacc	10200
cacctgtgtc	ctggtgtgct	gctgggtgca	gtaattcagc	agccaacaaa	gcaagcccat	10260
gcacttgtgc	cctcccattc	cagtggggaa	gacaatgttt	aaacaaattt	acaacatcgg	10320
tcctaagtat	tttgagggaa	agcagctaaa	taaaaggatg	aaagagggat	gggaacgagg	10380
gatgtttaga	tggcatagtc	aggggaaggcc	tctcttcgga	gatcacattt	gagcagactc	10440
tagaatgaaa	ggatagggtg	agccatgcaa	acccctgggc	aaagaggtct	ccaagtagag	10500
aatgaagcc	cttgaggcag	cagtgtcctc	agtaggttcc	agaaatagca	aggtcgctgg	10560
tacagagtca	gggactggga	gcgcaataga	agacatcaga	aagtagccag	ggaccaggat	10620
atttagggct	taggtgagga	cttggaattt	cattagagtg	aatgggaagc	cattgaaggc	10680
tttgaggcag	aggagtcaca	ggggcttatt	tacattttta	aaagaccatt	ctggaaaaca	10740
gtctttgtac	tcaggcaggg	acaagtggaa	aaaaggggca	aaataagatt	gttaccacag	10800
tattccaggg	gaggagacgg	tggtctcagac	taaggtagta	gagtataggt	gggacgaaat	10860
gggcagattc	aaggtacatt	ttga				10884

<210> 3
 <211> 10682
 <212> DNA
 <213> Homo sapiens

<220>
 <221> 5'UTR
 <222> 1..186

<220>
 <221> CDS
 <222> 187..5637

<220>
 <221> 3'UTR
 <222> 5638..10682

<220>

<221> polyA_signal
<222> 10631..10636

<220>

<221> allele

<222> 471

<223> 12-809-119 : polymorphic base G or C

<220>

<221> allele

<222> 5487

<223> 12-602-350 : polymorphic base A or C

<220>

<221> allele

<222> 6265

<223> 12-587-379 : polymorphic base A or C

<220>

<221> allele

<222> 7887

<223> 12-596-124 : polymorphic base A or G

<220>

<221> misc_feature

<222> 2,4,5924,7850

<223> n=a, g, c or t

<400> 3

```

gngnccgctg ctcgcgctga ggtgcgtccg gtgcccggcc ccccgcgccc ccgcgcgcgcg      60
cggtcctgtg tgaccggtc cgcccgctcg tctgcagcgc ggctgaggaa acttggtgtg      120
aacttcaaaa gaagatttga ttctttatct ctggactgca tatatatata taacaaggcc      180
attaaa atg tcg gga gcc tca gtg aag gtg gct gtc cgg gta agg ccc      228
      Met Ser Gly Ala Ser Val Lys Val Ala Val Arg Val Arg Pro
      1          5          10
ttc aat tct cga gag acc agc aag gaa tcc aaa tgc atc att cag atg      276
Phe Asn Ser Arg Glu Thr Ser Lys Glu Ser Lys Cys Ile Ile Gln Met
15          20          25          30
caa ggc aac tcg acc agt att att aac cca aag aat cca aag gaa gct      324
Gln Gly Asn Ser Thr Ser Ile Ile Asn Pro Lys Asn Pro Lys Glu Ala
35          40          45
cca aag tcc ttc agc ttc gac tat tcc tac tgg tct cat acc tca ccc      372
Pro Lys Ser Phe Ser Phe Asp Tyr Ser Tyr Trp Ser His Thr Ser Pro
50          55          60
gaa gat ccc tgt ttt gca tct caa aac cgt gtg tac aat gac att ggc      420
Glu Asp Pro Cys Phe Ala Ser Gln Asn Arg Val Tyr Asn Asp Ile Gly
65          70          75
aag gaa atg ctc tta cac gcc ttt gag gga tat aat gtc tgt att ttt      468
Lys Glu Met Leu Leu His Ala Phe Glu Gly Tyr Asn Val Cys Ile Phe
80          85          90
gcs tat ggg cag act ggt gct gga aaa tct tat aca atg atg ggt aaa      516
Ala Tyr Gly Gln Thr Gly Ala Gly Lys Ser Tyr Thr Met Met Gly Lys
95          100          105          110
caa gaa gaa agc cag gct ggc atc att cca cag tta tgt gaa gaa ctt      564
Gln Glu Glu Ser Gln Ala Gly Ile Ile Pro Gln Leu Cys Glu Glu Leu
115          120          125
ttt gag aaa atc aat gac aac tgt aat gaa gaa atg tct tac tct gta      612
Phe Glu Lys Ile Asn Asp Asn Cys Asn Glu Glu Met Ser Tyr Ser Val
130          135          140
gag gtg agc tac atg gaa att tac tgt gaa aga gta cga gat ttg ctg      660
Glu Val Ser Tyr Met Glu Ile Tyr Cys Glu Arg Val Arg Asp Leu Leu
145          150          155
aat cca aaa aac aag ggt aat ttg cgt gtg cgt gaa cac cca ctt ctt      708

```

53

Asn	Pro	Lys	Asn	Lys	Gly	Asn	Leu	Arg	Val	Arg	Glu	His	Pro	Leu	Leu	
160						165					170					
gga	ccc	tat	gtg	gag	gat	ctg	tcc	aag	ttg	gca	ggt	act	tcc	tac	aca	756
Gly	Pro	Tyr	Val	Glu	Asp	Leu	Ser	Lys	Leu	Ala	Val	Thr	Ser	Tyr	Thr	
175						180					185				190	
gac	att	gct	gac	ctc	atg	gat	gct	ggg	aac	aaa	gcc	agg	aca	gtg	gca	804
Asp	Ile	Ala	Asp	Leu	Met	Asp	Ala	Gly	Asn	Lys	Ala	Arg	Thr	Val	Ala	
				195					200					205		
gct	aca	aac	atg	aat	gaa	aca	agt	agc	cgt	tcc	cac	gct	gtg	ttt	acg	852
Ala	Thr	Asn	Met	Asn	Glu	Thr	Ser	Ser	Arg	Ser	His	Ala	Val	Phe	Thr	
			210						215					220		
att	gtt	ttc	acc	cag	aag	aaa	cac	gat	aat	gag	acc	aac	ctt	tcc	act	900
Ile	Val	Phe	Thr	Gln	Lys	Lys	His	Asp	Asn	Glu	Thr	Asn	Leu	Ser	Thr	
		225					230					235				
gag	aag	gtc	agt	aaa	atc	agc	ttg	gtg	gat	cta	gca	gga	agt	gaa	cga	948
Glu	Lys	Val	Ser	Lys	Ile	Ser	Leu	Val	Asp	Leu	Ala	Gly	Ser	Glu	Arg	
	240					245					250					
gct	gat	tca	act	ggt	gcc	aaa	ggg	act	cga	tta	aag	gaa	gga	gca	aat	996
Ala	Asp	Ser	Thr	Gly	Ala	Lys	Gly	Thr	Arg	Leu	Lys	Glu	Gly	Ala	Asn	
	255				260					265					270	
att	aat	aag	tct	ctt	aca	act	ttg	ggc	aaa	gtc	att	tca	gcc	ttg	gcc	1044
Ile	Asn	Lys	Ser	Leu	Thr	Thr	Leu	Gly	Lys	Val	Ile	Ser	Ala	Leu	Ala	
				275					280					285		
gag	gtg	gat	aac	tgc	act	agc	aag	agt	aaa	aag	aag	aag	aaa	aca	gat	1092
Glu	Val	Asp	Asn	Cys	Thr	Ser	Lys	Ser	Lys	Lys	Lys	Lys	Lys	Thr	Asp	
			290					295					300			
ttt	att	ccc	tac	agg	gat	tct	gta	ctt	act	tgg	ctc	ctt	cga	gaa	aat	1140
Phe	Ile	Pro	Tyr	Arg	Asp	Ser	Val	Leu	Thr	Trp	Leu	Leu	Arg	Glu	Asn	
		305					310					315				
tta	ggt	ggc	aat	tct	cgg	act	gca	atg	gtt	gct	gct	ctg	agc	ccc	gcg	1188
Leu	Gly	Asn	Ser	Arg	Thr	Ala	Met	Val	Ala	Ala	Leu	Ser	Pro	Ala		
	320				325					330						
gat	atc	aac	tac	gat	gag	act	ttg	agc	act	ctg	aga	tat	gca	gat	cgt	1236
Asp	Ile	Asn	Tyr	Asp	Glu	Thr	Leu	Ser	Thr	Leu	Arg	Tyr	Ala	Asp	Arg	
	335				340					345				350		
gca	aaa	caa	att	aaa	tgc	aat	gct	gtt	atc	aat	gag	gac	ccc	aat	gcc	1284
Ala	Lys	Gln	Ile	Lys	Cys	Asn	Ala	Val	Ile	Asn	Glu	Asp	Pro	Asn	Ala	
				355					360					365		
aaa	ctg	gtt	cgt	gaa	tta	aag	gag	gag	gtg	aca	cgg	ctg	aag	gac	ctt	1332
Lys	Leu	Val	Arg	Glu	Leu	Lys	Glu	Glu	Val	Thr	Arg	Leu	Lys	Asp	Leu	
			370					375					380			
ctt	cgt	gct	cag	ggc	ctg	gga	gat	att	att	gat	att	gat	cca	ttg	atc	1380
Leu	Arg	Ala	Gln	Gly	Leu	Gly	Asp	Ile	Ile	Asp	Ile	Asp	Pro	Leu	Ile	
	385				390					395						
gat	gat	tac	tct	gga	agt	gga	agc	aaa	tat	ctg	aaa	gat	ttt	cag	aac	1428
Asp	Asp	Tyr	Ser	Gly	Ser	Gly	Ser	Lys	Tyr	Leu	Lys	Asp	Phe	Gln	Asn	
	400				405					410						
aat	aag	cat	aga	tac	ttg	cta	gcc	tct	gag	aat	caa	cgc	cct	ggc	cat	1476
Asn	Lys	His	Arg	Tyr	Leu	Leu	Ala	Ser	Glu	Asn	Gln	Arg	Pro	Gly	His	
	415				420					425				430		
ttt	tcc	aca	gca	tcc	atg	ggg	tcc	ctc	act	tca	tcc	cca	tct	tcc	tgc	1524
Phe	Ser	Thr	Ala	Ser	Met	Gly	Ser	Leu	Thr	Ser	Ser	Pro	Ser	Ser	Cys	
				435					440					445		
tca	ctc	agt	agt	cag	gtg	ggc	ttg	acg	tct	gtg	acc	agt	att	caa	gag	1572
Ser	Leu	Ser	Ser	Gln	Val	Gly	Leu	Thr	Ser	Val	Thr	Ser	Ile	Gln	Glu	
			450				455						460			
agg	atc	atg	tct	aca	cct	gga	gga	gag	gaa	gct	att	gaa	cgt	tta	aag	1620
Arg	Ile	Met	Ser	Thr	Pro	Gly	Gly	Glu	Glu	Ala	Ile	Glu	Arg	Leu	Lys	
	465					470						475				
gaa	tca	gag	aag	atc	att	gct	gag	ttg	aat	gaa	act	tgg	gaa	gag	aag	1668
Glu	Ser	Glu	Lys	Ile	Ile	Ala	Glu	Leu	Asn	Glu	Thr	Trp	Glu	Glu	Lys	
	480					485						490				
ctt	cgt	aaa	aca	gag	gcc	atc	aga	atg	gag	aga	gag	gct	ttg	ttg	gct	1716

54

Leu	Arg	Lys	Thr	Glu	Ala	Ile	Arg	Met	Glu	Arg	Glu	Ala	Leu	Leu	Ala	
495					500					505					510	
gag	atg	gga	gtt	gcc	att	cgg	gaa	gat	gga	gga	acc	cta	ggg	gtt	ttc	1764
Glu	Met	Gly	Val	Ala	Ile	Arg	Glu	Asp	Gly	Gly	Thr	Leu	Gly	Val	Phe	
				515					520						525	
tca	cct	aaa	aag	acc	cca	cat	ctt	gtt	aac	ctc	aat	gaa	gac	cca	cta	1812
Ser	Pro	Lys	Lys	Thr	Pro	His	Leu	Val	Asn	Leu	Asn	Glu	Asp	Pro	Leu	
				530				535							540	
atg	tct	gag	tgc	cta	ctt	tat	tac	atc	aaa	gat	gga	att	aca	agg	gtt	1860
Met	Ser	Glu	Cys	Leu	Leu	Tyr	Tyr	Ile	Lys	Asp	Gly	Ile	Thr	Arg	Val	
				545			550								555	
ggc	caa	gca	gat	gct	gag	cgg	cgc	cag	gac	ata	gtg	ctg	agc	ggg	gct	1908
Gly	Gln	Ala	Asp	Ala	Glu	Arg	Arg	Gln	Asp	Ile	Val	Leu	Ser	Gly	Ala	
				560		565									570	
cac	att	aaa	gaa	gag	cat	tgt	atc	ttc	cgg	agt	gag	aga	agc	aac	agc	1956
His	Ile	Lys	Glu	Glu	His	Cys	Ile	Phe	Arg	Ser	Glu	Arg	Ser	Asn	Ser	
575					580					585					590	
ggg	gaa	gtt	atc	gtg	acc	tta	gag	ccc	tgt	gag	cgc	tca	gaa	acc	tac	2004
Gly	Glu	Val	Ile	Val	Thr	Leu	Glu	Pro	Cys	Glu	Arg	Ser	Glu	Thr	Tyr	
				595					600						605	
gta	aat	ggc	aag	agg	gtg	tcc	cag	cct	gtt	cag	ctg	cgc	tca	gga	aac	2052
Val	Asn	Gly	Lys	Arg	Val	Ser	Gln	Pro	Val	Gln	Leu	Arg	Ser	Gly	Asn	
				610				615							620	
cgt	atc	atc	atg	ggc	aaa	aac	cat	gtt	ttc	cgc	ttt	aac	cac	ccg	gaa	2100
Arg	Ile	Met	Gly	Lys	Asn	His	Val	Phe	Arg	Phe	Asn	His	Pro	Glu		
				625			630								635	
caa	gca	cga	gct	gag	cga	gag	aag	act	cct	tct	gct	gag	acc	ccc	tct	2148
Gln	Ala	Arg	Ala	Glu	Arg	Glu	Lys	Thr	Pro	Ser	Ala	Glu	Thr	Pro	Ser	
				640		645									650	
gag	cct	gtg	gac	tgg	aca	ttt	gcc	cag	agg	gag	ctt	ctg	gaa	aaa	caa	2196
Glu	Pro	Val	Asp	Trp	Thr	Phe	Ala	Gln	Arg	Glu	Leu	Leu	Glu	Lys	Gln	
655					660					665					670	
gga	att	gat	atg	aaa	caa	gag	atg	gag	aaa	agg	cta	cag	gaa	atg	gag	2244
Gly	Ile	Asp	Met	Lys	Gln	Glu	Met	Glu	Lys	Arg	Leu	Gln	Glu	Met	Glu	
				675					680						685	
atc	tta	tac	aaa	aag	gag	aag	gaa	gaa	gca	gat	ctt	ctt	ttg	gag	cag	2292
Ile	Leu	Tyr	Lys	Lys	Glu	Lys	Glu	Glu	Ala	Asp	Leu	Leu	Leu	Glu	Gln	
				690					695						700	
cag	aga	ctg	gac	tat	gag	agt	aaa	ttg	cag	gcc	ttg	cag	aag	cag	gtt	2340
Gln	Arg	Leu	Asp	Tyr	Glu	Ser	Lys	Leu	Gln	Ala	Leu	Gln	Lys	Gln	Val	
				705			710								715	
gaa	acc	cga	tct	ctg	gct	gca	gaa	aca	act	gaa	gag	gag	gaa	gaa	gag	2388
Glu	Thr	Arg	Ser	Leu	Ala	Ala	Glu	Thr	Thr	Glu	Glu	Glu	Glu	Glu	Glu	
				720		725									730	
gaa	gaa	gtt	cct	tgg	aca	cag	cat	gaa	ttt	gag	ttg	gcc	caa	tgg	gcc	2436
Glu	Glu	Val	Pro	Trp	Thr	Gln	His	Glu	Phe	Glu	Leu	Ala	Gln	Trp	Ala	
735					740					745					750	
ttc	cgg	aaa	tgg	aag	tct	cat	cag	ttt	act	tca	tta	cgg	gac	tta	ctc	2484
Phe	Arg	Lys	Trp	Lys	Ser	His	Gln	Phe	Thr	Ser	Leu	Arg	Asp	Leu	Leu	
				755					760						765	
tgg	ggc	aat	gcc	gtg	tac	cta	aag	gag	gcc	aat	gcc	atc	agt	gtg	gaa	2532
Trp	Gly	Asn	Ala	Val	Tyr	Leu	Lys	Glu	Ala	Asn	Ala	Ile	Ser	Val	Glu	
				770				775							780	
ctg	aaa	aag	aag	gtg	cag	ttt	cag	ttt	gtt	ctg	ctg	act	gac	aca	ctg	2580
Leu	Lys	Lys	Lys	Val	Gln	Phe	Gln	Phe	Val	Leu	Leu	Thr	Asp	Thr	Leu	
				785			790								795	
tac	tcc	cct	ttg	cct	cct	gaa	tta	ctt	ccc	act	gag	atg	gaa	aaa	act	2628
Tyr	Ser	Pro	Leu	Pro	Pro	Glu	Leu	Leu	Pro	Thr	Glu	Met	Glu	Lys	Thr	
				800		805									810	
cat	gag	gac	agg	cct	ttc	cgc	aca	gtg	gta	gca	gta	gaa	gtc	cag		2676
His	Glu	Asp	Arg	Pro	Phe	Pro	Arg	Thr	Val	Ala	Val	Glu	Val	Gln		
815					820										830	
gat	ttg	aag	aat	gga	gca	aca	cac	tat	tgg	tct	ttg	gag	aaa	ctc	aag	2724

55

Asp	Leu	Lys	Asn	Gly	Ala	Thr	His	Tyr	Trp	Ser	Leu	Glu	Lys	Leu	Lys	
			835						840					845		
cag	agg	ctg	gat	ttg	atg	cga	gag	atg	tat	gat	agg	gca	ggg	gag	atg	2772
Gln	Arg	Leu	Asp	Leu	Met	Arg	Glu	Met	Tyr	Asp	Arg	Ala	Gly	Glu	Met	
			850					855					860			
gcc	tcc	agt	gcc	caa	gac	gaa	agc	gaa	acc	act	gtg	act	ggc	agc	gat	2820
Ala	Ser	Ser	Ala	Gln	Asp	Glu	Ser	Glu	Thr	Thr	Val	Thr	Gly	Ser	Asp	
			865				870					875				
ccc	ttc	tat	gat	cgg	ttc	cac	tgg	ttc	aaa	ctt	gtg	ggg	agc	tcc	ccc	2868
Pro	Phe	Tyr	Asp	Arg	Phe	His	Trp	Phe	Lys	Leu	Val	Gly	Ser	Ser	Pro	
			880			885					890					
att	ttc	cac	ggc	tgt	gtg	aac	gag	cgc	ctt	gcc	gac	cgc	aca	ccc	tcc	2916
Ile	Phe	His	Gly	Cys	Val	Asn	Glu	Arg	Leu	Ala	Asp	Arg	Thr	Pro	Ser	
			895		900				905					910		
ccc	act	ttt	tcc	acg	gcc	gat	tcc	gac	atc	act	gag	ctg	gct	gac	gag	2964
Pro	Thr	Phe	Ser	Thr	Ala	Asp	Ser	Asp	Ile	Thr	Glu	Leu	Ala	Asp	Glu	
				915				920					925			
cag	caa	gat	gag	atg	gag	gat	ttt	gat	gat	gag	gca	ttc	gtg	gat	gac	3012
Gln	Gln	Asp	Glu	Met	Glu	Asp	Phe	Asp	Asp	Glu	Ala	Phe	Val	Asp	Asp	
			930				935					940				
gcc	ggc	tct	gac	gca	ggg	acg	gag	gag	gga	tca	gat	ctc	ttc	agt	gac	3060
Ala	Gly	Ser	Asp	Ala	Gly	Thr	Glu	Glu	Gly	Ser	Asp	Leu	Phe	Ser	Asp	
			945			950					955					
ggg	cat	gac	ccg	ttt	tac	gac	cga	tcc	cct	tgg	ttc	att	tta	gtg	gga	3108
Gly	His	Asp	Pro	Phe	Tyr	Asp	Arg	Ser	Pro	Trp	Phe	Ile	Leu	Val	Gly	
			960			965				970						
agg	gca	ttt	gtt	tac	ctg	agc	aat	ctg	ctg	tat	ccc	gtg	ccc	ctg	atc	3156
Arg	Ala	Phe	Val	Tyr	Leu	Ser	Asn	Leu	Leu	Tyr	Pro	Val	Pro	Leu	Ile	
					980				985					990		
cac	agg	gtg	gcc	atc	gtc	agt	gag	aaa	ggg	gaa	gtg	cgg	gga	ttt	ctg	3204
His	Arg	Val	Ala	Ile	Val	Ser	Glu	Lys	Gly	Glu	Val	Arg	Gly	Phe	Leu	
				995				1000					1005			
cgt	gtg	gct	gta	cag	gcc	atc	gca	gcg	gat	gaa	gaa	gct	cct	gat	tat	3252
Arg	Val	Ala	Val	Gln	Ala	Ile	Ala	Ala	Asp	Glu	Glu	Ala	Pro	Asp	Tyr	
			1010				1015					1020				
ggc	tct	gga	att	cga	cag	tca	gga	aca	gct	aaa	ata	tct	ttt	gat	aat	3300
Gly	Ser	Gly	Ile	Arg	Gln	Ser	Gly	Thr	Ala	Lys	Ile	Ser	Phe	Asp	Asn	
			1025			1030					1035					
gaa	tac	ttt	aat	cag	agt	gac	ttt	tcg	tct	gtt	gca	atg	act	cgt	tct	3348
Glu	Tyr	Phe	Asn	Gln	Ser	Asp	Phe	Ser	Ser	Val	Ala	Met	Thr	Arg	Ser	
			1040			1045					1050					
ggg	ctg	tcc	ttg	gag	gag	ttg	agg	att	gtg	gaa	gga	cag	ggg	cag	agt	3396
Gly	Leu	Ser	Leu	Glu	Glu	Leu	Arg	Ile	Val	Glu	Gly	Gln	Gly	Gln	Ser	
				1055		1060			1065				1070			
tct	gag	gtc	atc	act	cct	cca	gaa	gaa	atc	agt	cga	att	aat	gac	ttg	3444
Ser	Glu	Val	Ile	Thr	Pro	Pro	Glu	Glu	Ile	Ser	Arg	Ile	Asn	Asp	Leu	
				1075				1080					1085			
gat	ttg	aag	tca	agc	act	ttg	ctg	gat	ggg	aag	atg	gta	atg	gaa	ggg	3492
Asp	Leu	Lys	Ser	Ser	Thr	Leu	Leu	Asp	Gly	Lys	Met	Val	Met	Glu	Gly	
			1090				1095					1100				
ttt	tct	gaa	gag	att	ggc	aac	cac	ctg	aaa	ctg	ggc	agt	gcc	ttc	act	3540
Phe	Ser	Glu	Glu	Ile	Gly	Asn	His	Leu	Lys	Leu	Gly	Ser	Ala	Phe	Thr	
			1105			1110					1115					
ttc	cga	gta	aca	gtg	ttg	cag	gcc	agt	gga	atc	ctc	cca	gag	tat	gca	3588
Phe	Arg	Val	Thr	Val	Leu	Gln	Ala	Ser	Gly	Ile	Leu	Pro	Glu	Tyr	Ala	
			1120			1125				1130						
gat	atc	ttc	tgt	cag	ttc	aac	ttt	ttg	cat	cgc	cat	gat	gaa	gca	ttc	3636
Asp	Ile	Phe	Cys	Gln	Phe	Asn	Phe	Leu	His	Arg	His	Asp	Glu	Ala	Phe	
				1135		1140			1145				1150			
tcc	acg	gag	ccc	ctc	aaa	aac	aat	ggc	aga	gga	agt	ccc	ctg	gcc	ttt	3684
Ser	Thr	Glu	Pro	Leu	Lys	Asn	Asn	Gly	Arg	Gly	Ser	Pro	Leu	Ala	Phe	
				1155				1160					1165			
tat	cat	gtg	cag	aat	att	gca	gtg	gag	atc	act	gaa	tca	ttt	gtg	gat	3732

56

Tyr	His	Val	Gln	Asn	Ile	Ala	Val	Glu	Ile	Thr	Glu	Ser	Phe	Val	Asp	
			1170					1175					1180			
tac	atc	aaa	acc	aag	cct	att	gta	ttt	gaa	gtc	ttt	ggg	cat	tat	cag	3780
Tyr	Ile	Lys	Thr	Lys	Pro	Ile	Val	Phe	Glu	Val	Phe	Gly	His	Tyr	Gln	
			1185					1190				1195				
cag	cac	cca	ctt	cat	ctg	caa	gga	cag	gag	ctt	aac	agt	ccg	cct	cag	3828
Gln	His	Pro	Leu	His	Leu	Gln	Gly	Gln	Glu	Leu	Asn	Ser	Pro	Pro	Gln	
			1200					1205				1210				
ccg	tgc	cgc	cga	ttc	ttc	cct	cca	ccc	atg	cca	ctg	tcc	aag	cca	gtt	3876
Pro	Cys	Arg	Arg	Phe	Phe	Pro	Pro	Pro	Met	Pro	Leu	Ser	Lys	Pro	Val	
			1215					1220				1225			1230	
cca	gcc	acc	aag	tta	aac	acg	atg	agc	aaa	acc	agc	ctt	ggc	cag	agc	3924
Pro	Ala	Thr	Lys	Leu	Asn	Thr	Met	Ser	Lys	Thr	Ser	Leu	Gly	Gln	Ser	
			1235					1240				1245				
atg	agc	aag	tat	gac	ctc	ctg	gtt	tgg	ttt	gag	atc	agt	gaa	ctg	gag	3972
Met	Ser	Lys	Tyr	Asp	Leu	Leu	Val	Trp	Phe	Glu	Ile	Ser	Glu	Leu	Glu	
			1250					1255				1260				
cct	aca	gga	gag	tat	atc	cca	gct	gtg	gtt	gac	cac	aca	gca	ggc	ttg	4020
Pro	Thr	Gly	Glu	Tyr	Ile	Pro	Ala	Val	Val	Asp	His	Thr	Ala	Gly	Leu	
			1265					1270				1275				
cct	tgc	cag	ggg	aca	ttt	ttg	ctt	cat	cag	ggc	atc	cag	cga	agg	atc	4068
Pro	Cys	Gln	Gly	Thr	Phe	Leu	Leu	His	Gln	Gly	Ile	Gln	Arg	Arg	Ile	
			1280					1285				1290				
aca	gtg	acc	att	atc	cat	gag	aag	ggg	agc	gag	ctc	cat	tgg	aaa	gat	4116
Thr	Val	Thr	Ile	Ile	His	Glu	Lys	Gly	Ser	Glu	Leu	His	Trp	Lys	Asp	
			1295				1300				1305				1310	
gtt	cgt	gaa	ctg	gtg	gta	ggt	cgt	att	cgg	aat	aag	cct	gag	gtg	gat	4164
Val	Arg	Glu	Leu	Val	Val	Gly	Arg	Ile	Arg	Asn	Lys	Pro	Glu	Val	Asp	
			1315					1320				1325				
gaa	gct	gca	gtt	gat	gcc	atc	ctc	tcc	cta	aat	att	att	tct	gcc	aag	4212
Glu	Ala	Ala	Val	Asp	Ala	Ile	Leu	Ser	Leu	Asn	Ile	Ile	Ser	Ala	Lys	
			1330					1335				1340				
tac	ctg	aag	tct	tcc	cac	aac	tct	agc	agg	acc	ttc	tac	cgc	ttt	gag	4260
Tyr	Leu	Lys	Ser	Ser	His	Asn	Ser	Ser	Arg	Thr	Phe	Tyr	Arg	Phe	Glu	
			1345					1350				1355				
gct	gtg	tgg	gat	agc	tct	ctg	cat	aac	tcc	ctt	ctt	ctg	aac	cga	gtg	4308
Ala	Val	Trp	Asp	Ser	Ser	Leu	His	Asn	Ser	Leu	Leu	Leu	Asn	Arg	Val	
			1360				1365					1370				
aca	ccc	tat	gga	gaa	aag	atc	tac	atg	acc	ttg	tgc	gcc	tac	cta	gag	4356
Thr	Pro	Tyr	Gly	Glu	Lys	Ile	Tyr	Met	Thr	Leu	Ser	Ala	Tyr	Leu	Glu	
			1375				1380				1385				1390	
ctg	gat	cat	tgc	atc	cag	ccg	gct	gtc	atc	acc	aag	gat	gtg	tgc	atg	4404
Leu	Asp	His	Cys	Ile	Gln	Pro	Ala	Val	Ile	Thr	Lys	Asp	Val	Cys	Met	
			1395					1400				1405				
gtc	ttc	tac	tcc	cga	gat	gcc	aag	atc	tca	cca	cca	cgc	tct	ctg	cgt	4452
Val	Phe	Tyr	Ser	Arg	Asp	Ala	Lys	Ile	Ser	Pro	Pro	Arg	Ser	Leu	Arg	
			1410					1415				1420				
agc	ctc	ttt	ggc	agc	ggc	tac	tca	aag	tca	cca	gat	tgc	aat	cga	gtc	4500
Ser	Leu	Phe	Gly	Ser	Gly	Tyr	Ser	Lys	Ser	Pro	Asp	Ser	Asn	Arg	Val	
			1425				1430				1435					
act	ggc	att	tac	gaa	ctc	agc	tta	tgc	aaa	atg	tca	gac	aca	ggt	agt	4548
Thr	Gly	Ile	Tyr	Glu	Leu	Ser	Leu	Cys	Lys	Met	Ser	Asp	Thr	Gly	Ser	
			1440				1445				1450					
cca	ggt	atg	cag	aga	agg	aga	aga	aaa	atc	tta	gat	acg	tca	gtg	gca	4596
Pro	Gly	Met	Gln	Arg	Arg	Arg	Lys	Ile	Leu	Asp	Thr	Ser	Val	Ala		
			1455				1460				1465				1470	
tat	gtg	cgg	gga	gaa	gag	aac	tta	gca	ggc	tgg	cgg	ccc	cgt	gga	gac	4644
Tyr	Val	Arg	Gly	Glu	Glu	Asn	Leu	Ala	Gly	Trp	Arg	Pro	Arg	Gly	Asp	
			1475					1480				1485				
agc	ctc	atc	ctt	gag	cac	cag	tgg	gag	ctg	gag	aag	ctg	gag	ctc	cta	4692
Ser	Leu	Ile	Glu	His	Gln	Trp	Glu	Leu	Glu	Lys	Leu	Glu	Leu	Leu		
			1490					1495				1500				
cat	gag	gtg	gaa	aaa	acc	cgc	cac	ttt	ttg	ctg	ctg	cgt	gag	aga	ctt	4740

57

His Glu Val Glu Lys Thr Arg His Phe Leu Leu Leu Arg Glu Arg Leu	
1505 1510 1515	
ggt gac agc atc ccc aaa tcc ctg agc gac tcg tta tcc ccc agc ctc	4788
Gly Asp Ser Ile Pro Lys Ser Leu Ser Asp Ser Leu Ser Pro Ser Leu	
1520 1525 1530	
agc agt ggg acc ctc agc acc tcc acc agt atc tcc tct cag atc tca	4836
Ser Ser Gly Thr Leu Ser Thr Ser Thr Ser Ile Ser Ser Gln Ile Ser	
1535 1540 1545 1550	
acc act acc ttt gaa agc gcc atc aca cct agc gag agc agt ggc tat	4884
Thr Thr Thr Phe Glu Ser Ala Ile Thr Pro Ser Glu Ser Ser Gly Tyr	
1555 1560 1565	
gat tca gga gac atc gaa agc ctg gtg gac cga gag aaa gag ctg gct	4932
Asp Ser Gly Asp Ile Glu Ser Leu Val Asp Arg Glu Lys Glu Leu Ala	
1570 1575 1580	
acc aag tgc ctg caa ctt ctc acc cac act ttc aac aga gaa ttc agc	4980
Thr Lys Cys Leu Gln Leu Leu Thr His Thr Phe Asn Arg Glu Phe Ser	
1585 1590 1595	
cag gtg cac ggc agc gtc agt gac tgt aag ttg tct gat atc tct cca	5028
Gln Val His Gly Ser Val Ser Asp Cys Lys Leu Ser Asp Ile Ser Pro	
1600 1605 1610	
att gga cgg gat ccc tct gag tcc agt ttc agc agt gcc acc ctc act	5076
Ile Gly Arg Asp Pro Ser Glu Ser Ser Phe Ser Ser Ala Thr Leu Thr	
1615 1620 1625 1630	
ccc tcc tcc acc tgt ccc tct ctg gta gac tct agg agc aac tct ctg	5124
Pro Ser Ser Thr Cys Pro Ser Leu Val Asp Ser Arg Ser Asn Ser Leu	
1635 1640 1645	
gat cag aag acc cca gaa gcc aat tcc cgg gcc tct agt ccc tgc cca	5172
Asp Gln Lys Thr Pro Glu Ala Asn Ser Arg Ala Ser Ser Pro Cys Pro	
1650 1655 1660	
gaa ttt gaa cag ttt cag att gtc cca gct gtg gaa aca cca tat ttg	5220
Glu Phe Glu Gln Phe Gln Ile Val Pro Ala Val Glu Thr Pro Tyr Leu	
1665 1670 1675	
gcc cga gca gga aaa aac gaa ttt ctc aat ctt gtt cca gat att gaa	5268
Ala Arg Ala Gly Lys Asn Glu Phe Leu Asn Leu Val Pro Asp Ile Glu	
1680 1685 1690	
gaa att aga cca agc tca gtg gtc tct aag aaa gga tac ctt cat ttc	5316
Glu Ile Arg Pro Ser Ser Val Val Ser Lys Lys Gly Tyr Leu His Phe	
1695 1700 1705 1710	
aag gag cct ctt tac agt aac tgg gct aaa cat ttt gtt gtc gtc cgt	5364
Lys Glu Pro Leu Tyr Ser Asn Trp Ala Lys His Phe Val Val Val Arg	
1715 1720 1725	
cgg cct tat gtc ttc atc tat aac agt gac aaa gac cct gtg gag cgt	5412
Arg Pro Tyr Val Phe Ile Tyr Asn Ser Asp Lys Asp Pro Val Glu Arg	
1730 1735 1740	
gga atc att aac ctg tcc aca gca cag gtg gag tac agt gag gac cag	5460
Gly Ile Ile Asn Leu Ser Thr Ala Gln Val Glu Tyr Ser Glu Asp Gln	
1745 1750 1755	
cag gcc atg gtg aag aca cca aac acm ttt gct gtc tgc aca aag cac	5508
Gln Ala Met Val Lys Thr Pro Asn Thr Phe Ala Val Cys Thr Lys His	
1760 1765 1770	
cgt ggg gtc ctt ttg cag gcc ctc aat gac aaa gac atg aac gac tgg	5556
Arg Gly Val Leu Leu Gln Ala Leu Asn Asp Lys Asp Met Asn Asp Trp	
1775 1780 1785 1790	
ttg tat gcc ttc aac cca ctt cta gct ggc aca ata cgg tca aag ctt	5604
Leu Tyr Ala Phe Asn Pro Leu Leu Ala Gly Thr Ile Arg Ser Lys Leu	
1795 1800 1805	
tcc cgc aga tgc ccg agc cag tcg aaa tac taagtgactc tgccgagtgc	5654
Ser Arg Arg Cys Pro Ser Gln Ser Lys Tyr	
1810 1815	
cctcactcgc cttcagagaga taaagaaagc gttacctctc atttctcttt gtgattcttg	5714
acgggtgactc ttgtatgtaa tcctgtggct taactacttc tccctccttg tccagcactt	5774
ttctagctct cccgttcccc atctccattg cctgtactc ttttcttttt tcttgtgctg	5834
agaatctcgt tagtagcatg tggcctaaca aaaggaaaaa atgttttttaa acacacacac	5894

acacacacac	acacacacac	acatacacan	acaaaaaac	aaaaactctg	aggggatctg	5954
gtgaatctcc	aaattattgt	gggtgtactt	tggcttcctt	ttgtatgata	gggtcccatc	6014
atgaccacct	ctgatgtctg	tgctgctgtc	accaggcacc	tttgtttttc	aagacaacat	6074
actttttttt	tcttttctct	gtttgtgata	tcactttaat	ttttcttggg	tggcttagag	6134
actaaggag	gagacatctg	gccttttttag	aacctgagag	gaaaaaaaaga	gtcttttttt	6194
ccctctgtc	tctttttgcc	atggctaate	cctgcatttc	cattcaggga	aaagggtgta	6254
gtgagcatag	mactgcaaca	gttatattct	gagtcaaaagt	tggggctttt	tacggcataa	6314
ttatggaatt	tttattttact	ggtagagagg	agacgagagg	ctttttcagt	gggcctggga	6374
cagtggctgc	tcttgacttt	gtgtgaagg	aaatgccaa	gatgcttctg	gtggacttca	6434
ggggacccca	gggttttgcc	gtgggcccgtg	atggcagcag	gcgggtggg	gctttagact	6494
cctcacagca	ggattcctgc	ccactgtttt	ttctctgttg	ggaggggaagc	tcttttctag	6554
gagtgtctca	gttctgtctt	tggcattagt	gatgggtggtg	gtacagttgg	aattagtgcc	6614
atgtcataca	caaagtgtcc	acaaggcggg	agtgtttcac	tttctggtga	taaacttgat	6674
ggctattgtt	atgattaaga	taatgccggg	caggccgggc	acagtggctc	acgcctgtaa	6734
tccaagcact	tggggaggcc	gaggcgggca	gatcacgaga	tcaggagttc	aagaccagcc	6794
tggccaatat	gatgaaaccc	cgtctctact	aaaaatacaa	aattagtccg	gtatggtggc	6854
acatgcctgt	aattccagct	gcttgggagc	ctgaggcagg	agaactgctt	gaaccagga	6914
ggcagagggt	gcagtgaacc	aagatcgccg	tattgcactc	cagcctgggt	gacagagcaa	6974
gactctgcct	cagaaaaaaa	aaaaaaaata	ataatgctgg	gtagtacact	tgtgattgtt	7034
acagctccct	ttgatcaaa	aaatatagct	ttcaggcata	aacctggaag	tctccctctg	7094
aatccagcag	ttgttttcat	tgtatgctgt	cagggtgaag	atgctttcag	tgatgtctc	7154
tatactcata	aataagcaaa	atgtggcagg	ctttgtcttc	tggatcccag	gattaaaaat	7214
aaccgtgacc	actactcaa	acaaaacaca	atatgcctag	gggcacggat	gaacgtccag	7274
ggagcccggg	ccccaggctt	tgttgccgtg	tccctgctcc	tctccatctg	gtgtggaaac	7334
actgcccagg	gagaaaagg	gaagctcact	gtggacagtc	ttctttcctt	ctgacagacc	7394
aggtcatctg	gcttccgaga	tcacacagaga	agataagtct	gtctctttca	gctgccagta	7454
agttttccag	gatgagagg	gaaaaagaaa	gcctccagtg	acttcagttg	ctttgccagt	7514
tgtcttgga	ttgttttaca	ccatccctta	cttcccttgc	tcagacctct	ctgtttcacc	7574
attgctcagg	cattcaggaa	agtatctgct	cactcccact	tgggtgagtcc	tcggccttga	7634
ggttgctgac	tctcaggcgt	taggcagctg	gatgacttcc	cgcttcatgc	agcaaaggcc	7694
aggggcttgc	gcgcctctgc	agagttgttg	ctaggagagac	ttgtgtcatc	atccacaacc	7754
ttgtttctca	cttccctggt	gggtccatct	ctgaagaaca	ggtctcccag	cttcgctcct	7814
tatcactgca	ttgtgaagag	gaggaaaagt	gaatcncgga	gagagaaaag	aaaggataga	7874
atcacaggct	gertctgcac	ctgaaaagt	accgcggaa	actctatggc	ggattttttt	7934
tttaactttc	ttcttccgtg	taaaacatag	gtcactaact	gtgatgttat	ttgttttcta	7994
agtgggtatgt	gagattttct	aatgtagtta	gaagtttcat	tgtctgatgg	acacaatatg	8054
cccttccggt	tctattcaaa	ccagcaggat	ctgtcgggtg	ttagagatgg	ctgcctggac	8114
tggaatcaaa	tctaatttca	gggaaatgaa	gatggaaattt	gaaggctcact	tttaaaatta	8174
agtcattgat	gctgctgtta	cagagtgtga	cagaggatcc	atgtctgtga	cacaggacgg	8234
tgggaagcct	gagagagagt	gaaattatgt	gatacactga	aatgactttt	gtttttcttc	8294
taactcatac	aaaactgggt	tggaaaagtct	ttgctttgga	agcgtcagac	attagaacag	8354
gccaactgg	actgtctgtt	catagcgtgc	ctgaataaga	aggcctctta	gggagccaga	8414
gggagcagag	tggctgtgtc	ctgcgtgtc	ttcaccctct	ggggcgcccc	tgtcgggct	8474
ggcagggtgca	gacagccttt	gctggtcccc	agcacgtcca	gggtgggtgc	tcccttgccc	8534
gacagaacca	tccccactgt	gaggctgtga	gagattttgtg	gcaggaaactg	tttatgaggc	8594
tctagttgtt	gctgttgttg	cgggaaagt	aagaaacata	gcccttaagg	aaaccacctt	8654
tatgtatttt	cttaaaagcac	gcctttaaat	aagcaaaaac	tttaaaaggc	aggaaagaga	8714
attcttaggc	aaattcagag	aaataagtgc	tagttaatac	taatcacctc	ctcctctgtc	8774
tctcatctc	ctttctccca	tcaaagcaaa	atatggcctc	accaccagcc	ccaaatcagt	8834
gctcagaccc	tctctgtgtc	tgtgtgccct	cctgggagtc	agtcagcgct	caggccagga	8894
ctgtgcagg	ccagccagcc	catgcgctag	tcaggagcac	aggcaagggg	tgcttgtggc	8954
agtggccggg	cacctgagcc	ccagcttgtt	gttaaacgtg	ctgacggcaa	ggggcaatgg	9014
agtgaatttc	ccaactaaga	aaccactatt	atatattttt	tcccttcagt	cacatagact	9074
tcagacaact	ctcctatttt	ttatggattt	ttcagctcat	ttcagatgaa	ggaactaagt	9134
cattgtgaac	tgtctcttga	gatctaaaaa	caagatgact	tttcttgcca	catattccaa	9194
agcaaagact	ttgttgccctg	ctgcttattg	tctaatttac	agggatattt	aattttgtca	9254
ggtctatgta	tatttatcca	gctatactta	cttgcacagt	ggattggaga	gaaaggattc	9314
tccagtgctg	acactcatcg	gtactctttc	tgcatttccc	tcgtgctgtg	tcccgtcgg	9374
gttccaatgg	acagtatcag	ggcttgtttg	acttaggtct	ttcagttttc	ctttcggttc	9434
ccttttaaaa	atgtgattgt	taacctgcct	cttgaagat	tcaaccgggt	gtggtggctc	9494
acgcctgtaa	tcccagcaca	tgggaggcc	aaggcagggtg	gatcacctga	ggtcaggagt	9554
ttgagaccag	cctggccaac	atggtgaaac	cccgtctcta	ctaaaaatac	aaaaagtagc	9614
caggcgtggt	ggcgtgtgcc	tgtagtccca	gctacttggg	aggctgaggc	aggagaatag	9674


```

cttgaacctg ggaggtggag gctgcagtga gttgagactg caccattgca ctccagcctg 9734
ggtgacaaag caggactccg tctcaaaaaa aaagaaaaga ttcgatgatgc tgctgctccc 9794
agaaggtttg ctggatgtgt ttacatagga ctctaacttg tgtgcactac agttgttcac 9854
cagggccagt gattcacccc agtgtgtggc cagaccatga ctgtgtagca ggaatgtttt 9914
aatttgtgct tccttagtaa attgaaatat cagctgagag attatttgct gctgttattc 9974
aaaaggccat ttatgaagtt agtatttgag ccccataaga tctttaaaaa gcctccaatc 10034
atttaaagga agaaatcaga gttgctataa aattcagtaa aaagctcata gccaaaacgg 10094
ctgtgctcag atggaaagtc tgagctgagg ttggtctctt gccaaaccgt ggctgttggtg 10154
tggtgttctt catgtcttcg agttcatttt ttttcattct gcctattctg gcatcagctc 10214
acttgaggag tccctcagcc ttcttgattt taaggcatcg tcttagactt tgtggctcta 10274
aagtacctgt ctgttgagat ttcaagtctc ttgtcaccat cctcacacat gacaacaaaa 10334
cccataatgc ataagtggcc tttttgaacc aagactttgc aaactgatct ctcccccggtg 10394
aaggagtgtg gcacattagc aacaatgtac attaatTTTg gatTTTcatt ttcatgtttt 10454
attttgtaaa tattatctga tgtttggagc ttgagtatac agactgtaaa tatagtctct 10514
gtatttgtag taattctgat tcttttgctg tatagcctta gatgtgcaat gcagacacta 10574
tctaactgtg tgtggtaacc ttgcgtcacg gagctgttag tgaacgaggt aaaaaataa 10634
aaggtacagc cagtgcacat aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 10682

```

<210> 4

<211> 1816

<212> PRT

<213> Homo sapiens

<400> 4

```

Met Ser Gly Ala Ser Val Lys Val Ala Val Arg Val Arg Pro Phe Asn
1          5          10          15
Ser Arg Glu Thr Ser Lys Glu Ser Lys Cys Ile Ile Gln Met Gln Gly
20          25          30
Asn Ser Thr Ser Ile Ile Asn Pro Lys Asn Pro Lys Glu Ala Pro Lys
35          40          45
Ser Phe Ser Phe Asp Tyr Ser Tyr Trp Ser His Thr Ser Pro Glu Asp
50          55          60
Pro Cys Phe Ala Ser Gln Asn Arg Val Tyr Asn Asp Ile Gly Lys Glu
65          70          75          80
Met Leu Leu His Ala Phe Glu Gly Tyr Asn Val Cys Ile Phe Ala Tyr
85          90          95
Gly Gln Thr Gly Ala Gly Lys Ser Tyr Thr Met Met Gly Lys Gln Glu
100          105          110
Glu Ser Gln Ala Gly Ile Ile Pro Gln Leu Cys Glu Glu Leu Phe Glu
115          120          125
Lys Ile Asn Asp Asn Cys Asn Glu Glu Met Ser Tyr Ser Val Glu Val
130          135          140
Ser Tyr Met Glu Ile Tyr Cys Glu Arg Val Arg Asp Leu Leu Asn Pro
145          150          155          160
Lys Asn Lys Gly Asn Leu Arg Val Arg Glu His Pro Leu Leu Gly Pro
165          170          175
Tyr Val Glu Asp Leu Ser Lys Leu Ala Val Thr Ser Tyr Thr Asp Ile
180          185          190
Ala Asp Leu Met Asp Ala Gly Asn Lys Ala Arg Thr Val Ala Ala Thr
195          200          205
Asn Met Asn Glu Thr Ser Ser Arg Ser His Ala Val Phe Thr Ile Val
210          215          220
Phe Thr Gln Lys Lys His Asp Asn Glu Thr Asn Leu Ser Thr Glu Lys
225          230          235          240
Val Ser Lys Ile Ser Leu Val Asp Leu Ala Gly Ser Glu Arg Ala Asp
245          250          255
Ser Thr Gly Ala Lys Gly Thr Arg Leu Lys Glu Gly Ala Asn Ile Asn
260          265          270
Lys Ser Leu Thr Thr Leu Gly Lys Val Ile Ser Ala Leu Ala Glu Val
275          280          285
Asp Asn Cys Thr Ser Lys Ser Lys Lys Lys Lys Thr Asp Phe Ile
290          295          300
Pro Tyr Arg Asp Ser Val Leu Thr Trp Leu Leu Arg Glu Asn Leu Gly

```

[illegible]

61

Asp Arg Pro Phe Pro Arg Thr Val Val Ala Val Glu Val Gln Asp Leu
 820 825 830
 Lys Asn Gly Ala Thr His Tyr Trp Ser Leu Glu Lys Leu Lys Gln Arg
 835 840 845
 Leu Asp Leu Met Arg Glu Met Tyr Asp Arg Ala Gly Glu Met Ala Ser
 850 855 860
 Ser Ala Gln Asp Glu Ser Glu Thr Thr Val Thr Gly Ser Asp Pro Phe
 865 870 875 880
 Tyr Asp Arg Phe His Trp Phe Lys Leu Val Gly Ser Ser Pro Ile Phe
 885 890 895
 His Gly Cys Val Asn Glu Arg Leu Ala Asp Arg Thr Pro Ser Pro Thr
 900 905 910
 Phe Ser Thr Ala Asp Ser Asp Ile Thr Glu Leu Ala Asp Glu Gln Gln
 915 920 925
 Asp Glu Met Glu Asp Phe Asp Asp Glu Ala Phe Val Asp Asp Ala Gly
 930 935 940
 Ser Asp Ala Gly Thr Glu Glu Gly Ser Asp Leu Phe Ser Asp Gly His
 945 950 955 960
 Asp Pro Phe Tyr Asp Arg Ser Pro Trp Phe Ile Leu Val Gly Arg Ala
 965 970 975
 Phe Val Tyr Leu Ser Asn Leu Leu Tyr Pro Val Pro Leu Ile His Arg
 980 985 990
 Val Ala Ile Val Ser Glu Lys Gly Glu Val Arg Gly Phe Leu Arg Val
 995 1000 1005
 Ala Val Gln Ala Ile Ala Ala Asp Glu Glu Ala Pro Asp Tyr Gly Ser
 1010 1015 1020
 Gly Ile Arg Gln Ser Gly Thr Ala Lys Ile Ser Phe Asp Asn Glu Tyr
 1025 1030 1035 1040
 Phe Asn Gln Ser Asp Phe Ser Ser Val Ala Met Thr Arg Ser Gly Leu
 1045 1050 1055
 Ser Leu Glu Glu Leu Arg Ile Val Glu Gly Gln Gly Gln Ser Ser Glu
 1060 1065 1070
 Val Ile Thr Pro Pro Glu Glu Ile Ser Arg Ile Asn Asp Leu Asp Leu
 1075 1080 1085
 Lys Ser Ser Thr Leu Leu Asp Gly Lys Met Val Met Glu Gly Phe Ser
 1090 1095 1100
 Glu Glu Ile Gly Asn His Leu Lys Leu Gly Ser Ala Phe Thr Phe Arg
 1105 1110 1115 1120
 Val Thr Val Leu Gln Ala Ser Gly Ile Leu Pro Glu Tyr Ala Asp Ile
 1125 1130 1135
 Phe Cys Gln Phe Asn Phe Leu His Arg His Asp Glu Ala Phe Ser Thr
 1140 1145 1150
 Glu Pro Leu Lys Asn Asn Gly Arg Gly Ser Pro Leu Ala Phe Tyr His
 1155 1160 1165
 Val Gln Asn Ile Ala Val Glu Ile Thr Glu Ser Phe Val Asp Tyr Ile
 1170 1175 1180
 Lys Thr Lys Pro Ile Val Phe Glu Val Phe Gly His Tyr Gln Gln His
 1185 1190 1195 1200
 Pro Leu His Leu Gln Gly Gln Glu Leu Asn Ser Pro Pro Gln Pro Cys
 1205 1210 1215
 Arg Arg Phe Phe Pro Pro Pro Met Pro Leu Ser Lys Pro Val Pro Ala
 1220 1225 1230
 Thr Lys Leu Asn Thr Met Ser Lys Thr Ser Leu Gly Gln Ser Met Ser
 1235 1240 1245
 Lys Tyr Asp Leu Leu Val Trp Phe Glu Ile Ser Glu Leu Glu Pro Thr
 1250 1255 1260
 Gly Glu Tyr Ile Pro Ala Val Val Asp His Thr Ala Gly Leu Pro Cys
 1265 1270 1275 1280
 Gln Gly Thr Phe Leu Leu His Gln Gly Ile Gln Arg Arg Ile Thr Val
 1285 1290 1295
 Thr Ile Ile His Glu Lys Gly Ser Glu Leu His Trp Lys Asp Val Arg
 1300 1305 1310
 Glu Leu Val Val Gly Arg Ile Arg Asn Lys Pro Glu Val Asp Glu Ala

1315 1320 1325
 Ala Val Asp Ala Ile Leu Ser Leu Asn Ile Ile Ser Ala Lys Tyr Leu
 1330 1335 1340
 Lys Ser Ser His Asn Ser Ser Arg Thr Phe Tyr Arg Phe Glu Ala Val
 1345 1350 1355 1360
 Trp Asp Ser Ser Leu His Asn Ser Leu Leu Asn Arg Val Thr Pro
 1365 1370 1375
 Tyr Gly Glu Lys Ile Tyr Met Thr Leu Ser Ala Tyr Leu Glu Leu Asp
 1380 1385 1390
 His Cys Ile Gln Pro Ala Val Ile Thr Lys Asp Val Cys Met Val Phe
 1395 1400 1405
 Tyr Ser Arg Asp Ala Lys Ile Ser Pro Pro Arg Ser Leu Arg Ser Leu
 1410 1415 1420
 Phe Gly Ser Gly Tyr Ser Lys Ser Pro Asp Ser Asn Arg Val Thr Gly
 1425 1430 1435 1440
 Ile Tyr Glu Leu Ser Leu Cys Lys Met Ser Asp Thr Gly Ser Pro Gly
 1445 1450 1455
 Met Gln Arg Arg Arg Arg Lys Ile Leu Asp Thr Ser Val Ala Tyr Val
 1460 1465 1470
 Arg Gly Glu Glu Asn Leu Ala Gly Trp Arg Pro Arg Gly Asp Ser Leu
 1475 1480 1485
 Ile Leu Glu His Gln Trp Glu Leu Glu Lys Leu Glu Leu His Glu
 1490 1495 1500
 Val Glu Lys Thr Arg His Phe Leu Leu Leu Arg Glu Arg Leu Gly Asp
 1505 1510 1515 1520
 Ser Ile Pro Lys Ser Leu Ser Asp Ser Leu Ser Pro Ser Leu Ser Ser
 1525 1530 1535
 Gly Thr Leu Ser Thr Ser Thr Ser Ile Ser Ser Gln Ile Ser Thr Thr
 1540 1545 1550
 Thr Phe Glu Ser Ala Ile Thr Pro Ser Glu Ser Ser Gly Tyr Asp Ser
 1555 1560 1565
 Gly Asp Ile Glu Ser Leu Val Asp Arg Glu Lys Glu Leu Ala Thr Lys
 1570 1575 1580
 Cys Leu Gln Leu Leu Thr His Thr Phe Asn Arg Glu Phe Ser Gln Val
 1585 1590 1595 1600
 His Gly Ser Val Ser Asp Cys Lys Leu Ser Asp Ile Ser Pro Ile Gly
 1605 1610 1615
 Arg Asp Pro Ser Glu Ser Ser Phe Ser Ser Ala Thr Leu Thr Pro Ser
 1620 1625 1630
 Ser Thr Cys Pro Ser Leu Val Asp Ser Arg Ser Asn Ser Leu Asp Gln
 1635 1640 1645
 Lys Thr Pro Glu Ala Asn Ser Arg Ala Ser Ser Pro Cys Pro Glu Phe
 1650 1655 1660
 Glu Gln Phe Gln Ile Val Pro Ala Val Glu Thr Pro Tyr Leu Ala Arg
 1665 1670 1675 1680
 Ala Gly Lys Asn Glu Phe Leu Asn Leu Val Pro Asp Ile Glu Glu Ile
 1685 1690 1695
 Arg Pro Ser Ser Val Val Ser Lys Lys Gly Tyr Leu His Phe Lys Glu
 1700 1705 1710
 Pro Leu Tyr Ser Asn Trp Ala Lys His Phe Val Val Val Arg Arg Pro
 1715 1720 1725
 Tyr Val Phe Ile Tyr Asn Ser Asp Lys Asp Pro Val Glu Arg Gly Ile
 1730 1735 1740
 Ile Asn Leu Ser Thr Ala Gln Val Glu Tyr Ser Glu Asp Gln Gln Ala
 1745 1750 1755 1760
 Met Val Lys Thr Pro Asn Thr Phe Ala Val Cys Thr Lys His Arg Gly
 1765 1770 1775
 Val L u Leu Gln Ala Leu Asn Asp Lys Asp Met Asn Asp Trp Leu Tyr
 1780 1785 1790
 Ala Phe Asn Pro Leu Leu Ala Gly Thr Ile Arg Ser Lys Leu Ser Arg
 1795 1800 1805
 Arg Cys Pro Ser Gln Ser Lys Tyr
 1810 1815

<210> 5
 <211> 357
 <212> DNA
 <213> Homo sapiens

<220>
 <221> allele
 <222> 178
 <223> 10-265-178 : polymorphic base A or G

<220>
 <221> misc_binding
 <222> 166..190
 <223> 10-265-178.probe

<220>
 <221> primer_bind
 <222> 159..177
 <223> 10-265-178.mis

<220>
 <221> primer_bind
 <222> 179..197
 <223> 10-265-178.mis complement

<220>
 <221> primer_bind
 <222> 1..18
 <223> 10-265.pu

<220>
 <221> primer_bind
 <222> 338..357
 <223> 10-265.rp complement

<400> 5
 ccaaatgtgg aaggaccgga cccctgggtt gcagcgcgtc gagcgggtgct gactctttcc 60
 ttgttctgt ttctgctct ctagagctga catcgcgctg atcggattgg ccgtcatggg 120
 ccagaactta attctgaaca tgaatgacca cggcttttg gtaagcggcg tgggcgcrtt 180
 gtcttctctc tggttcccgg gcgctttagc cgaggccggc gataggtttg ggagcttacg 240
 ggtctcctgg ccgtgctttg ctaatgtgct ctgttgctgc tcgtggcatt tttgtatgga 300
 aaggagaagc accctgtagg cgtgggcggc cgatcccga cttagtcctg cggagtg 357

<210> 6
 <211> 420
 <212> DNA
 <213> Homo sapiens

<220>
 <221> allele
 <222> 203
 <223> 10-266-203 : polymorphic base C or T

<220>
 <221> misc_binding
 <222> 191..215
 <223> 10-266-203.probe

<220>
 <221> primer_bind
 <222> 184..202
 <223> 10-266-203.mis

```

<220>
<221> primer_bind
<222> 204..222
<223> 10-266-203.mis complement

<220>
<221> primer_bind
<222> 1..20
<223> 10-266.pu

<220>
<221> primer_bind
<222> 401..420
<223> 10-266.rp complement

<400> 6
ggtgacataa ctttacagtg aagctccac agctgggga agaaaaggca aaggcaggtc      60
tgacctcccc agaacttggg ttgaatggaa aggtcattgt tactttggcc acagtctgaa      120
agtcttgtgt gtcctggatc tcctactcag gacttttgtc cttctagggtc tgtgtcttta      180
ataggactgt ctccaaagtt gaygatttct tggccaatga ggcaaaggga accaaagtgg      240
tgggtgcccc gtcctgaaa gagatgggtc ccaagctgaa gaagccccgg cggatcatcc      300
tcctggtgaa ggctgggcaa gctgtggatg atttcacga gaaattggtg aggccagctg      360
tgctctcagc tgctaccacg atagcagctg tttttggtt cttccttag ttctcttct      420

<210> 7
<211> 465
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> 118
<223> 12-592-118 : polymorphic base A or T

<220>
<221> misc_binding
<222> 106..130
<223> 12-592-118.probe

<220>
<221> primer_bind
<222> 99..117
<223> 12-592-118.mis

<220>
<221> primer_bind
<222> 119..137
<223> 12-592-118.mis complement

<220>
<221> primer_bind
<222> 1..19
<223> 12-592.pu

<220>
<221> primer_bind
<222> 448..465
<223> 12-592.rp complement

<400> 7
gagttcagtt tgagttcagc tacgagaatt atttagttga gtgagtgtca gagctgggat      60
tttcaaactc gcccttactg gtatgttgct ttacacaccc tcttacgtaa atcaaacwag      120

```

65

gattccaccc	tgggaggttt	gcatagaggg	ctgttcttgt	agaacttggtg	ctcatgcttt	180
gatttgggat	ttggggagtt	agggcaagcc	agaaagtgtt	tctggtggat	aataatgtgg	240
gttgactttc	ttaagcattt	taagccaagc	acttgagttt	ctaacaacta	aaaagctaag	300
tcagcctgac	acagctctag	cgcgccctgg	cttgattctg	ttcatcccca	gggggagact	360
tgcctttgtt	ccagtcctgc	tctccaagc	cagcttactg	tagttttcca	gcaattctga	420
gaagcagtat	tttttactgc	tgattagacc	ttaacatgga	aatgg		465

<210> 8

<211> 449

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> 420

<223> 12-783-421 : polymorphic base C or T

<220>

<221> misc_binding

<222> 408..432

<223> 12-783-421.probe

<220>

<221> primer_bind

<222> 401..419

<223> 12-783-421.mis

<220>

<221> primer_bind

<222> 421..439

<223> 12-783-421.mis complement

<220>

<221> primer_bind

<222> 1..19

<223> 12-783.pu

<220>

<221> primer_bind

<222> 429..449

<223> 12-783.rp complement

<220>

<221> allele

<222> 72

<223> 12-783-73 : polymorphic base G or C

<220>

<221> misc_binding

<222> 60..84

<223> 12-783-73.probe

<220>

<221> primer_bind

<222> 53..71

<223> 12-783-73.mis

<220>

<221> primer_bind

<222> 73..91

<223> 12-783-73.mis complement

<400> 8

66

atgactgtga	gaacgaaggg	gtaaactggg	acaagatccg	gtgattgcag	aaggcttctg	60
aggaggcgac	astgcacaga	gacctaagtg	aagtgaggag	cccttgctc	tgaggagaca	120
acacttctgg	tggccagaaa	ttcagtataa	ccaggttcag	aacaaacaca	actgactctg	180
ggttagcata	aaactcaacc	agcaggagca	gaagtcccca	ggcagacgcg	agcagagcct	240
gctggcaacc	gtgagcattg	gtcagcgtgg	acattggaca	aggggcttcc	ttgctcagcc	300
ctattcctc	taacatggtt	ctctcctgtg	ttctgcatgt	aggcctttga	ggattggaat	360
aagacagagc	tagactcatt	cctgattgaa	atcacagcca	atattctcaa	gttccaagay	420
accgatggca	aacacctgct	gccaaagat				449

<210> 9

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> sequencing oligonucleotide PrimerPU

<400> 9

tgtaaaacga cggccagt

18

<210> 10

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> sequencing oligonucleotide PrimerRP

<400> 10

caggaaacag ctatgacc

18

INTERNATIONAL SEARCH REPORT

International Application No
PCT/IB 00/00562

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/47 A01K67/027 C07K16/18 C12Q1/68
G06F17/30

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12N C07K A01K C12Q G06F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

STRAND, BIOSIS, EMBASE, CAB Data, EPO-Internal, PAJ, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Y. OKADA ET AL.: "The neuron-specific kinesin superfamily protein KIF1A is a unique monomeric motor for anterograde axonal transport of synaptic vesicle precursors." CELL, vol. 81, February 1906 (1906-02), pages 769-780, XP000929277 the whole document --- -/--	1-67

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"8" document member of the same patent family

Date of the actual completion of the international search

4 September 2000

Date of mailing of the international search report

08/09/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3018

Authorized officer

Hix, R

INTERNATIONAL SEARCH REPORT

national Application No
PCT/IB 00/00562

C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	M. NANGAKU ET AL.: "KIF1B, a novel microtubule plus end-directed monomeric motor protein for transport of mitochondria." CELL, vol. 79, 30 December 1994 (1994-12-30), pages 1209-1220, XP000915469 cited in the application the whole document	1-67
Y	R.A. FURLONG ET AL.: "Characterization of a kinesin-related gene ATSV, within the tuberous sclerosis locus (TSC1) candidate region on chromosome 9Q34." GENOMICS, vol. 33, no. 3, 1 May 1996 (1996-05-01), pages 421-429, XP000925509 the whole document	1-67
Y	AIZAWA H ET AL: "Kinesin family in murine central nervous system" JOURNAL OF CELL BIOLOGY, US, ROCKEFELLER UNIVERSITY PRESS, NEW YORK, US, vol. 119, 1 December 1992 (1992-12-01), pages 1287-1296, XP002080692 ISSN: 0021-9525 the whole document	1-67
P, X	T-W. L. GONG ET AL.: "A novel mouse kinesin of the UNC-104/KIF1 subfamily encoded by the Kif1b gene" GENE, vol. 239, 1999, pages 117-127, XP000925542 the whole document	17-22, 46, 47
A	Y. YONEKAWA ET AL.: "Defect in synaptic vesicle precursor transport and neuronal cell death in KIF1A motor protein-deficient mice." JOURNAL OF CELL BIOLOGY, vol. 141, no. 2, 20 April 1998 (1998-04-20), pages 431-441, XP000925196 cited in the application the whole document	
P, A	WO 00 18919 A (INCYTE PHARMA INC ; PATTERSON CHANDRA (US); CORLEY NEIL C (US); GUE) 6 April 2000 (2000-04-06) the whole document	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 00/00562

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0018919 A	06-04-2000	US 6013454 A	11-01-2000
		AU 6403599 A	17-04-2000